

**f –LACUNARY STATISTICAL CONVERGENCE AND STRONG
 f –LACUNARY SUMMABILITY OF ORDER α OF DOUBLE
SEQUENCES**

Hacer Şengül, Mikail Et and Yavuz Altın

© 2020 by University of Niš, Serbia | Creative Commons Licence: CC BY-NC-ND

Abstract. The main objective of this article is to introduce the concepts of f –lacunary statistical convergence of order α and strong f –lacunary summability of order α of double sequences and give some inclusion relations between these concepts.

Keywords: f –lacunary statistical convergence; strong f –lacunary summability; sequence spaces.

1. Introduction

In 1951, Steinhaus [41] and Fast [19] introduced the concept of statistical convergence while later in 1959, Schoenberg [40] reintroduced it independently. Bhardwaj and Dhawan [4], Caserta et al. [5], Connor [6], Çakallı [11], Çınar et al. [12], Çolak [13], Et et al. ([15],[17]), Fridy [21], Işık [27], Salat [39], Di Maio and Kočinac [14], Mursaleen et al. ([31],[30],[32]), Belen and Mohiuddine [3] and many authors investigated the arguments related to this notion.

A modulus f is a function from $[0, \infty)$ to $[0, \infty)$ such that

- i) $f(x) = 0$ if and only if $x = 0$,
- ii) $f(x + y) \leq f(x) + f(y)$ for $x, y \geq 0$,
- iii) f is increasing,
- iv) f is continuous from the right at 0.

It follows that f must be continuous everywhere on $[0, \infty)$. A modulus may be unbounded or bounded.

Aizpuru et al. [1] defined f –density of a subset $E \subset \mathbb{N}$ for any unbounded modulus f by

$$d^f(E) = \lim_{n \rightarrow \infty} \frac{f(|\{k \leq n : k \in E\}|)}{f(n)}, \text{ if the limit exists}$$

and defined f -statistical convergence for any unbounded modulus f by

$$d^f(\{k \in \mathbb{N} : |x_k - \ell| \geq \varepsilon\}) = 0$$

i.e.

$$\lim_{n \rightarrow \infty} \frac{1}{f(n)} f(|\{k \leq n : |x_k - \ell| \geq \varepsilon\}|) = 0,$$

and we write it as $S^f - \lim x_k = \ell$ or $x_k \rightarrow \ell (S^f)$. Every f -statistically convergent sequence is statistically convergent, but a statistically convergent sequence does not need to be f -statistically convergent for every unbounded modulus f .

By a lacunary sequence we mean an increasing integer sequence $\theta = (k_r)$ of non-negative integers such that $k_0 = 0$ and $h_r = (k_r - k_{r-1}) \rightarrow \infty$ as $r \rightarrow \infty$. The intervals determined by θ will be denoted by $I_r = (k_{r-1}, k_r]$ and the ratio $\frac{k_r}{k_{r-1}}$ will be abbreviated by q_r , and $q_1 = k_1$ for convenience.

In [22], Fridy and Orhan introduced the concept of lacunary statistically convergence in the sense that a sequence (x_k) of real numbers is called lacunary statistically convergent to a real number ℓ , if

$$\lim_{r \rightarrow \infty} \frac{1}{h_r} |\{k \in I_r : |x_k - \ell| \geq \varepsilon\}| = 0$$

for every positive real number ε .

Lacunary sequence spaces were studied in ([7],[8],[9],[10],[18],[20],[22],[23],[25],[26],[28],[36],[43]).

A double sequence $x = (x_{j,k})_{j,k=0}^{\infty}$ has Pringsheim limit ℓ provided that given for every $\varepsilon > 0$ there exists $N \in \mathbb{N}$ such that $|x_{j,k} - \ell| < \varepsilon$ whenever $j, k > N$. In this case, we write $P - \lim x = \ell$ (see Pringsheim [38]).

Let $K \subseteq \mathbb{N} \times \mathbb{N}$ and $K(m, n) = \{(j, k) : j \leq m, k \leq n\}$. The double natural density of K is defined by

$$\delta_2(K) = P - \lim_{m,n} \frac{1}{mn} |K(m, n)|, \text{ if the limit exists.}$$

A double sequence $x = (x_{jk})_{j,k \in \mathbb{N}}$ is said to be statistically convergent to a number ℓ if for every $\varepsilon > 0$ the set $\{(j, k) : j \leq m, k \leq n : |x_{jk} - \ell| \geq \varepsilon\}$ has double natural density zero (see Mursaleen and Edely [31]).

In [35], Patterson and Savaş introduced the concept of double lacunary sequence in the sense that double sequence $\theta'' = \{(k_r, l_s)\}$ is called double lacunary sequence, if there exists two increasing sequences of integers such that

$$k_0 = 0, h_r = k_r - k_{r-1} \rightarrow \infty \text{ as } r \rightarrow \infty$$

and

$$l_0 = 0, \bar{h}_s = l_s - l_{s-1} \rightarrow \infty \text{ as } s \rightarrow \infty.$$

where $k_{r,s} = k_r l_s$, $h_{r,s} = h_r \bar{h}_s$ and the following intervals are determined by θ'' , $I_r = \{(k) : k_{r-1} < k \leq k_r\}$, $I_s = \{(l) : l_{s-1} < l \leq l_s\}$, $I_{r,s} = \{(k, l) : k_{r-1} < k \leq k_r \text{ and } l_{s-1} < l \leq l_s\}$, $q_r = \frac{k_r}{k_{r-1}}$, $\bar{q}_s = \frac{l_s}{l_{s-1}}$ and $q_{r,s} = q_r \bar{q}_s$.

The double number sequence x is $S_{\theta''}$ -convergent to ℓ provided that for every $\varepsilon > 0$,

$$P - \lim_{r,s} \frac{1}{h_{r,s}} |\{(k, l) \in I_{r,s} : |x_{k,l} - \ell| \geq \varepsilon\}| = 0.$$

In this case, we write $S_{\theta''} - \lim x_{k,l} = \ell$ or $x_{k,l} \rightarrow \ell (S_{\theta''})$ (see [35]).

The notion of a modulus was given by Nakano [33]. Maddox [29] used a modulus function to construct some sequence spaces. Afterwards, different sequence spaces defined by modulus have been studied by Altın and Et [2], Et et al. [16], Işık [27], Gaur and Mursaleen [24], Nuray and Savaş [34], Pehlivan and Fisher [37], Şengül [42] and many others.

2. Main Results

In this section, we will introduce the concepts of f -lacunary statistical convergence of order α and strong f -lacunary summability of order α of double sequences, where f is an unbounded modulus and also give some results related to these concepts.

Definition 2.1. Let f be an unbounded modulus, $\theta'' = \{(k_r, l_s)\}$ be a double lacunary sequence and α be a real number such that $0 < \alpha \leq 1$. We say that the double sequence $x = (x_{k,l})$ is f -lacunary statistically convergent of order α , if there is a real number ℓ such that

$$\lim_{r,s \rightarrow \infty} \frac{1}{[f(h_{r,s})]^\alpha} f(|\{(k, l) \in I_{r,s} : |x_{k,l} - \ell| \geq \varepsilon\}|) = 0.$$

This space will be denoted by $S_{\theta''}^{f,\alpha}$. In this case, we write $S_{\theta''}^{f,\alpha} - \lim x_{k,l} = \ell$ or $x_{k,l} \rightarrow \ell (S_{\theta''}^{f,\alpha})$. In the special case $\theta'' = \{(2^r, 2^s)\}$, we shall write $S''^{f,\alpha}$ instead of $S_{\theta''}^{f,\alpha}$.

Definition 2.2. Let f be a modulus function, $\theta'' = \{(k_r, l_s)\}$ be a double lacunary sequence, $p = (p_k)$ be a sequence of strictly positive real numbers and α be a positive real number. We say that the double sequence $x = (x_{k,l})$ is strongly $w^\alpha [\theta'', f, p]$ -summable to ℓ (a real number), if there is a real number ℓ such that

$$\lim_{r,s \rightarrow \infty} \frac{1}{[h_{r,s}]^\alpha} \sum_{(k,l) \in I_{r,s}} [f(|x_{k,l} - \ell|)]^{p_k} = 0.$$

In this case we write $w^\alpha [\theta'', f, p] - \lim x_{k,l} = \ell$. The set of all strongly $w^\alpha [\theta'', f, p]$ -summable sequences will be denoted by $w^\alpha [\theta'', f, p]$. If we take $p_k = 1$ for all $k \in \mathbb{N}$, we write $w^\alpha [\theta'', f]$ instead of $w^\alpha [\theta'', f, p]$.

Definition 2.3. Let f be an unbounded modulus, $\theta'' = \{(k_r, l_s)\}$ be a double lacunary sequence, $p = (p_k)$ be a sequence of strictly positive real numbers and α be a positive real number. We say that the double sequence $x = (x_{k,l})$ is strongly $w_{\theta''}^{f,\alpha}(p)$ -summable to ℓ (a real number), if there is a real number ℓ such that

$$\lim_{r,s \rightarrow \infty} \frac{1}{[f(h_{r,s})]^\alpha} \sum_{(k,l) \in I_{r,s}} [f(|x_{k,l} - \ell|)]^{p_k} = 0.$$

In the present case, we write $w_{\theta''}^{f,\alpha}(p) - \lim x_{k,l} = \ell$. The set of all strongly $w_{\theta''}^{f,\alpha}(p)$ -summable sequences will be denoted by $w_{\theta''}^{f,\alpha}(p)$. In case of $p_k = p$ for all $k \in \mathbb{N}$ we write $w_{\theta''}^{f,\alpha}[p]$ instead of $w_{\theta''}^{f,\alpha}(p)$.

Definition 2.4. Let f be an unbounded modulus, $\theta'' = \{(k_r, l_s)\}$ be a double lacunary sequence, $p = (p_k)$ be a sequence of strictly positive real numbers and α be a positive real number. We say that the double sequence $x = (x_{k,l})$ is strongly $w_{\theta'',f}^\alpha(p)$ -summable to ℓ (a real number), if there is a real number ℓ such that

$$\frac{1}{[f(h_{r,s})]^\alpha} \sum_{(k,l) \in I_{r,s}} |x_{k,l} - \ell|^{p_k} = 0.$$

In the present case, we write $w_{\theta'',f}^\alpha(p) - \lim x_{k,l} = \ell$. The set of all strongly $w_{\theta'',f}^\alpha(p)$ -summable sequences will be denoted by $w_{\theta'',f}^\alpha(p)$. In case of $p_k = p$ for all $k \in \mathbb{N}$ we write $w_{\theta'',f}^\alpha[p]$ instead of $w_{\theta'',f}^\alpha(p)$.

The proof of each of the following results is fairly straightforward, so we choose to state these results without proof, where we shall assume that the sequence $p = (p_k)$ is bounded and $0 < h = \inf_k p_k \leq p_k \leq \sup_k p_k = H < \infty$.

Theorem 2.1. *The space $w_{\theta''}^{f,\alpha}(p)$ is paranormed by*

$$g(x) = \sup_{r,s} \left\{ \frac{1}{[f(h_{r,s})]^\alpha} \sum_{(k,l) \in I_{r,s}} [f(|x_{k,l}|)]^{p_k} \right\}^{\frac{1}{M}}$$

where, $M = \max(1, H)$.

Proposition 2.1. ([37]) *Let f be a modulus and $0 < \delta < 1$. Then for each $\|u\| \geq \delta$, we have $f(\|u\|) \leq 2f(1)\delta^{-1}\|u\|$.*

Theorem 2.2. Let f be an unbounded modulus, α be a real number such that $0 < \alpha \leq 1$ and $p > 1$. If $\lim_{u \rightarrow \infty} \inf \frac{f(u)}{u} > 0$, then $w_{\theta'',f}^{f,\alpha} [p] = w_{\theta'',f}^\alpha [p]$.

Proof. Let $p > 1$ be a positive real number and $x \in w_{\theta'',f}^{f,\alpha} [p]$. If $\lim_{u \rightarrow \infty} \inf \frac{f(u)}{u} > 0$ then there exists a number $c > 0$ such that $f(u) > cu$ for $u > 0$. Clearly

$$\begin{aligned} \frac{1}{[f(h_{r,s})]^\alpha} \sum_{(k,l) \in I_{r,s}} [f(|x_{k,l} - \ell|)]^p &\geq \frac{1}{[f(h_{r,s})]^\alpha} \sum_{(k,l) \in I_{r,s}} [c|x_{k,l} - \ell|]^p \\ &= \frac{c^p}{[f(h_{r,s})]^\alpha} \sum_{(k,l) \in I_{r,s}} |x_{k,l} - \ell|^p, \end{aligned}$$

and therefore $w_{\theta'',f}^{f,\alpha} [p] \subset w_{\theta'',f}^\alpha [p]$.

Now let $x \in w_{\theta'',f}^\alpha [p]$. Then we have

$$\frac{1}{[f(h_{r,s})]^\alpha} \sum_{(k,l) \in I_{r,s}} |x_{k,l} - \ell|^p \rightarrow 0 \text{ as } r, s \rightarrow \infty.$$

Let $0 < \delta < 1$. We can write

$$\begin{aligned} \frac{1}{[f(h_{r,s})]^\alpha} \sum_{(k,l) \in I_{r,s}} |x_{k,l} - \ell|^p &\geq \frac{1}{[f(h_{r,s})]^\alpha} \sum_{\substack{(k,l) \in I_{r,s} \\ |x_{k,l} - \ell| \geq \delta}} |x_{k,l} - \ell|^p \\ &\geq \frac{1}{[f(h_{r,s})]^\alpha} \sum_{\substack{(k,l) \in I_{r,s} \\ |x_{k,l} - \ell| \geq \delta}} \left[\frac{f(|x_{k,l} - \ell|)}{2f(1)\delta^{-1}} \right]^p \\ &\geq \frac{1}{[f(h_{r,s})]^\alpha} \frac{\delta^p}{2^p f(1)^p} \sum_{(k,l) \in I_{r,s}} [f(|x_{k,l} - \ell|)]^p \end{aligned}$$

by Proposition 2.1. Therefore $x \in w_{\theta'',f}^{f,\alpha} [p]$.

If $\lim_{u \rightarrow \infty} \inf \frac{f(u)}{u} = 0$, the equality $w_{\theta'',f}^{f,\alpha} [p] = w_{\theta'',f}^\alpha [p]$ cannot be hold as shown in the following example:

Let $f(x) = 2\sqrt{x}$ and define a double sequence $x = (x_{k,l})$ by

$$x_{k,l} = \begin{cases} \sqrt[3]{h_{r,s}}, & \text{if } k = k_r \text{ and } l = l_s \\ 0, & \text{otherwise} \end{cases} \quad r, s = 1, 2, \dots$$

For $\ell = 0$, $\alpha = \frac{3}{4}$ and $p = \frac{6}{5}$, we have

$$\frac{1}{[f(h_{r,s})]^\alpha} \sum_{(k,l) \in I_{r,s}} [f(|x_{k,l}|)]^p = \frac{(2[h_{r,s}]^{\frac{1}{6}})^{\frac{6}{5}}}{(2\sqrt{h_{r,s}})^{\frac{3}{4}}} = \frac{(2(h_r \bar{h}_s)^{\frac{1}{6}})^{\frac{6}{5}}}{(2\sqrt{h_r \bar{h}_s})^{\frac{3}{4}}} \rightarrow 0 \text{ as } r, s \rightarrow \infty$$

hence $x \in w_{\theta''}^{f,\alpha} [p]$, but

$$\frac{1}{[f(h_{r,s})]^\alpha} \sum_{(k,l) \in I_{r,s}} |x_{k,l}|^p = \frac{(\sqrt[3]{h_{r,s}})^{\frac{6}{5}}}{(2\sqrt{h_{r,s}})^{\frac{3}{4}}} \rightarrow \infty \text{ as } r, s \rightarrow \infty$$

and so $x \notin w_{\theta'',f}^{\alpha} [p]$. \square

Maddox [29] showed that the existence of an unbounded modulus f for which there is a positive constant c such that $f(xy) \geq cf(x)f(y)$, for all $x \geq 0, y \geq 0$.

Theorem 2.3. *Let f be an unbounded modulus and α be a positive real number. If $\lim_{u \rightarrow \infty} \frac{[f(u)]^\alpha}{u^\alpha} > 0$, then $w^\alpha [\theta'', f] \subset S_{\theta''}^{f,\alpha}$.*

Proof. Let $x \in w^\alpha [\theta'', f]$ and $\lim_{u \rightarrow \infty} \frac{f(u)^\alpha}{u^\alpha} > 0$. For $\varepsilon > 0$, we have

$$\begin{aligned} \frac{1}{[h_{r,s}]^\alpha} \sum_{(k,l) \in I_{r,s}} f(|x_{k,l} - \ell|) &\geq \frac{1}{[h_{r,s}]^\alpha} f\left(\sum_{(k,l) \in I_{r,s}} |x_{k,l} - \ell|\right) \geq \frac{1}{[h_{r,s}]^\alpha} f\left(\sum_{\substack{(k,l) \in I_{r,s} \\ |x_{k,l} - \ell| \geq \varepsilon}} |x_{k,l} - \ell|\right) \\ &\geq \frac{1}{[h_{r,s}]^\alpha} f(|\{(k,l) \in I_{r,s} : |x_{k,l} - \ell| \geq \varepsilon\}| \varepsilon) \\ &\geq \frac{c}{[h_{r,s}]^\alpha} f(|\{(k,l) \in I_{r,s} : |x_{k,l} - \ell| \geq \varepsilon\}|) f(\varepsilon) \\ &= \frac{c}{[h_{r,s}]^\alpha} \frac{f(|\{(k,l) \in I_{r,s} : |x_{k,l} - \ell| \geq \varepsilon\}|)}{[f(h_{r,s})]^\alpha} [f(h_{r,s})]^\alpha f(\varepsilon). \end{aligned}$$

Therefore, $w^\alpha [\theta'', f] - \lim x_{k,l} = \ell$ implies $S_{\theta''}^{f,\alpha} - \lim x_{k,l} = \ell$. \square

Theorem 2.4. *Let α_1, α_2 be two real numbers such that $0 < \alpha_1 \leq \alpha_2 \leq 1$, f be an unbounded modulus function and let $\theta'' = \{(k_r, l_s)\}$ be a double lacunary sequence, then we have $w_{\theta''}^{f,\alpha_1} (p) \subset S_{\theta''}^{f,\alpha_2}$.*

Proof. Let $x \in w_{\theta''}^{f,\alpha_1} (p)$ and $\varepsilon > 0$ be given and \sum_1, \sum_2 denote the sums over $(k, l) \in I_{r,s}, |x_{k,l} - \ell| \geq \varepsilon$ and $(k, l) \in I_{r,s}, |x_{k,l} - \ell| < \varepsilon$ respectively. Since

$f(h_{r,s})^{\alpha_1} \leq f(h_{r,s})^{\alpha_2}$ for each r and s , we may write

$$\begin{aligned} & \frac{1}{[f(h_{r,s})]^{\alpha_1}} \sum_{(k,l) \in I_{r,s}} [f(|x_{k,l} - \ell|)]^{p_k} \\ &= \frac{1}{[f(h_{r,s})]^{\alpha_1}} \left[\sum_1 [f(|x_{k,l} - \ell|)]^{p_k} + \sum_2 [f(|x_{k,l} - \ell|)]^{p_k} \right] \\ &\geq \frac{1}{[f(h_{r,s})]^{\alpha_2}} \left[\sum_1 [f(|x_{k,l} - \ell|)]^{p_k} + \sum_2 [f(|x_{k,l} - \ell|)]^{p_k} \right] \\ &\geq \frac{1}{[f(h_{r,s})]^{\alpha_2}} \left[\sum_1 [f(\varepsilon)]^{p_k} \right] \\ &\geq \frac{1}{H \cdot [f(h_{r,s})]^{\alpha_2}} \left[f \left(\sum_1 [\varepsilon]^{p_k} \right) \right] \\ &\geq \frac{1}{H \cdot [f(h_{r,s})]^{\alpha_2}} \left[f \left(\sum_1 \min([\varepsilon]^h, [\varepsilon]^H) \right) \right] \\ &\geq \frac{1}{H \cdot [f(h_{r,s})]^{\alpha_2}} f \left(|\{(k, l) \in I_{r,s} : |x_{k,l} - \ell| \geq \varepsilon\}| \left[\min([\varepsilon]^h, [\varepsilon]^H) \right] \right) \\ &\geq \frac{c}{H \cdot [f(h_{r,s})]^{\alpha_2}} f \left(|\{(k, l) \in I_{r,s} : |x_{k,l} - \ell| \geq \varepsilon\}| \right) f \left(\left[\min([\varepsilon]^h, [\varepsilon]^H) \right] \right). \end{aligned}$$

Hence $x \in S_{\theta''}^{f,\alpha_2}$. \square

Theorem 2.5. Let $\theta'' = \{(k_r, l_s)\}$ be a double lacunary sequence and α be a fixed real number such that $0 < \alpha \leq 1$. If $\liminf_r q_r > 1$, $\liminf_s q_s > 1$ and $\lim_{u \rightarrow \infty} \frac{[f(u)]^\alpha}{u^\alpha} > 0$, then $S''f,\alpha \subset S_{\theta''}^{f,\alpha}$.

Proof. Suppose first that $\liminf_r q_r > 1$ and $\liminf_s q_s > 1$; then there exists $a, b > 0$ such that $q_r \geq 1 + a$ and $q_s \geq 1 + b$ for sufficiently large r and s , which implies that

$$\frac{h_r}{k_r} \geq \frac{a}{1+a} \implies \left(\frac{h_r}{k_r} \right)^\alpha \geq \left(\frac{a}{1+a} \right)^\alpha$$

and

$$\frac{\bar{h}_s}{l_s} \geq \frac{b}{1+b} \implies \left(\frac{\bar{h}_s}{l_s} \right)^\alpha \geq \left(\frac{b}{1+b} \right)^\alpha.$$

If $S''f,\alpha - \lim x_{k,l} = \ell$, then for every $\varepsilon > 0$ and for sufficiently large r and s , we

have

$$\begin{aligned}
& \frac{1}{[f(k_r l_s)]^\alpha} f(|\{k \leq k_r, l \leq l_s : |x_{k,l} - \ell| \geq \varepsilon\}|) \\
& \geq \frac{1}{[f(k_r l_s)]^\alpha} f(|\{(k, l) \in I_{r,s} : |x_{k,l} - \ell| \geq \varepsilon\}|) \\
& = \frac{[f(h_{r,s})]^\alpha}{[f(k_r l_s)]^\alpha} \frac{1}{[f(h_{r,s})]^\alpha} f(|\{(k, l) \in I_{r,s} : |x_{k,l} - \ell| \geq \varepsilon\}|) \\
& = \frac{[f(h_{r,s})]^\alpha}{[h_{r,s}]^\alpha} \frac{k_r^\alpha}{[f(k_r l_s)]^\alpha} \frac{[h_{r,s}]^\alpha}{k_r^\alpha} \frac{f(|\{(k, l) \in I_{r,s} : |x_{k,l} - \ell| \geq \varepsilon\}|)}{[f(h_{r,s})]^\alpha} \\
& = \frac{[f(h_{r,s})]^\alpha}{[h_{r,s}]^\alpha} \frac{k_r^\alpha l_s^\alpha}{[f(k_r l_s)]^\alpha} \frac{h_r^\alpha \bar{h}_s^\alpha}{k_r^\alpha l_s^\alpha} f(|\{(k, l) \in I_{r,s} : |x_{k,l} - \ell| \geq \varepsilon\}|) \\
& \geq \frac{[f(h_{r,s})]^\alpha}{[h_{r,s}]^\alpha} \frac{(k_r l_s)^\alpha}{[f(k_r l_s)]^\alpha} \left(\frac{a}{1+a}\right)^\alpha \left(\frac{b}{1+b}\right)^\alpha \frac{f(|\{(k, l) \in I_{r,s} : |x_{k,l} - \ell| \geq \varepsilon\}|)}{[f(h_{r,s})]^\alpha}.
\end{aligned}$$

This proves the sufficiency. \square

Theorem 2.6. Let f be an unbounded modulus, $\theta = (k_r)$ and $\theta' = (l_s)$ be two lacunary sequences, $\theta'' = \{(k_r, l_s)\}$ be a double lacunary sequence and $0 < \alpha \leq 1$. If $S_{f,\theta}^\alpha - \lim x_k = \ell$ and $S_{f,\theta'}^\alpha - \lim x_l = \ell$, then $S_{f,\theta''}^\alpha - \lim x_{k,l} = \ell$.

Proof. Suppose $S_{f,\theta}^\alpha - \lim x_k = \ell$ and $S_{f,\theta'}^\alpha - \lim x_l = \ell$. Then for $\varepsilon > 0$ we can write

$$\lim_r \frac{1}{[f(h_r)]^\alpha} |\{k \in I_r : |x_k - \ell| \geq \varepsilon\}| = 0$$

and

$$\lim_s \frac{1}{[f(\bar{h}_s)]^\alpha} |\{l \in I_s : |x_l - \ell| \geq \varepsilon\}| = 0.$$

So we have

$$\begin{aligned}
& \frac{1}{[f(h_{r,s})]^\alpha} |\{(k, l) \in I_{r,s} : |x_{k,l} - \ell| \geq \varepsilon\}| \\
& \leq \frac{1}{[cf(h_r) f(\bar{h}_s)]^\alpha} |\{(k, l) \in I_{r,s} : |x_{k,l} - \ell| \geq \varepsilon\}| \\
& \leq \frac{1}{c^\alpha [f(h_r)]^\alpha [f(\bar{h}_s)]^\alpha} |\{(k, l) \in I_{r,s} : |x_{k,l} - \ell| \geq \varepsilon\}| \\
& \leq \left[\frac{1}{[f(h_r)]^\alpha} |\{k \in I_r : |x_k - \ell| \geq \varepsilon\}| \right] \left[\frac{1}{[f(\bar{h}_s)]^\alpha} |\{l \in I_s : |x_l - \ell| \geq \varepsilon\}| \right].
\end{aligned}$$

Hence $S_{f,\theta''}^\alpha - \lim x_{k,l} = \ell$. \square

Theorem 2.7. *Let f be an unbounded modulus. If $\lim p_k > 0$, then $w_{\theta''}^{f,\alpha}(p) - \lim x_{k,l} = \ell$ uniquely.*

Proof. Let $\lim p_k = s > 0$. Assume that $w_{\theta''}^{f,\alpha}(p) - \lim x_{k,l} = \ell_1$ and $w_{\theta''}^{f,\alpha}(p) - \lim x_{k,l} = \ell_2$. Then

$$\lim_{r,s} \frac{1}{[f(h_{r,s})]^\alpha} \sum_{(k,l) \in I_{r,s}} [f(|x_{k,l} - \ell_1|)]^{p_k} = 0,$$

and

$$\lim_{r,s} \frac{1}{[f(h_{r,s})]^\alpha} \sum_{(k,l) \in I_{r,s}} [f(|x_{k,l} - \ell_2|)]^{p_k} = 0.$$

By definition of f , we have

$$\begin{aligned} & \frac{1}{[f(h_{r,s})]^\alpha} \sum_{(k,l) \in I_{r,s}} [f(|\ell_1 - \ell_2|)]^{p_k} \\ & \leq \frac{D}{[f(h_{r,s})]^\alpha} \left(\sum_{(k,l) \in I_{r,s}} [f(|x_{k,l} - \ell_1|)]^{p_k} + \sum_{(k,l) \in I_{r,s}} [f(|x_{k,l} - \ell_2|)]^{p_k} \right) \\ & = \frac{D}{[f(h_{r,s})]^\alpha} \sum_{(k,l) \in I_{r,s}} [f(|x_{k,l} - \ell_1|)]^{p_k} + \frac{D}{[f(h_{r,s})]^\alpha} \sum_{(k,l) \in I_{r,s}} [f(|x_{k,l} - \ell_2|)]^{p_k} \end{aligned}$$

where $\sup_k p_k = H$ and $D = \max(1, 2^{H-1})$. Hence

$$\lim_{r,s} \frac{1}{[f(h_{r,s})]^\alpha} \sum_{(k,l) \in I_{r,s}} [f(|\ell_1 - \ell_2|)]^{p_k} = 0.$$

Since $\lim_{k \rightarrow \infty} p_k = s$ we have $\ell_1 - \ell_2 = 0$. Thus the limit is unique. \square

Theorem 2.8. *Let $\theta''_1 = \{(k_r, l_s)\}$ and $\theta''_2 = \{(s_r, t_s)\}$ be two double lacunary sequences such that $I_{r,s} \subset J_{r,s}$ for all $r, s \in \mathbb{N}$ and α_1, α_2 two real numbers such that $0 < \alpha_1 \leq \alpha_2 \leq 1$. If*

$$(2.1) \quad \lim_{r,s \rightarrow \infty} \inf \frac{[f(h_{r,s})]^{\alpha_1}}{[f(l_{r,s})]^{\alpha_2}} > 0$$

then $w_{\theta''_2}^{f,\alpha_2}(p) \subset w_{\theta''_1}^{f,\alpha_1}(p)$, where $I_{r,s} = \{(k,l) : k_{r-1} < k \leq k_r \text{ and } l_{s-1} < l \leq l_s\}$, $k_{r,s} = k_r l_s$, $h_{r,s} = h_r \bar{h}_s$ and $J_{r,s} = \{(s,t) : s_{r-1} < s \leq s_r \text{ and } t_{s-1} < t \leq t_s\}$, $s_{r,s} = s_r t_s$, $l_{r,s} = l_r \bar{l}_s$.

Proof. Let $x \in w_{\theta_2}^{f, \alpha_2}(p)$. We can write

$$\begin{aligned} \frac{1}{[f(\ell_{r,s})]^{\alpha_2}} \sum_{(k,l) \in J_{r,s}} [f(|x_{k,l} - \ell|)]^{p_k} &= \frac{1}{[f(\ell_{r,s})]^{\alpha_2}} \sum_{(k,l) \in J_{r,s} - I_{r,s}} [f(|x_{k,l} - \ell|)]^{p_k} \\ &\quad + \frac{1}{[f(\ell_{r,s})]^{\alpha_2}} \sum_{(k,l) \in I_{r,s}} [f(|x_{k,l} - \ell|)]^{p_k} \\ &\geq \frac{1}{[f(\ell_{r,s})]^{\alpha_2}} \sum_{(k,l) \in I_{r,s}} [f(|x_{k,l} - \ell|)]^{p_k} \\ &\geq \frac{[f(h_{r,s})]^{\alpha_1}}{[f(\ell_{r,s})]^{\alpha_2}} \frac{1}{[f(h_{r,s})]^{\alpha_1}} \sum_{(k,l) \in I_{r,s}} [f(|x_{k,l} - \ell|)]^{p_k}. \end{aligned}$$

Thus if $x \in w_{\theta_2}^{f, \alpha_2}(p)$, then $x \in w_{\theta_1}^{f, \alpha_1}(p)$. \square

From Theorem 2.8. we have the following results.

Corollary 2.1. Let $\theta_1'' = \{(k_r, l_s)\}$ and $\theta_2'' = \{(s_r, t_s)\}$ be two double lacunary sequences such that $I_{r,s} \subset J_{r,s}$ for all $r, s \in \mathbb{N}$ and α_1, α_2 two real numbers such that $0 < \alpha_1 \leq \alpha_2 \leq 1$. If (2.1) holds then

- (i) $w_{\theta_2}^{f, \alpha}(p) \subset w_{\theta_1}^{f, \alpha}(p)$, if $\alpha_1 = \alpha_2 = \alpha$,
- (ii) $w_{\theta_2}^f(p) \subset w_{\theta_1}^{f, \alpha_1}(p)$, if $\alpha_2 = 1$,
- (iii) $w_{\theta_2}^f(p) \subset w_{\theta_1}^f(p)$, if $\alpha_1 = \alpha_2 = 1$.

REFERENCES

1. A. AIZPURU, M. C. LISTÁN-GARCÍA and F. RAMBLA-BARRENO: *Density by moduli and statistical convergence*. Quaest. Math. **37(4)** (2014), 525–530.
2. Y. ALTIN and M. ET: *Generalized difference sequence spaces defined by a modulus function in a locally convex space*. Soochow J. Math. **31(2)** (2005), 233–243.
3. C. BELEN and S. A. MOHIUDDINE: *Generalized weighted statistical convergence and application*. Applied Mathematics and Computation **219** (2013), 9821–9826.
4. V. K. BHARDWAJ and S. DHAWAN: *Density by moduli and lacunary statistical convergence*. Abstr. Appl. Anal. **2016** (2016), Art. ID 9365037, 11 pp.
5. A. CASERTA, G. DI MAIO and L. D. R. KOČINAC: *Statistical convergence in function spaces*. Abstr. Appl. Anal. **2011** (2011), Art. ID 420419, 11 pp.
6. J. S. CONNOR: *The statistical and strong p -Cesaro convergence of sequences*. Analysis **8** (1988), 47–63.
7. H. ÇAKALLI: *Lacunary statistical convergence in topological groups*. Indian J. Pure Appl. Math. **26(2)** (1995), 113–119.
8. H. ÇAKALLI, C. G. ARAS and A. SÖNMEZ: *Lacunary statistical ward continuity*. AIP Conf. Proc. **1676** 020042 (2015), <http://dx.doi.org/10.1063/1.4930468>.

9. H. ÇAKALLI and H. KAPLAN: *A variation on lacunary statistical quasi Cauchy sequences*. Commun. Fac. Sci. Univ. Ank. Sér. A1 Math. Stat. **66(2)** (2017), 71–79.
10. H. ÇAKALLI and H. KAPLAN: *A study on N_θ -quasi-Cauchy sequences*. Abstr. Appl. Anal. **2013** (2013), Article ID 836970, 4 pages.
11. H. ÇAKALLI: *A study on statistical convergence*. Funct. Anal. Approx. Comput. **1(2)** (2009), 19–24.
12. M. ÇINAR, M. KARAKAŞ and M. ET: *On pointwise and uniform statistical convergence of order α for sequences of functions*. Fixed Point Theory And Applications, **2013(33)** (2013), 11 pages.
13. R. ÇOLAK: *Statistical convergence of order α* . Modern Methods in Analysis and Its Applications, New Delhi, India: Anamaya Pub. **2010** (2010), 121–129.
14. G. DI MAIO and L. D. R. KOČINAC: *Statistical convergence in topology*. Topology Appl. **156** (2008), 28–45.
15. M. ET, M. ÇINAR and M. KARAKAŞ: *On λ -statistical convergence of order α of sequences of functions*. J. Inequal. Appl. **2013(204)** (2013), 8 pages.
16. M. ET, Y. ALTIN and H. ALTINOK: *On some generalized difference sequence spaces defined by a modulus function*. Filomat **17** (2003), 23–33.
17. M. ET, A. ALOTAIBI and S. A. MOHIUDDINE: *On (Δ^m, I) statistical convergence of order α* . Scientific World Journal, Article Number: 535419 (2014), 5 pages.
18. M. ET and H. ŞENGÜL: *Some Cesaro-type summability spaces of order α and lacunary statistical convergence of order α* . Filomat **28(8)** (2014), 1593–1602.
19. H. FAST: *Sur la convergence statistique*. Colloq. Math. **2** (1951), 241–244.
20. A. R. FREEDMAN, J. J. SEMBER and M. RAPHAEL: *Some Cesàro-type summability spaces*. Proc. London Math. Soc. (3) **37(3)** (1978), 508–520.
21. J. FRIDY: *On statistical convergence*. Analysis **5** (1985), 301–313.
22. J. A. FRIDY and C. ORHAN: *Lacunary statistical convergence*. Pacific J. Math. **160** (1993), 43–51.
23. J. A. FRIDY and C. ORHAN: *Lacunary statistical summability*. J. Math. Anal. Appl. **173(2)** (1993), 497–504.
24. A. K. GAUR and M. MURSALEEN: *Difference sequence spaces defined by a sequence of moduli*. Demonstratio Math. **31(2)** (1998), 275–278.
25. M. IŞIK and K. E. ET: *On lacunary statistical convergence of order α in probability*. AIP Conference Proceedings **1676** 020045 (2015), doi: <http://dx.doi.org/10.1063/1.4930471>.
26. M. IŞIK and K. E. AKBAŞ: *On λ -statistical convergence of order α in probability*. J. Inequal. Spec. Funct. **8(4)** (2017), 57–64.
27. M. IŞIK: *Generalized vector-valued sequence spaces defined by modulus functions*. J. Inequal. Appl. **2010** Art. ID 457892 (2010), 7 pp.
28. H. KAPLAN and H. ÇAKALLI: *Variations on strong lacunary quasi-Cauchy sequences*. J. Nonlinear Sci. Appl. **9(6)** (2016), 4371–4380.
29. I. J. MADDOX: *Sequence spaces defined by a modulus*. Math. Proc. Camb. Philos. Soc. **100** (1986), 161–166.
30. S. A. MOHIUDDINE, A. ALOTAIBI and M. MURSALEEN: *Statistical convergence of double sequences in locally solid Riesz spaces*. Abstr. Appl. Anal. **2012** Article ID 719729 (2012), 9 pages.

31. M. MURSALEEN and O. H. H. EDELY: *Statistical convergence of double sequences*. Journal of Mathematical Analysis and Applications **288(1)** (2003), 223–231.
32. M. MURSALEEN and S. A. MOHIUDDINE: *On lacunary statistical convergence with respect to the intuitionistic fuzzy normed space*. J. Comput. Appl. Math. **233(2)** (2009), 142–149.
33. H. NAKANO: *Modulared sequence spaces*. Proc. Japan Acad. **27** (1951), 508–512.
34. F. NURAY and E. SAVAŞ: *Some new sequence spaces defined by a modulus function*. Indian J. Pure Appl. Math. **24(11)** (1993), 657–663.
35. R. F. PATTERSON and E. SAVAŞ: *Lacunary statistical convergence of double sequences*. Math. Commun. **10(1)** (2005), 55–61.
36. S. PEHLIVAN and B. FISHER: *Lacunary strong convergence with respect to a sequence of modulus functions*. Comment. Math. Univ. Carolin. **36(1)** (1995), 69–76.
37. S. PEHLIVAN and B. FISHER: *Some sequence spaces defined by a modulus*. Math. Slovaca **45(3)** (1995), 275–280.
38. A. PRINGSHEIM: *Zur Theorie der zweifach unendlichen Zahlenfolgen*. Mathematische Annalen **53(3)** (1900), 289–321.
39. T. SALAT: *On statistically convergent sequences of real numbers*. Math. Slovaca **30** (1980), 139–150.
40. I. J. SCHOENBERG: *The integrability of certain functions and related summability methods*. Amer. Math. Monthly **66** (1959), 361–375.
41. H. STEINHAUS: *Sur la convergence ordinaire et la convergence asymptotique*. Colloq. Math. **2** (1951), 73–74.
42. H. ŞENGÜL: *Some Cesàro-type summability spaces defined by a modulus function of order (α, β)* . Commun. Fac. Sci. Univ. Ank. Sér. A1 Math. Stat. **66(2)** (2017), 80–90.
43. H. ŞENGÜL and M. ET: *On lacunary statistical convergence of order α* . Acta Math. Sci. Ser. B Engl. Ed. **34(2)** (2014), 473–482.

Hacer Şengül
Faculty of Education
Harran University
Osmanbey Campus
63190 Şanlıurfa, Turkey
hacer.sengul@hotmail.com

Mikail Et
Department of Mathematics
Fırat University
23119 Elazığ, Turkey
mikaillet68@gmail.com

Yavuz Altın
Department of Mathematics
Fırat University
23119 Elazığ, Turkey
yaltin23@yahoo.com