



ONE-SIDED GENERALIZED (α, β) –REVERSE DERIVATIONS OF ASSOCIATIVE RINGS

Ayşe Engin and Neşet Aydın

Department of Mathematics, Faculty of Science
Çanakkale Onsekiz Mart University, 17100 Çanakkale, Türkiye

ORCID IDs: Ayşe Engin
Neşet Aydın

 <https://orcid.org/0000-0002-1626-5498>
 <https://orcid.org/0000-0002-7193-3399>

Abstract. In this paper, we introduce the notion of the one-sided generalized (α, β) -reverse derivation of a ring R . Let R be a semiprime ring, ϱ be a non-zero ideal of R , α be an epimorphism of ϱ , β be a homomorphism of ϱ (α be a homomorphism of ϱ , β be an epimorphism of ϱ) and $\gamma : \varrho \rightarrow R$ be a non-zero (α, β) -reverse derivation. We show that there exists $F : \varrho \rightarrow R$, an l -generalized (α, β) -reverse derivation (an r -generalized (α, β) -reverse derivation) associated with γ iff $F(\varrho), \gamma(\varrho) \subset C_R(\varrho)$ and F is an r -generalized (β, α) -derivation (an l -generalized (β, α) -derivation) associated with (β, α) -derivation γ on ϱ . This theorem generalizes the results of A. Aboubakr and S. Gonzalez proved in [1, Theorem 3.1, and Theorem 3.2].

Keywords: Semiprime ring, prime ring, one-sided generalized (α, β) –reverse derivation, (α, β) –reverse derivation.

1. Introduction

Throughout the paper, R is an associative ring with Z , which the center of R denotes. Recall that a ring R is prime if for any $r_1, r_2 \in R$, $r_1 R r_2 = (0)$ implies $r_1 = 0$ or $r_2 = 0$, and is a semiprime in case $r_1 \in R$, $r_1 R r_1 = (0)$ implies $r_1 = 0$. For $r_1, r_2 \in R$, $[r_1, r_2]$ denotes the element $r_1 r_2 - r_2 r_1$. The symbol $[r_1, r_2]$ stands for Lie commutator of r_1 and r_2 and it satisfies the basic commutator identities: for each $r_1, r_2, r_3 \in R$, $[r_1 + r_2, r_3] = [r_1, r_3] + [r_2, r_3]$, $[r_1, r_2 + r_3] = [r_1, r_2] + [r_1, r_3]$, $[r_1 r_2, r_3] = r_1 [r_2, r_3] + [r_1, r_3] r_2$, $[r_1, r_2 r_3] = [r_1, r_2] r_3 + r_2 [r_1, r_3]$. We denote the

Received March 22, 2022, revised: July 11, 2023, accepted: September 04, 2023

Communicated by Dijana Mosić

Corresponding Author: Ayşe Engin. E-mail addresses: ayseayraan@gmail.com (A. Engin), neseta@comu.edu.tr (N. Aydın)

2010 *Mathematics Subject Classification.* Primary 16N60; Secondary 16W30

identity mapping of R by id_R ; that is, the mapping $id_R : R \rightarrow R$ is defined as $id_R(r_1) = r_1$, for all $r_1 \in R$. For a non-empty subset A of R , $C_R(A)$ is defined as $C_R(A) = \{r \in R : [r, x] = 0, \text{ for all } x \in A\}$.

Let α, β be any two mapping of R . An additive mapping $\delta : R \rightarrow R$ is called an (α, β) -derivation if $\delta(r_1r_2) = \delta(r_1)\alpha(r_2) + \beta(r_1)\delta(r_2)$ holds, for all $r_1, r_2 \in R$. An additive mapping $\varphi : R \rightarrow R$ is called a right generalized (α, β) -derivation (a left generalized (α, β) -derivation) of R associated with δ , if $\varphi(r_1r_2) = \delta(r_1)\alpha(r_2) + \beta(r_1)\varphi(r_2)$ ($\varphi(r_1r_2) = \varphi(r_1)\alpha(r_2) + \beta(r_1)\delta(r_2)$), for all $r_1, r_2 \in R$ and φ is said to be a generalized (α, β) -derivation of R with δ if it is both a right and a left generalized (α, β) -derivation of R associated with δ .

Many authors have investigated the relationship between the commutativity of a ring and the act of derivation ((α, β) -derivation, reverse derivation, (α, β) -reverse derivation, generalized reverse derivation, etc.) defined on the ring. Herstein (1957) was the first to introduce the concept of reverse derivation. An additive mapping $g : R \rightarrow R$ is a reverse derivation if $g(r_1r_2) = g(r_2)r_1 + r_2g(r_1)$, for all $r_1, r_2 \in R$. In [4], it is shown that if a prime ring R with a characteristic different from two admits non-zero reverse derivation g , then g is a derivation of R . An additive mapping $d : R \rightarrow R$ is an (α, β) -reverse derivation if $d(r_1r_2) = d(r_2)\alpha(r_1) + \beta(r_2)d(r_1)$, for all $r_1, r_2 \in R$. In [8], Chaudhry and Thaheem shown that if a semiprime ring R admits non-zero (α, β) -reverse derivation d , then d is (α, β) -reverse derivation of R . Here, α and β are automorphism of R . An additive mapping $H : R \rightarrow R$ is called l -generalized reverse derivation (r -generalized reverse derivation) In [1], A. Aboubakr and S. Gonzalez (2015) introduced one-sided generalized reverse derivation. An additive mapping $H : R \rightarrow R$ is called an l -generalized reverse derivation (r -generalized reverse derivation) if there exists a reverse derivation $g : R \rightarrow R$ such that $H(r_1r_2) = H(r_2)r_1 + r_2g(r_1)$ ($H(r_1r_2) = g(r_2)r_1 + r_2H(r_1)$), for all $r_1, r_2 \in R$. In [1], they have indicated that if a semiprime ring R admits non-zero one-sided generalized reverse derivation H associated with reverse derivation g , then H is a one-sided generalized derivation with associated derivation g . Reverse derivation, generalized reverse derivation, (α, β) -reverse derivation, generalized (α, β) -reverse derivation, multiplicative reverse derivation, multiplicative generalized reverse derivation, multiplicative (α, β) -reverse derivation, and multiplicative generalized (α, β) -reverse derivation of prime or semiprime rings have been studied by a lot of scholars in the literature. (see [2],[3],[4], [9],[10],[12],[13],[14],[15],[16].)

This paper extends the notion of one-sided reverse derivation to one-sided generalized (α, β) -reverse derivation.

Definition 1.1. Let R be a ring, α, β be a mapping of R , and γ be an (α, β) -reverse derivation of R . An additive mapping $F : R \rightarrow R$ is said to be an r -generalized (α, β) -reverse derivation of R associated with γ if

$$F(r_1r_2) = \gamma(r_2)\alpha(r_1) + \beta(r_2)F(r_1)$$

for all $r_1, r_2 \in R$, F is said to be an l -generalized (α, β) -reverse derivation of R associated with γ if

$$F(r_1r_2) = F(r_2)\alpha(r_1) + \beta(r_2)\gamma(r_1)$$

for all $r_1, r_2 \in R$ and F said to be a generalized (α, β) -reverse derivation of R associated with γ if it is both an r -generalized and l -generalized (α, β) -reverse derivation of R associated with γ .

When $\alpha = \beta = id_R$, an r -generalized (l -generalized) (α, β) -reverse derivation is a r -generalized (l -generalized) reverse derivation. Thus, the one-sided generalized reverse derivation is a special case of one-sided generalized (α, β) -reverse derivation.

This study consists of 2 parts. In the first part, we show that If R is a 2-torsion free semiprime ring, α, β are automorphisms of R , and $\gamma : R \rightarrow R$ is a non-zero (α, β) -reverse derivation, then γ is an (α, β) -derivation on R . With this result, we will show that the concepts of (α, β) -reverse derivation and (α, β) -derivation overlap in 2 torsion-free semiprime rings in which α and β are automorphisms of the ring. In the second part, we give a generalization of [1, Theorem 3.1, Theorem 3.2, and Corollary 3.3], which is the main result of the article. In that case, one-sided generalized (α, β) -reverse derivation and one-sided generalized (β, α) -derivation overlap in a semiprime ring where only one of α and β is an epimorphism of the ring. Thus we will show that the intersection of the set of all generalized (α, β) -derivation and the set of all generalized (α, β) -reverse derivation is different from the empty set. At the end of the paper, we showed that in case α is a homomorphism of R and β is an epimorphism of R ; there is no non-zero generalized (α, β) -reverse derivation associated with (α, β) -reverse derivation of noncommutative prime ring R .

From now on, R is an associative ring, Z is the center of R , and $\alpha, \beta : R \rightarrow R$ are homomorphisms.

2. Preliminary

In this section, we give some auxiliary results that will need later. We begin our discussion with several examples related to (α, β) -reverse derivation and one-sided generalized (α, β) -reverse derivation.

Lemma 2.1. [7, Lemma 3] *If the prime ring R contains a commutative non-zero right ideal I , then R is commutative.*

Lemma 2.2. [7, Lemma 4] *Let b and ab be in the center of a prime ring R . If b is not zero, then a is in Z , the center of R .*

Lemma 2.3. [11, Corollary 2.1] *Let R be a 2-torsion free semiprime ring, α, β be automorphisms of R and $L \not\subseteq Z(R)$ be a non-zero square-closed Lie ideal of R . If $\delta : R \rightarrow L$ satisfying*

$$(2.1) \quad (a^2)^\delta = a^\delta \alpha(a) + \beta(a) a^\delta, \text{ for all } a \in L$$

and $a^\delta, \beta(a) \in L$, then δ is a (α, β) -derivation on L .

Example 2.1. Consider the ring $R = \left\{ \begin{bmatrix} a_{11} & a_{12} \\ 0 & a_{22} \end{bmatrix} : a_{11}, a_{12}, a_{22} \in \mathbb{Z} \right\}$, where \mathbb{Z} the ring of integers. Let us define $\alpha : R \rightarrow R$, $\beta : R \rightarrow R$, and $d : R \rightarrow R$ as follows:

$$\begin{aligned} \alpha \left(\begin{bmatrix} a_{11} & a_{12} \\ 0 & a_{22} \end{bmatrix} \right) &= \begin{bmatrix} 0 & 0 \\ 0 & a_{11} \end{bmatrix} \\ \beta \left(\begin{bmatrix} a_{11} & a_{12} \\ 0 & a_{22} \end{bmatrix} \right) &= \begin{bmatrix} a_{22} & 0 \\ 0 & 0 \end{bmatrix} \\ d \left(\begin{bmatrix} a_{11} & a_{12} \\ 0 & a_{22} \end{bmatrix} \right) &= \begin{bmatrix} 0 & a_{11} - a_{22} \\ 0 & 0 \end{bmatrix}. \end{aligned}$$

It is easy to check that d is both an (α, β) -reverse derivation and an (α, β) -derivation.

Example 2.2. Consider the ring $R = \left\{ \begin{bmatrix} a_{11} & a_{12} \\ 0 & a_{22} \end{bmatrix} : a_{11}, a_{12}, a_{22} \in \mathbb{Z} \right\}$, where \mathbb{Z} the ring of integers. Define the mappings $\alpha : R \rightarrow R$, $\beta : R \rightarrow R$, and $d : R \rightarrow R$ as follows:

$$\begin{aligned} \alpha \left(\begin{bmatrix} a_{11} & a_{12} \\ 0 & a_{22} \end{bmatrix} \right) &= \begin{bmatrix} 0 & 0 \\ 0 & a_{11} \end{bmatrix} \\ \beta \left(\begin{bmatrix} a_{11} & a_{12} \\ 0 & a_{22} \end{bmatrix} \right) &= \begin{bmatrix} a_{22} & 0 \\ 0 & 0 \end{bmatrix} \\ d \left(\begin{bmatrix} a_{11} & a_{12} \\ 0 & a_{22} \end{bmatrix} \right) &= \begin{bmatrix} 0 & -a_{12} \\ 0 & 0 \end{bmatrix}. \end{aligned}$$

It is easy to check that d is an (α, β) -reverse derivation. But d is not an (α, β) -derivation.

Example 2.3. Consider the ring $R = \left\{ \begin{bmatrix} a_{11} & a_{12} \\ 0 & a_{22} \end{bmatrix} : a_{11}, a_{12}, a_{22} \in \mathbb{Z} \right\}$, where \mathbb{Z} the ring of integers. Define the mappings $\alpha : R \rightarrow R$, $\beta : R \rightarrow R$, and $d : R \rightarrow R$ as follows:

$$\begin{aligned} \alpha \left(\begin{bmatrix} a_{11} & a_{12} \\ 0 & a_{22} \end{bmatrix} \right) &= \begin{bmatrix} 0 & 0 \\ 0 & a_{22} \end{bmatrix} \\ \beta \left(\begin{bmatrix} a_{11} & a_{12} \\ 0 & a_{22} \end{bmatrix} \right) &= \begin{bmatrix} a_{11} & 0 \\ 0 & 0 \end{bmatrix} \\ d \left(\begin{bmatrix} a_{11} & a_{12} \\ 0 & a_{22} \end{bmatrix} \right) &= \begin{bmatrix} 0 & a_{12} \\ 0 & 0 \end{bmatrix}. \end{aligned}$$

It is easy to check that d is an (α, β) -derivation. But d is not an (α, β) -reverse derivation.

Example 2.4. Let $(R_1, +, *)$ be a commutative ring and (R_2, \oplus, \otimes) be a noncommutative ring. Let's consider operation $\otimes : R_2 \times R_2 \rightarrow R_2$, $r \otimes s = s \otimes r$. With these operations (R_2, \oplus, \otimes) called opposite ring and it is shown R_2^{op} . α, β are homomorphisms of R_2 , $\delta : R_2 \rightarrow R_2^{op}$ is an (β, α) -derivation, and $\varphi : R_2 \rightarrow R_2^{op}$ is a left generalized (β, α) -derivation with δ . Define the mappings $\tilde{\alpha}, \tilde{\beta} : R_2 \times R_1 \rightarrow R_2 \times R_1$, and $\tilde{\delta}, \tilde{\varphi} : R_2 \times R_1 \rightarrow R_2^{op} \times R_1$ as follows:

$$\begin{aligned} \tilde{\alpha}(r, s) &= (\alpha(r), s) \\ \tilde{\beta}(r, s) &= (\beta(r), s) \\ \tilde{\delta}(r, s) &= (\delta(r), s) \\ \tilde{\varphi}(r, s) &= (\varphi(r), s). \end{aligned}$$

Then it is straightforward to verify that $\tilde{\varphi}$ is an l -generalized (α, β) -reverse derivation with (α, β) -reverse derivation $\tilde{\delta}$ of $R_2 \times R_1$. But $\tilde{\varphi}$ is not a generalized (α, β) -derivation with (α, β) -derivation $\tilde{\delta}$ of $R_2 \times R_1$.

Example 2.5. Let $(R_1, +, *)$ and (R_2, \oplus, \otimes) be rings as defined in example 2.4. Let α, β be homomorphisms of R_2 , $\delta : R_2 \rightarrow R_2^{\text{op}}$ be an (β, α) -derivation, and $\varphi : R_2 \rightarrow R_2^{\text{op}}$ be a right generalized (β, α) -derivation with δ . Define the mappings $\tilde{\alpha}, \tilde{\beta} : R_2 \times R_1 \rightarrow R_2 \times R_1$ and $\tilde{\delta}, \tilde{\varphi} : R_2 \times R_1 \rightarrow R_2^{\text{op}} \times R_1$ as follows:

$$\begin{aligned}\tilde{\alpha}(r, s) &= (\alpha(x), s) \\ \tilde{\beta}(r, s) &= (\beta(x), s) \\ \tilde{\delta}(r, s) &= (\delta(x), s) \\ \tilde{\varphi}(r, s) &= (\varphi(x), s).\end{aligned}$$

Then it is straightforward to verify that $\tilde{\varphi}$ is an r -generalized (α, β) -reverse derivation with (α, β) -reverse derivation $\tilde{\delta}$ of $R_2 \times R_1$. But $\tilde{\varphi}$ is not a generalized (α, β) -derivation with (α, β) -derivation $\tilde{\delta}$ of $R_2 \times R_1$.

3. (α, β) -Reverse Derivation

Theorem 3.1. *Let R be a 2-torsion free semiprime ring, α, β be automorphisms of R . If $\gamma : R \rightarrow R$ is a non-zero (α, β) -reverse derivation, then γ is an (α, β) -derivation on R .*

Proof. Suppose that R is non-commutative ring. Let $r_1 \in R$. From the hypothesis, we get

$$\gamma(r_1^2) = \gamma(r_1)\alpha(r_1) + \beta(r_1)\gamma(r_1).$$

This equation ensures equality of (2.1). We know that the ring R is a square closed Lie ideal of R . So, we can think of R instead of L in Lemma 2.3. Thus, γ is an (α, β) -derivation on R because of Lemma 2.3. While R is a commutative ring, (α, β) -reverse derivation of R is (α, β) -derivation of R . So, the proof ends. \square

Theorem 3.2. *Let R be a semiprime ring, ϱ is a non-zero two-sided ideal of R , α be an epimorphism of ϱ and β be a homomorphism of ϱ (or α be a homomorphism of ϱ and β be an epimorphism of ϱ). There exists $\gamma : \varrho \rightarrow R$ a non-zero (α, β) -reverse derivation iff $\gamma(\varrho) \subset C_R(\varrho)$ and γ is (β, α) -derivation on ϱ .*

Proof. We only prove case of no parenthesis. The another one has the same argument. Let $x_1, x_2, x_3 \in \varrho$. Since γ is (α, β) -reverse derivation on ϱ , we have

$$(3.1) \quad \gamma(x_1x_2x_3) = \gamma(x_1(x_2x_3)) = \gamma(x_3)\alpha(x_2)\alpha(x_1) + \beta(x_3)\gamma(x_2)\alpha(x_1) + \beta(x_2)\beta(x_3)\gamma(x_1)$$

and

$$(3.2) \quad \gamma(x_1x_2x_3) = \gamma((x_1x_2)x_3) = \gamma(x_3)\alpha(x_1)\alpha(x_2) + \beta(x_3)\gamma(x_2)\alpha(x_1) + \beta(x_3)\beta(x_2)\gamma(x_1).$$

From (3.1) and (3.2),

$$(3.3) \quad \gamma(x_3) [\alpha(x_1), \alpha(x_2)] = [\beta(x_3), \beta(x_2)] \gamma(x_1).$$

Replacing x_3 by x_2 in (3.3),

$$\gamma(x_2) [\alpha(x_1), \alpha(x_2)] = 0$$

for all $x_1, x_2 \in \varrho$. Because α is an epimorphism of ϱ , for each $x_1, x_2 \in \varrho$, we get

$$(3.4) \quad \gamma(x_2) [x_1, \alpha(x_2)] = 0.$$

Take $r \in R$. Substituting $x_1 x_3 r$ for x_1 in (3.4), we obtain $\gamma(x_2) x_1 x_3 [r, \alpha(x_2)] = 0$, for all $x_1, x_2, x_3 \in \varrho, r \in R$. So implies that

$$(3.5) \quad \gamma(x_2) \varrho \varrho [R, \alpha(x_2)] = (0)$$

for all $x_2 \in \varrho$. Because ϱ is a semiprime ring, it must contain a family ρ of prime ideals such that $\cap \rho = (0)$. Let ρ_φ be a typical member of this family and $x_2 \in \varrho$; by (3.5),

$$\gamma(x_2) \varrho \subset \rho_\varphi \text{ or } [R, \alpha(x_2)] \subset \rho_\varphi.$$

Let $M = \{x_2 \in \varrho : \gamma(x_2) \varrho \subset \rho_\varphi\}$ and $N = \{x_2 \in \varrho : [R, \alpha(x_2)] \subset \rho_\varphi\}$. Clearly, each group M and N is additive subgroup of ϱ such that $\varrho = M \cup N$. But a group cannot be a set union of two proper subgroups. Hence, $M = \varrho$ or $N = \varrho$. Since ρ_φ is ideal of ϱ , it holds that $\gamma(\varrho) \varrho [R, \alpha(\varrho)] \subset \rho_\varphi$. Thus $\gamma(\varrho) \varrho [R, \alpha(\varrho)] \subset \cap \rho = (0)$. Because α is an epimorphism of ϱ , it provides that $\gamma(\varrho) \varrho [R, \varrho] = 0$. Let $x_1, x_2, x_3 \in \varrho, r \in R$. Means that,

$$(3.6) \quad \gamma(x_1) x_2 [r, x_3] = 0.$$

Let $x_4 \in \varrho$. In (3.6), replacing r by $x_4 \gamma(x_1)$ and x_3 by x_2 , we get

$$(3.7) \quad \gamma(x_1) x_2 x_4 [\gamma(x_1), x_2] = 0.$$

In (3.6), substituting x_2 by x_4 , we get $\gamma(x_1) x_4 [r, x_3] = 0$. In this equation replacing x_3 by x_2 , r by $\gamma(x_1)$ and multiply from the left by x_2 , it holds

$$(3.8) \quad x_2 \gamma(x_1) x_4 [\gamma(x_1), x_2] = 0.$$

From (3.7) and (3.8),

$$(3.9) \quad [\gamma(x_1), x_2] x_4 [\gamma(x_1), x_2] = 0, \text{ for all } x_1, x_2, x_4 \in \varrho.$$

Since ϱ is a semiprime ring,

$$[\gamma(x_1), x_2] = 0, \text{ for all } x_1, x_2 \in \varrho.$$

That is $\gamma(\varrho) \subset C_R(\varrho)$. We get

$$\begin{aligned} \gamma(x_1 x_2) &= \gamma(x_2) \alpha(x_1) + \beta(x_2) \gamma(x_1) \\ &= \gamma(x_1) \beta(x_2) + \alpha(x_1) \gamma(x_2) \end{aligned}$$

for all $x_1, x_2 \in \varrho$. This means that γ is (β, α) -derivation on ϱ . The converse is trivial. \square

If consider R instead of ϱ in Theorem 3.2, we get

Corollary 3.1. *Let R be a semiprime ring, α be an epimorphism of R and β be a homomorphism of R (or α be a homomorphism of R and β be an epimorphism of R). There exists $\gamma : R \rightarrow R$ a non-zero (α, β) -reverse derivation iff central γ is (β, α) -derivation on R .*

Corollary 3.2. *Let R be a prime ring, α be an epimorphism of R and β be a homomorphism of R (or α be a homomorphism of R and β be an epimorphism of R). There exists $\gamma : R \rightarrow R$ a non-zero (α, β) -reverse derivation iff R is commutative and γ is an (α, β) -derivation of R .*

Proof. We only prove a case in which α is an epimorphism of R and β is a homomorphism of R . Another case has the similar argument. By Corollary 3.1, γ is a central (β, α) -derivation of R . Let $r_1, r_2 \in R$. It is clear that

$$[\gamma(r_1 r_2), \beta(r_2)] = 0.$$

Applying Lie commutator features, we get

$$\begin{aligned} [\gamma(r_2)\alpha(r_1) + \beta(r_2)\gamma(r_1), \beta(r_2)] &= [\gamma(r_2)\alpha(r_1), \beta(r_2)] + [\beta(r_2)\gamma(r_1), \beta(r_2)] \\ &= \gamma(r_2) [\alpha(r_1), \beta(r_2)] + [\gamma(r_2), \beta(r_2)] \alpha(r_1) \\ &\quad + \beta(r_2) [\gamma(r_1), \beta(r_2)] + [\beta(r_2), \beta(r_2)] \gamma(r_1) \end{aligned}$$

for all $r_1, r_2 \in R$. In the last equation, since $\gamma(r_1), \gamma(r_2) \in Z$, we have

$$\gamma(r_2) [\alpha(r_1), \beta(r_2)] = 0$$

for all $r_1, r_2 \in R$. Let $r_3 \in R$. Since α is an epimorphism of R , we get

$$\gamma(r_2) r_3 [\alpha(r_1), \beta(r_2)] = 0.$$

Thus, for each $r_2 \in R$, we write

$$\gamma(r_2) R [\alpha(r_1), \beta(r_2)] = (0).$$

By the primeness of R , for each $r_2 \in R$, we get

$$\gamma(r_2) = 0 \text{ or } \beta(r_2) \in Z.$$

Let $M = \{r_2 \in R : \gamma(r_2) = 0\}$ and $N = \{r_2 \in R : \beta(r_2) \in Z\}$. Clearly, each group M and N is additive subgroup of R such that $R = M \cup N$. But a subgroup cannot be a set union of two proper subgroups. Hence, $M = R$ or $N = R$. Since γ is a non-zero (α, β) -reverse derivation of R , it happens $\beta(R) \subset Z$. Since $\gamma(r_1 r_2) \in Z$ and Z is a subring of R , we have

$$\gamma(r_2)\alpha(r_1) \in Z, \text{ for all } r_1, r_2 \in R.$$

In view of Lemma 2.2, for each $r_1 \in R$, we have $\alpha(r_1) \in Z$. In addition, since α is an epimorphism of R , we have R is commutative. Therefore, we conclude that

$$\gamma(r_1 r_2) = \gamma(r_2 r_1) = \gamma(r_1)\alpha(r_2) + \beta(r_1)\gamma(r_2)$$

for all $r_1, r_2 \in R$. This implies γ is an (α, β) -derivation of R . \square

4. One-Sided Generalized (α, β) –Reverse Derivation

Theorem 4.1. *Let R be a semiprime ring, ϱ is a non-zero two-sided ideal of R , α be an epimorphism of ϱ , β be homomorphism of ϱ and $\gamma : \varrho \rightarrow R$ be a non-zero (α, β) -reverse derivation. There exists $F : \varrho \rightarrow R$, a l -generalized (α, β) -reverse derivation associated with γ iff $F(\varrho), \gamma(\varrho) \subset C_R(\varrho)$ and F is r -generalized (β, α) -derivation associated with (β, α) -derivation γ on ϱ .*

Proof. Let $x_1, x_2, x_3 \in \varrho$. Using the definition of l -generalized (α, β) -reverse derivation one can easily see that

$$(4.1) \quad F(x_1(x_2x_3)) = F(x_3)\alpha(x_2)\alpha(x_1) + \beta(x_3)\gamma(x_2)\alpha(x_1) + \beta(x_2)\beta(x_3)\gamma(x_1)$$

and

$$(4.2) \quad F((x_1x_2)x_3) = F(x_3)\alpha(x_1)\alpha(x_2) + \beta(x_3)\gamma(x_2)\alpha(x_1) + \beta(x_3)\beta(x_2)\gamma(x_1)$$

Combining (4.1) and (4.2),

$$(4.3) \quad F(x_3) [\alpha(x_2), \alpha(x_1)] = [\beta(x_3), \beta(x_2)] \gamma(x_1).$$

Substituting x_3 by x_2 in (4.3),

$$F(x_2) [\alpha(x_2), \alpha(x_1)] = 0$$

for all $x_1, x_2 \in \varrho$. Since α is an epimorphism of ϱ , for each $x_1, x_2 \in \varrho$, we have

$$(4.4) \quad F(x_2) [\alpha(x_2), x_1] = 0.$$

Taking $x_3 \in \varrho, r \in R$. Replacing x_1 by x_1x_3r in (4.4), $F(x_2)x_1x_3r [\alpha(x_2), r] = 0$. For each $x_2 \in \varrho$, we have $F(x_2)\varrho\varrho [\alpha(x_2), R] = (0)$. Now, when similar steps are applied to the steps from equality (3.5) to equality (3.9), for each $x_1, x_2, x_3 \in \varrho$, we have $[F(x_1), x_2] x_3 [F(x_1), x_2] = 0$. Since ϱ is a semiprime ring,

$$[F(x_1), x_2] = 0$$

for all $x_1, x_2 \in \varrho$. That is $F(\varrho) \subset C_R(\varrho)$. Moreover, if γ is (α, β) –reverse derivation of R , then by Theorem 3.2, $\gamma(\varrho) \subset C_R(\varrho)$ and γ is an (β, α) –derivation on ϱ . Hence,

$$\begin{aligned} F(x_1x_2) &= F(x_2)\alpha(x_1) + \beta(x_2)\gamma(x_1) \\ &= \gamma(x_1)\beta(x_2) + \alpha(x_1)F(x_2) \end{aligned}$$

for all $x_1, x_2 \in \varrho$ and F is a r -generalized (β, α) –derivation associated with (β, α) –derivation γ on ϱ . The converse is a trivial. \square

Corollary 4.1. *Let R be a semiprime ring, α be an epimorphism of R , β be homomorphism of R and $\gamma : R \rightarrow R$ be a non-zero (α, β) -reverse derivation. There exists $F : R \rightarrow R$, a l -generalized (α, β) -reverse derivation associated with γ iff $F(I), \gamma(I) \subset Z$ and F is r -generalized (β, α) -derivation associated with (β, α) -derivation γ of R .*

Theorem 4.2. *Let R be a semiprime ring, ϱ is a non-zero two-sided ideal of R , α be a homomorphism of ϱ , β be an epimorphism of ϱ and $\gamma : \varrho \rightarrow R$ be a non-zero (α, β) -reverse derivation. There exists $F : \varrho \rightarrow R$, a r -generalized (α, β) -reverse derivation associated with γ iff $F(\varrho), \gamma(\varrho) \subset C_R(\varrho)$ and F is l -generalized (β, α) -derivation associated with (β, α) -derivation γ on ϱ .*

Proof. By a similar proof in Theorem 4.1, desired is achieved. \square

Corollary 4.2. *Let R be a semiprime ring, α be an homomorphism of R , β be an epimorphism of R and $\gamma : R \rightarrow R$ be a non-zero (α, β) -reverse derivation. There exists $F : R \rightarrow R$, a r -generalized (α, β) -reverse derivation associated with γ iff $F(R), \gamma(R) \subset Z$ and F is l -generalized (β, α) -derivation associated with (β, α) -derivation γ of R .*

Theorem 4.3. *Let R be a semiprime ring, α and β be an epimorphisms of R and $\gamma : R \rightarrow R$ be a non-zero (α, β) -reverse derivation. If there exists $F : R \rightarrow R$, a non-zero l -generalized (α, β) -reverse derivation (r -generalized (α, β) -reverse derivation) associated with γ then R contains a non-zero central ideal.*

Proof. Assume that $F : R \rightarrow R$ is a l -generalized (α, β) -reverse derivation associated with non-zero (α, β) -reverse derivation γ of R . From Corollary 4.1, it holds $\gamma(R), F(R) \subset Z$. For all $r_1, r_2 \in R$,

$$[F(r_1 r_2), \beta(r_2)] = 0$$

is obtained. This means

$$F(r_2) [\alpha(r_1), \beta(r_2)] = 0$$

for all $r_1, r_2 \in R$. Because α is an epimorphism of R , for each $r_1, r_2 \in R$, we get $F(r_2) [r_1, \beta(r_2)] = 0$. Let r_3 . Replacing r_1 by $r_1 r_3$ in $F(r_2) [r_1, \beta(r_2)] = 0$, we get

$$F(r_2) r_1 [r_3, \beta(r_2)] = 0.$$

Now, when similar steps are applied to the steps from equality (3.5) to equality (3.9), for each $r_1, r_2, r_3 \in R$, we have $F(r_1) [r_2, \beta(r_3)] = 0$. Because β is an epimorphism of R , we get $F(r_1) [r_2, r_3] = 0$. That is

$$[F(r_1) r_2, r_3] = 0$$

for all $r_1, r_2, r_3 \in R$. This means $F(R)R \subset Z$. Since F is non-zero l -generalized (α, β) -reverse derivation and R is semiprime, $F(R)R \neq (0)$. $F(R)R$ is obviously central ideal of R . The proof has a similar argument if F is r -generalized (α, β) -reverse derivation of R . \square

Corollary 4.3. *Let R be a semiprime ring, α and β be an epimorphisms of R and $\gamma : R \rightarrow R$ be a non-zero (α, β) -reverse derivation. If there exists $F : R \rightarrow R$, a non-zero generalised (α, β) -reverse derivation associated with γ then R contains a non-zero central ideal.*

Corollary 4.4. *Let R be a prime ring, α and β be an epimorphisms of R and $\gamma : R \rightarrow R$ be a non-zero (α, β) -reverse derivation. If there exists $F : R \rightarrow R$, a non-zero generalized (α, β) -reverse derivation associated with d then R is commutative ring and F is a generalized (α, β) -derivation associated with an (α, β) -derivation γ of R .*

Theorem 4.4. *Let R be a noncommutative prime ring, α be a homomorphism of R and β be an epimorphism of R . If $F : R \rightarrow R$ is a generalized (α, β) -reverse derivation associated with non-zero (α, β) -reverse derivation γ of R then $F = \gamma$.*

Proof. Assume that $F : R \rightarrow R$ is a generalized (α, β) -reverse derivation associated with non-zero (α, β) -reverse derivation γ of R . Let $r_1, r_2 \in R$. Then,

$$F(r_1r_2) = F(r_2)\alpha(r_1) + \beta(r_2)\gamma(r_1) = \gamma(r_2)\alpha(r_1) + \beta(r_2)F(r_1).$$

That is,

$$(F - \gamma)(r_2)\alpha(r_1) - \beta(r_2)(F - \gamma)(r_1) = 0.$$

Let us introduce mapping $\varphi : R \rightarrow R$, $\varphi(r_1) = (F - \gamma)(r_1)$. Moreover, the last equation implies that

$$(4.5) \quad \varphi(r_2)\alpha(r_1) = \beta(r_2)\varphi(r_1).$$

Let $r_1, r_2 \in R$. Since F is an r -generalized (α, β) -reverse derivation (l - of generalized (α, β) -reverse derivation) R and γ is an (α, β) -reverse derivation of R , the mapping φ respectively ensures:

$$\begin{aligned} \varphi(r_1r_2) &= (F - \gamma)(r_1r_2) = \gamma(r_2)\alpha(r_1) + \beta(r_2)F(r_1) - \gamma(r_2)\alpha(r_1) + \beta(r_2)\gamma(r_1) \\ &= \beta(r_2)\varphi(r_1) \end{aligned}$$

and

$$\begin{aligned} \varphi(r_1r_2) &= (F - \gamma)(r_1r_2) = F(r_2)\alpha(r_1) + \beta(r_2)\gamma(r_1) - \gamma(r_2)\alpha(r_1) + \beta(r_2)\gamma(r_1) \\ &= \varphi(r_2)\alpha(r_1). \end{aligned}$$

That is,

$$(4.6) \quad \varphi(r_1r_2) = \beta(r_2)\varphi(r_1).$$

$$(4.7) \quad \varphi(r_1r_2) = \varphi(r_2)\alpha(r_1).$$

Let $r_3 \in R$. Writing r_2r_3 by r_2 in (4.5), we get

$$\varphi(r_2r_3)\alpha(r_1) - \beta(r_2r_3)\varphi(r_1) = 0.$$

In the last equality, using (4.6) and (4.7), we get

$$\beta([r_3, r_2])\varphi(r_1) = 0$$

for all $r_1, r_2, r_3 \in R$. Because β is an epimorphism, for each $r_1, r_2, r_3 \in R$, we have $[r_3, r_2]\varphi(r_1) = 0$. Given that R is a noncommutative prime ring, we get $\varphi = 0$. That is, $F = \gamma$. \square

Acknowledgment

This paper is derived from the first author's doctoral dissertation supervised by the second author.

REFERENCES

1. A. ABOUBAKR and S. GONZALEZ: (α, β) -reverse derivation on semiprime rings. *Siberian Mathematical Journal* **56(2)** (2015), 199–205.
2. A. ALI and A. BANO: *Multiplicative (generalized) reverse derivations on semiprime ring*. *European Journal of Pure and Applied Mathematics* **11(3)** (2018), 717–729.
3. C. J. S. REDDY, N. SUBBARAYUDU and C. V. S. R. REDDY: *A Note on multiplicative (generalized)- (α, β) -reverse derivations on left Ideals in prime rings*. *Global Journal of Pure and Applied Mathematics* **18(2)** (2022), 425–431.
4. G. S. SANDHU and D. KUMAR: *Annihilator conditions of multiplicative reverse derivations on prime rings*. *International electronic journal of algebra* **25(25)** (2019), 87–103.
5. I. N. HERSTEIN: *Jordan derivations of prime rings*. *Proc. Amer. Math. Soc.* **8(6)** (1957), 1104–1110.
6. I. N. HERSTEIN: *Rings with involution*. The University of Chicago Press, 1976.
7. J. H. MAYNE: *Centralizing mappings of prime rings*. *Canad. Math. Bull.* **27(1)** (1984), 122–126.
8. M. A. CHAUDHRY and A. B. THAHEEM: *On (α, β) -derivations of semiprime rings II*. *Demonstratio Mathematica* **37(4)** (2004), 793–802.
9. M. S. SAMMAN and A. B. THAHEEM: *Derivations on semiprime rings*. *Int. Journal of Pure and Applied Math.* **2(4)** (2003), 465–472.
10. M. S. SAMMAN and N. ALYAMANI: *Derivations and reverse derivations in semiprime rings*. *Int. Math. Forum* **39(2)** (2007), 1895–1902.
11. N. U. REHMAN and E. K. SÖĞÜTÇÜ: *Lie ideals and jordan Triple (α, β) -derivations in rings*. *Commun. Fac. Sci. Univ. Ank. Ser. A1 Math. Stat.* **69(1)** (2020), 528–539.
12. S. K. TIWARI, R. K. SHARMA and B. DHARA: *Some theorems of commutativity on semiprime rings with mappings*. *Southeast Asian Bull. Math* **42(2)** (2018), 279–292.
13. S. HUANG: *Generalized reverse derivations and commutativity of prime rings*. *Communications in Mathematics* **27** (2019), 43–50.
14. Y. AHMED and M. ASLAM: *On diversity of generalized reverse derivations in rings*. *Journal of Mechanics of Continua and Mathematical Sciences* **15(6)** (2020), 118–126.
15. Z. Z. M. ALHAIDARY and A. H. MAJEED: *Square closed Lie ideals and multiplicative (generalized) (α, β) reverse derivation of prime rings*. *Journal of Discrete Mathematical Sciences & Cryptography* **24(7)** (2021), 2037–2046.
16. Z. Z. M. ALHAIDARY and A. H. MAJEED: *Commutativity results for multiplicative (generalized) (α, β) reverse derivations on prime rings*. *Iraqi Journal of Science* **62(9)** (2021), 3102–3113.