

ON CONFORMALLY FLAT p -POWER (α, β) -METRICS

Mosayeb Zohrehvand¹, Shahroud Azami²
and Ghodratalah Fasihi-Ramandi²

¹Faculty of Mathematical Sciences and Statistics, Department of Mathematics
Malayer University, P. O. Box 65719-95863, Malayer, Iran

²Faculty of Sciences, Department of pure Mathematics
Imam Khomeini International University, P. O. Box 65719-95863, Gazvin, Iran

ORCID IDs: Mosayeb Zohrehvand  <https://orcid.org/0000-0002-3876-4060>
Shahroud Azami  <https://orcid.org/0000-0002-8976-2014>
Ghodratallah Fasihi-Ramandi  <https://orcid.org/0009-0000-1452-9376>

Abstract. The purpose of this paper is to study the class of conformally flat p -power (α, β) -metrics $F = \alpha(1 + \frac{\beta}{\alpha})^p$, where $p \neq 0$ is a constant. This metric is interesting, because for $p = -1, \frac{1}{2}, 1, 2$ it reduces to the Matsumoto, square-root, Randers and square metric, respectively. We prove that if a p -power (α, β) -metric has relatively isotropic mean Landsberg curvature, then it is either a Riemannian metric or a locally Minkowski metric.

Keywords: metrics, manifold, conformal geometry.

1. Introduction

Conformal geometry has many important and interesting applications in physical theories, which have led to increased attention and research. In general relativity, light-like geodesics remain invariant under the conformal relation between pseudo-Riemannian metrics. The Weyl theorem states that the study of conformal and projective properties of a Finsler metric characterizes the metric properties as uniquely [13, 19].

Two Finsler metrics F and $\tilde{F}$, defined on a differentiable manifold M , are said to be conformally related if $F = e^{\kappa(x)}\tilde{F}$, where $\kappa(x)$ is a scalar function on M and

Received: February 28, 2024, revised: March 31, 2024, accepted: March 31, 2024

Communicated by Uday Chand De

Corresponding Author: M. Zohrehvand

E-mail addresses: m.zohrehvand@malayeru.ac.ir (M. Zohrehvand), azami@sci.ikiu.ac.ir (S. Azami), fasihi@sci.ikiu.ac.ir (G. Fasihi-Ramandi)

2020 *Mathematics Subject Classification.* Primary 53B40; Secondary 53C60

is referred to as the conformal factor. If $\tilde{F}$ is a locally Minkowski metric, we say that F is a conformally flat metric.

S. Kikuchi found a Finsler connection that is conformally invariant and he expressed the conformal flatness of a Finsler metric in terms of this connection [12]. Based on Kikuchi's idea, M. Matsumoto introduced a conformal invariant Finsler connection for Finsler metrics that have a tensor satisfying in a certain condition, weaker than Kikuchi's condition, then the condition that a Finsler metric is conformally flat stated by the terms of this connection [16].

Ichijyō and Hashiguchi proposed a condition that a Randers metric can be conformally flat [10]. Randers metrics are the simplest examples of (α, β) -metrics which made an important class of Finsler metrics. They have wide-ranging applications in physics, biology, etc (see [2]).

A Finsler metric that is represented as $F = \alpha\phi(s)$, $s := \beta/\alpha$ is called a (α, β) -metric where $\alpha(x, y) := \sqrt{a_{ij}(x)y^i y^j}$ is a Riemannian metric, $\beta(x, y) := b_i(x)y^i$ is a 1-form and $\phi(s)$ is a C^∞ function that satisfies a certain inequality [15]. The one-form β can be considered as an external force like wind and current. Thus (α, β) -metrics can express the geometry of a Riemannian space impacted by an external force. These metrics have been extensively studied because they are computable and the research on (α, β) -metrics have contributed significantly to the field of Finsler geometry and have suggested several ideas for further studies.

L. Kang considered conformally flat Randers metrics of scalar flag curvature and proved that they are projectively flat and also provided a complete classification of such metrics [11]. Conformally flat (α, β) -metrics with isotropic S-curvature are considered in [3]. It is shown that they are Riemannian or locally Minkowski metric and also classified conformally flat weak Einstein (α, β) -metrics of polynomial type. In [5] it is proved that every non-Riemannian conformally flat weakly Landsberg (α, β) -metric must be a locally Minkowski metric. Chen et al. studied conformally flat (α, β) -metrics with constant flag curvature. Their research showed that these metrics are either locally Minkowski or Riemannian metrics [4]. Tayebi and Razgordani studied conformally flat weak Einstein fourth root (α, β) -metrics and proved that they are also either locally Minkowskian or Riemannian [23]. For more references, refer to [1, 17, 21, 22]

The class of p -power (α, β) -metrics are of the form

$$F = \alpha \left(1 + \frac{\beta}{\alpha} \right)^p,$$

where $p \neq 0$ is a constant. For $p = 1$, F is a Randers metric. The Randers metrics were introduced by G. Randers, when he was studying general relativity [18]. Later on, R. S. Ingarden applied this metric to the theory of the electron microscope [2]. If $p = -1$ then $F = \alpha^2/(\beta + \alpha)$ is a Matsumoto metric. This metric was introduced by M. Matsumoto and is also called the slope metric. As a geometrical motivation for the Matsumoto metric, suppose that a person is walking with the constant speed v on a surface S that has the angle θ with respect to sea level and under the gravitational field g . We can embed S in the Euclidean space $\mathbb{E}^3$ using

the parametrization

$$(x, y) \longmapsto (x, y, f(x, y)),$$

where $f(x, y)$ is a smooth function. By use of Okubo's method [2], we obtain the Matsumoto metric

$$F(x, y, \dot{x}, \dot{y}) = \frac{\alpha^2}{v\alpha - \frac{g}{2}\beta},$$

where

$$\begin{aligned}\alpha &:= \sqrt{(1 + f_x^2)\dot{x}^2 + 2f_x f_y \dot{x}\dot{y} + (1 + f_y^2)\dot{y}^2}, \\ \beta &:= f_x \dot{x} + f_y \dot{y}.\end{aligned}$$

In order to obtain the usual form of the Matsumoto metric, we set $v = -\frac{g}{2}$ and in this case, we have $F = \frac{\alpha^2}{\alpha + \beta}$. For more details see [20].

Also, in the case of $p = \frac{1}{2}$, $F = \sqrt{\alpha(\alpha + \beta)}$ that is called square-root (α, β) -metric. In [24] p -power (α, β) -metrics of Einstein-reversible type have been considered and the local structure of a two-dimensional square-root metric is determined. Thus, the class of p -power (α, β) -metrics deserve more attention.

For a Finsler metric F , we have the basic tensors, fundamental tensor $\mathbf{g}_y$ and Cartan torsion $\mathbf{C}$. By taking horizontal covariant derivative of Cartan torsion along the geodesics we obtain the tensor field $\mathbf{L}$ that is called Landsberg curvature. The trace of $\mathbf{C}$ and $\mathbf{L}$ are called the mean Cartan torsion $\mathbf{I}$ and the mean Landsberg curvature $\mathbf{J}$, respectively. A Finsler metric F is called relatively isotropic mean Landsberg curvature if there exists a scalar function $c = c(x)$ on M such that

$$(1.1) \quad \mathbf{J} + cF\mathbf{I} = 0.$$

As to find an explicit example of conformally flat Finsler metric, this paper is devoted to the study of the conformally flat p -power (α, β) -metric, that has relatively isotropic mean Landsberg curvature. The main result is the following.

Theorem 1.1. *Let $F = \alpha(1 + \frac{\beta}{\alpha})^p$ be the conformally flat (α, β) -metric on a differentiable manifold M of dimension $n \geq 3$, where $p \neq 0$ is a real constant. Suppose that F has relatively isotropic mean Landsberg curvature. Then F reduces to a Riemannian metric or a locally Minkowski metric.*

As the consequences of Theorem 1.1, we have the following conclusions, that retrieve other researcher's results.

Corollary 1.1. *Let $F = \alpha + \beta$ be the conformally flat Randers metric on a differentiable manifold M of the dimension $n \geq 3$. Suppose that F has relatively isotropic mean Landsberg curvature. Then F is a Riemannian metric or a locally Minkowski metric.*

It would be noted that, for $F = \alpha + \beta$, it is proved in [6] that every Randers metric is of isotropic mean Landsberg curvature if and only if it is of isotropic S-curvature. On the other hand every conformally flat Randers metric with isotropic S-curvature is either a locally Minkowski or a Riemannian metric [11]. Thus, using these results, we have additional proof for Corollary 1.1.

Corollary 1.2. *Let $F = \sqrt{\alpha(\alpha + \beta)}$ be the conformally flat square-root (α, β) -metric on a differentiable manifold M of the dimension $n \geq 3$. Suppose that F has relatively isotropic mean Landsberg curvature. Then F is a Riemannian metric or a locally Minkowski metric.*

In [17] the conformally flat square-root (α, β) -metrics of relatively isotropic mean Landsberg curvature is studied and Corollary 1.2 is proved.

Corollary 1.3. *Let $F = \frac{\alpha^2}{\alpha + \beta}$ be the conformally flat Matsumoto metric on a differentiable manifold M of the dimension $n \geq 3$. Suppose that F has relatively isotropic mean Landsberg curvature. Then F is a Riemannian metric or a locally Minkowski metric.*

2. Preliminaries

Let $F = F(x, y)$ be a Finsler metric on an n -dimensional differentiable manifold M and $TM_0 := \bigcup_{x \in M} T_x M - \{0\}$ the slit tangent bundle. The fundamental tensor $(\mathbf{g}_y) = (g_{ij}(x, y))$ of F is a quadratic form on $T_x M$ that is defined

$$g_{ij}(x, y) := \frac{1}{2} [F^2]_{y^i y^j}(x, y).$$

A curve $x = x^i(t)$ on Finsler space (M, F) is called geodesic if satisfies in the following system of ODEs:

$$\frac{d^2 x^i}{dt^2} + G^i(x, \frac{dx}{dt}) = 0,$$

where $G^i = G^i(x, y)$ are called the geodesic coefficients of F and defined by

$$G^i = \frac{1}{4} g^{il} \left\{ [F^2]_{x^m y^l} y^m - [F^2]_{x^l} \right\}.$$

In Finsler geometry, there are some geometric quantities that are vanishing for Riemannian metrics and are called non-Riemannian quantities. The Cartan torsion $\mathbf{C}$ is a symmetric trilinear form $\mathbf{C} := C_{ijk} dx^i \otimes dx^j \otimes dx^k$ on TM_0 that is defined as follow

$$C_{ijk} := \frac{1}{4} [F^2]_{y^i y^j y^k} = \frac{1}{2} \frac{\partial g_{ij}}{\partial y^k}.$$

One can see that F is a Riemannian metric if and only if $\mathbf{C} = 0$. Thus it is a non-Riemannian quantity.

The mean Cartan torsion of F is the tensor field $\mathbf{I} := I_i dx^i$, that is defined by

$$I_i := g^{jk} C_{ijk}.$$

Furthermore, one can see that

$$I_i = \frac{\partial}{\partial y^i} \left[\ln \sqrt{\det(g_{jk})} \right]$$

The horizontal covariant derivative of Cartan torsion along the geodesics defines the tensor field $\mathbf{L} := L_{ijk} dx^i \otimes dx^j \otimes dx^k$ on the slit tangent bundle TM_0 , that is called the Landsberg curvature of F . Thus $L_{ijk} := C_{ijk;m} y^m$, where ";" denoted the horizontal covariant derivative with respect to the Berwald connection of F . Also, the Landsberg curvature can be expressed as the following

$$(2.1) \quad L_{ijk} = -\frac{1}{2} F F_{y^m} [G^m]_{y^i y^j y^k}.$$

A Finsler metric F is called the Landsberg metric if $\mathbf{L} = 0$.

The mean Landsberg curvature $\mathbf{J} := J_i dx^i$ is a non-Riemannian quantity that is obtained by horizontal covariant derivative of the mean Cartan torsion $\mathbf{I}$ along the geodesics of F . Thus

$$(2.2) \quad J_i := I_{i;m} y^m.$$

Also, the mean Landsberg curvature $\mathbf{J}$ can be obtained as following

$$J_i := g^{jk} L_{ijk}.$$

A Finsler metric F is called weak Landsberg metric if $\mathbf{J} = 0$.

A Finsler metric F is called of relatively isotropic mean Landsberg curvature if $\mathbf{J}/\mathbf{I}$, the relative growth rate of the mean Cartan torsion along geodesics of F is isotropic, i.e. there exists a scalar function $c = c(x)$ on M such that

$$\mathbf{J} + cF\mathbf{I} = 0.$$

A Finsler metric F is an (α, β) -metric if $F = \alpha\phi(s)$, $s := \beta/\alpha$, where $\alpha = \sqrt{a_{ij}(x)y^i y^j}$ is a Riemannian metric, $\beta = b_i(x)y^i$ is a 1-form with $\|\beta_x\| < b_0$, $x \in M$ and $\phi(s)$ is a positive C^∞ function on $(-b_0, b_0)$ satisfying

$$(2.3) \quad \phi(s) - s\phi'(s) + (b^2 - s^2)\phi''(s) > 0, \quad |s| \leq b < b_0.$$

In this case, the metric $F = \alpha\phi(s)$ is a positive definite Finsler metric [9]. The fundamental tensor $F = \alpha\phi(s)$ is given by

$$g_{ij} = \rho a_{ij} + \rho_0 b_i b_j + \rho_1 (b_i \alpha_i + b_j \alpha_j) + \rho_2 \alpha_i \alpha_j,$$

where $\alpha_i := \alpha^{-1} a_{ij} y^j$, and

$$\begin{aligned} \rho &:= \phi(\phi - s\phi'), & \rho_0 &:= \phi\phi'' + \phi'\phi', \\ \rho_1 &:= -s(\phi\phi'' + \phi'\phi') + \phi\phi', & \rho_2 &:= s\{s(\phi\phi'' + \phi'\phi') - \phi\phi'\}. \end{aligned}$$

Let

$$r_{ij} := \frac{1}{2}(b_{i|j} + b_{j|i}), \quad s_{ij} := \frac{1}{2}(b_{i|j} - b_{j|i})$$

where $b_{i|j}$ denote the coefficients of the covariant derivative of β with respect to α . Let's denote

$$\begin{aligned} r_j^i &:= a^{im} r_{mj}, & r_{00} &:= r_{ij} y^i y^j, & r_i &:= b^m r_{mi}, \\ r_0 &:= r_i y^i, & r_{i0} &:= r_{im} y^m, & s_j^i &:= a^{im} s_{mj}, \\ s_{i0} &:= s_{im} y^m, & s_i &:= b^m s_{mi}, & s_0 &:= s_i y^i, \end{aligned}$$

The geodesic coefficients G^i of an (α, β) -metric $F = \alpha\phi(s)$ are given by

$$(2.4) \quad G^i = G_\alpha^i + \alpha Q s_0^i + \{-2Q\alpha s_0 + r_{00}\} \left\{ \Psi b^i + \Theta \alpha^{-1} y^i \right\},$$

where G_α^i are the geodesic coefficients of α and

$$\begin{aligned} Q &:= \frac{\phi'}{\phi - s\phi'}, \\ \Theta &:= \frac{\phi\phi' - s(\phi\phi'' + \phi'\phi')}{2\phi[\phi - s\phi' + (b^2 - s^2)\phi'']}, \\ \Psi &:= \frac{\phi''}{2[\phi - s\phi' + (b^2 - s^2)\phi'']}. \end{aligned}$$

For more details, see [9].

3. Proof of Theorem 1.1

In this section, we focus on a class of (α, β) -metrics, that contain Randers metric, square metric, square-root metric, Matsumoto metric, etc. This class of metrics is called p -power (α, β) -metrics and is of the form

$$F = \alpha(1 + s)^p, \quad s := \beta/\alpha,$$

where $p \neq 0$ is a real constant. For more details see [24].

We consider p -power (α, β) -metric with relatively isotropic mean Landsberg curvature and prove Theorem 1.1. In [8], the mean Cartan torsion of an (α, β) -metric is computed.

Lemma 3.1. ([8]) *For an (α, β) -metric $F = \alpha\phi(s)$, $s = \beta/\alpha$, the mean Cartan torsion is given by*

$$(3.1) \quad I_i = -\frac{1}{2F} \frac{\Phi}{\Delta} (\phi - s\phi') h_i,$$

where

$$\begin{aligned}\Delta &:= 1 + sQ + (b^2 - s^2)Q', \\ \Phi &:= -(n\Delta + 1 + sQ)(Q - sQ') - (b^2 - s^2)(1 + sQ)Q'', \\ \Psi_1 &:= \sqrt{b^2 - s^2}\Delta^{\frac{1}{2}} \left[\frac{\sqrt{b^2 - s^2}\Phi}{\Delta^{\frac{3}{2}}} \right]', \\ h_j &:= b_j - \alpha^{-1}sy_j.\end{aligned}$$

It is well known that, by Deicke's theorem, F is a Riemannian metric if and only if $\mathbf{I} = 0$. Thus from (3.1) we have

Lemma 3.2. *An (α, β) -metric $F = \alpha\phi(s)$, $s := \beta/\alpha$ is a Riemannian metric if and only if $\Phi = 0$.*

From (2.2) and (3.1), one can see that the mean Landsberg curvature of an (α, β) -metric $F = \alpha\phi(s)$, $s = \beta/\alpha$, is given by

$$\begin{aligned}J_j &= \frac{1}{2\alpha^4\Delta} \left\{ \frac{2\alpha^3}{b^2 - s^2} \left[\frac{\Phi}{\Delta} + (n+1)(Q - sQ') \right] (s_0 + r_0)h_j \right. \\ &\quad + \frac{\alpha^2}{b^2 - s^2} \left[\Psi_1 + s\frac{\Phi}{\Delta} \right] (r_{00} - 2\alpha Qs_0)h_j \\ &\quad + \alpha \left[-\alpha^2 Q' s_0 h_j + \alpha Q(\alpha^2 s_j - y_j s_0) + \alpha^2 \Delta s_{j0} \right. \\ &\quad \left. \left. + \alpha^2 (r_{j0} - 2\alpha Qs_j) - (r_{00} - 2\alpha Qs_0)y_j \right] \frac{\Phi}{\Delta} \right\},\end{aligned}\tag{3.2}$$

where $y_j := a_{ij}y^i$. For more details see [8, 14]. From (3.1) and (3.2), we obtain

$$\begin{aligned}J_j + c(x)FI_j &= -\frac{1}{2\alpha^4\Delta} \left\{ \frac{2\alpha^3}{b^2 - s^2} \left[\frac{\Phi}{\Delta} + (n+1)(Q - sQ') \right] (s_0 + r_0)h_j \right. \\ &\quad + \frac{\alpha^2}{b^2 - s^2} \left[\Psi_1 + s\frac{\Phi}{\Delta} \right] (r_{00} - 2\alpha Qs_0)h_j + \alpha \left[-\alpha^2 Q' s_0 h_j \right. \\ &\quad + \alpha Q(\alpha^2 s_j - y_j s_0) + \alpha^2 \Delta s_{j0} + \alpha^2 (r_{j0} - 2\alpha Qs_j) \\ &\quad \left. \left. - (r_{00} - 2\alpha Qs_0)y_j \right] \frac{\Phi}{\Delta} + c(x)\alpha^4\Phi(\phi - s\phi')h_j \right\}.\end{aligned}\tag{3.3}$$

Since we study conformally flat (α, β) -metrics, we need the following Lemma.

Lemma 3.3. ([2]) *Let $F = \alpha\phi(s)$, $s = \beta/\alpha$, be an (α, β) -metric. Then F is locally Minkowski metric if and only if α is flat and β is parallel with respect to α .*

Now, let $F = \alpha\phi(s)$, $s = \beta/\alpha$, be a conformally flat Finsler metric, it means that, there exists a Minkowski metric $\tilde{F}$ such that $\tilde{F} = e^{\kappa(x)}F$, where $\kappa(x)$ is a scalar function on the manifold. Since $F = \alpha\phi(\beta/\alpha)$, we deduce $\tilde{F} = \tilde{\alpha}\phi(\tilde{\beta}/\tilde{\alpha})$ is an (α, β) -metric, where

$$(3.4) \quad \tilde{\alpha} = e^{\kappa(x)}\alpha, \quad \tilde{\beta} = e^{\kappa(x)}\beta.$$

From (3.4), we have

$$\tilde{a}_{ij} = e^{2\kappa(x)}a_{ij}, \quad \tilde{b}_i = e^{\kappa(x)}b_i.$$

The Christoffel symbols Γ_{jk}^i of α and the Christoffel symbols $\tilde{\Gamma}_{jk}^i$ of $\tilde{\alpha}$ are related by

$$\tilde{\Gamma}_{jk}^i = \Gamma_{jk}^i + \delta_j^i \kappa_k + \delta_k^i \kappa_j - \kappa^i a_{jk},$$

where $\kappa_i := \frac{\partial \kappa}{\partial x^i}$ and $\kappa^i := a^{ij}\kappa_j$ [7]. Thus, we obtain

$$(3.5) \quad \tilde{b}_{i||j} = \frac{\partial \tilde{b}_i}{\partial x^j} - \tilde{b}_s \tilde{\Gamma}_{jk}^i = e^\kappa (b_{i||j} - b_j \kappa_i + b_r \kappa^r a_{ij}).$$

where $\tilde{b}_{i||j}$ denote the coefficients of the covariant derivative of $\tilde{\beta}$ with respect to $\tilde{\alpha}$.

Since $\tilde{F}$ is a Minkowski metric, from Lemma 3.3, we have $\tilde{b}_{i||j} = 0$. Thus

$$(3.6) \quad b_{i||j} = b_j \kappa_i - b_r \kappa^r a_{ij}.$$

From (3.6), we conclude

$$(3.7) \quad r_{ij} = \frac{1}{2}(\kappa_i b_j + \kappa_j b_i) - b_r \kappa^r a_{ij}, \quad r_j = -\frac{1}{2}(b^r \kappa_r) b_j + \frac{1}{2} \kappa_j b^2,$$

$$(3.8) \quad r_{i0} = \frac{1}{2}[\kappa_i \beta + (\kappa_r y^r) b_i] - \kappa_r b^r y_i, \quad s_{ij} = \frac{1}{2}(\kappa_i b_j - \kappa_j b_i),$$

$$(3.9) \quad s_j = \frac{1}{2}(b^r \kappa_r) b_j - \kappa_j b^2, \quad s_{i0} = \frac{1}{2}[\kappa_i \beta - (\kappa_r y^r) b_i].$$

Further, we have

$$(3.10) \quad r_{00} = (\kappa_r y^r) \beta - (\kappa_r b^r) \alpha^2,$$

$$(3.11) \quad r_0 = \frac{1}{2}(\kappa_r y^r) b^2 - \frac{1}{2}(\kappa_r b^r) \beta,$$

$$(3.12) \quad s_0 = \frac{1}{2}(\kappa_r b^r) \beta - \frac{1}{2}(\kappa_r y^r) b^2.$$

From (3.11) and (3.12), we see that a conformally flat (α, β) -metric satisfies $r_0 + s_0 = 0$ which means that the 1-form β has constant length with respect to α .

In order to simplify the computations, we take an orthonormal basis at any point x with respect to α such that $\alpha = \sqrt{\sum_{i=1}^n (y^i)^2}$ and $\beta = b y^1$, where $b := \|\beta_x\|_\alpha$. Then, we take the following coordinate transformation

$$\psi : (s, u^A) \longrightarrow (y^i),$$

in $T_x M$, that is

$$(3.13) \quad y^1 = \frac{s}{\sqrt{b^2 - s^2}} \bar{\alpha}, \quad y^A = u^A, \quad 2 \leq A \leq n,$$

where $\bar{\alpha} = \sqrt{\sum_{i=2}^n (u^i)^2}$. In this case, we have

$$(3.14) \quad \alpha = \frac{b}{\sqrt{b^2 - s^2}} \bar{\alpha}, \quad \beta = \frac{bs}{\sqrt{b^2 - s^2}} \bar{\alpha}.$$

Then, by (3.6)-(3.14) one can obtain

$$(3.15) \quad r_{00} = -b\kappa_1 \bar{\alpha}^2 + \frac{bs\bar{\kappa}_0 \bar{\alpha}}{\sqrt{b^2 - s^2}}, \quad r_0 = -s_0 = \frac{1}{2} b^2 \bar{\kappa}_0,$$

$$(3.16) \quad r_{A0} = \frac{1}{2} \frac{\kappa_A bs \bar{\alpha}}{\sqrt{b^2 - s^2}} - (b\kappa_1) u_A, \quad r_{10} = \frac{1}{2} b \bar{\kappa}_0,$$

$$(3.17) \quad s_A = -\frac{1}{2} \kappa_A b^2, \quad s_1 = 0,$$

$$(3.18) \quad s_{A0} = \frac{1}{2} \frac{\kappa_A bs \bar{\alpha}}{\sqrt{b^2 - s^2}}, \quad s_{10} = -\frac{1}{2} b \bar{\kappa}_0,$$

$$(3.19) \quad h_A = -\frac{\sqrt{b^2 - s^2} s u_A}{b \bar{\alpha}}, \quad h_1 = b - \frac{s^2}{b}.$$

where $\bar{\kappa}_0 := \kappa_A u^A$.

Now, we can prove Theorem 1.1.

Proof of Theorem 1.1: Since $\tilde{b}_{i||j} = 0$, we have that $\tilde{b}$ is a real constant. If $\tilde{b} = 0$, then $F = e^{k(x)} \tilde{\alpha}$ is a Riemannian metric. Now, let F is not Riemannian. Suppose that F is a conformally flat (α, β) -metric with relatively isotropic mean Landsberg curvature. By (3.3) and $r_0 + s_0 = 0$, we obtain

$$(3.20) \quad \begin{aligned} & \frac{\alpha^2}{b^2 - s^2} \left\{ \Psi_1 + s \frac{\Phi}{\Delta} \right\} (r_{00} - 2\alpha Q s_0) h_j + \alpha \left\{ -\alpha^2 Q' s_0 h_j \right. \\ & \quad \left. + \alpha Q (\alpha^2 s_j - y_j s_0) + \alpha^2 \Delta s_{j0} + \alpha^2 (r_{j0} - 2\alpha Q s_j) \right. \\ & \quad \left. - (r_{00} - 2\alpha Q s_0) y_j \right\} \frac{\Phi}{\Delta} - c(x) \alpha^4 \Phi (\phi - s\phi') h_j = 0. \end{aligned}$$

Putting $j = 1$ in (3.20), we have

$$(3.21) \quad \begin{aligned} & \frac{\alpha^2}{b^2 - s^2} \left\{ \Psi_1 + s \frac{\Phi}{\Delta} \right\} (r_{00} - 2\alpha Q s_0) h_1 + \alpha \left\{ -\alpha^2 Q' s_0 h_1 \right. \\ & \quad \left. + \alpha Q (\alpha^2 s_1 - y_1 s_0) + \alpha^2 \Delta s_{10} + \alpha^2 (r_{10} - 2\alpha Q s_1) \right. \\ & \quad \left. - (r_{00} - 2\alpha Q s_0) y_1 \right\} \frac{\Phi}{\Delta} - c(x) \alpha^4 \Phi (\phi - s\phi') h_1 = 0. \end{aligned}$$

Substituting (3.14)-(3.19) into (3.21) and then multiplying the resulting equation with $-2\Delta(b^2 - s^2)^{3/2}$ we have

$$b^2 \bar{\alpha}^3 \left\{ 2\sqrt{b^2 - s^2} \Delta [bc(x) \Phi (\phi - s\phi') + \Psi_1 \sigma_1] \bar{\alpha} - \bar{\kappa}_0 [b^2 \Phi Q' (b^2 - s^2) \right.$$

$$(3.22) \quad \left. + \Phi b^2(sQ + 1) - \Delta \Phi b^2 - 2\Psi_1 \Delta(b^2Q + s) \right\} = 0.$$

From (3.22), we get

$$(3.23) \quad \Delta [bc(x)\Phi(\phi - s\phi') + \Psi_1 \kappa_1] = 0,$$

$$(3.24) \quad \bar{\kappa}_0 [b^2 \Phi Q'(b^2 - s^2) + \Phi b^2(sQ + 1) - \Delta \Phi b^2 - 2\Psi_1 \Delta(b^2Q + s)] = 0.$$

Since $\Delta = Q'(b^2 - s^2) + sQ + 1$, one can see that (3.24) simplify as follow

$$(b^2 \Psi_1 \Delta Q + \Psi_1 \Delta s) \bar{\kappa}_0 = 0.$$

This means that

$$(3.25) \quad \Psi_1 \Delta(b^2Q + s) \bar{\kappa}_0 = 0.$$

Now let $j = A$ in (3.20), thus we have

$$(3.26) \quad \begin{aligned} & \frac{\alpha^2}{b^2 - s^2} [\Psi_1 + s \frac{\Phi}{\Delta}] (r_{00} - 2\alpha Q s_0) h_A + \alpha [-\alpha^2 Q' s_0 h_A + \\ & \alpha Q (\alpha^2 s_A - y_A s_0) + \alpha^2 \Delta s_{A0} + \alpha^2 (r_{A0} - 2\alpha Q s_A) \\ & - (r_{00} - 2\alpha Q s_0) y_A] \frac{\Phi}{\Delta} + c(x) \alpha^4 \Phi(\phi - s\phi') h_A = 0. \end{aligned}$$

Putting (3.14)-(3.19) into (3.26) and using the same method as used in the case of $j = 1$ and from $\Delta = Q'(b^2 - s^2) + sQ + 1$, we get

$$(3.27) \quad \begin{aligned} & (s\Delta + s + b^2Q) b^2 \Phi \kappa_A \bar{\alpha}^2 - [(s\Delta + s + b^2Q) b^2 \Phi \\ & + 2s(b^2Q + s) \Psi_1 \Delta] \bar{\kappa}_0 u_A = 0, \end{aligned}$$

$$(3.28) \quad s\sqrt{b^2 - s^2} [\Phi bc(x)(\phi - s\phi') + \Psi_1 \kappa_1] \Delta u_A = 0.$$

One can easily see that (3.28) is equivalent to (3.23). Also, multiplying (3.27) with u^A implies that

$$(3.29) \quad s(b^2Q + s) \Psi_1 \Delta \bar{\kappa}_0 \bar{\alpha}^2 = 0.$$

It is obvious that (3.29) is equivalent to (3.25). Anyway, we showed that a conformally flat (α, β) -metric with relatively isotropic mean Landsberg curvature satisfy (3.23) and (3.25).

If $b^2Q + s = 0$, then we obtain $\phi = k\sqrt{b^2 - s^2}$, where k is a constant. This is a contradiction with the assumption that $\phi = (1 + s)^p$. Thus $b^2Q + s \neq 0$ and then from (3.25) we conclude that $\Psi_1 = 0$ or $\kappa_A = 0$.

If $\Psi_1 = 0$, then using (3.23) we obtain that $\Phi = 0$, and from Lemma 3.2, we see that F is a Riemannian metric.

If $\Psi_1 \neq 0$, then $\kappa_A = 0$. In this case, we prove that $\kappa_1 = 0$. Simplifying (3.23) and multiplying it by Δ^2 , we get

$$(3.30) \quad \left\{ [(b^2 - s^2)\Phi' - s\Phi] \Delta - \frac{3}{2}(b^2 - s^2)\Phi \Delta' \right\} \kappa_1 - bc(x) \Delta^2 \Phi(\phi - s\phi') = 0.$$

Let $A_1 := (p-1)s - 1$ and $A_2 := (1-p^2)s^2 + (2-p)s + p(p-1)b^2 + 1$. Putting $\phi(s) = (1+s)^p$ into (3.30) and multiplying by $A_1^4 A_2^2$ and using Maple program, we obtain

$$(3.31) \quad A_1 A_2^2 E c(x) b p (1+s)^{p-1} + \mathcal{M}_7 s^7 + \dots + \mathcal{M}_0 = 0,$$

where

$$(3.32) \quad \begin{aligned} E := & 2n(p+1)(p-1)^2 s^3 + (p-1)[(p-5)n-3]s^2 \\ & - \{2(np^3 - (2n+1)p^2 + (n+2)p-1)b^2 + (n+1)(3p-4)\}s \\ & + [(n+2)p^2 - (n+4)p+2]b^2 + (1+n), \end{aligned}$$

and $\mathcal{M}_i$, $(0 \leq i \leq 7)$ are polynomials independent from s . Specially

$$(3.33) \quad \mathcal{M}_7 := 3(n-1)p(p+1)(p-1)^4 \kappa_1.$$

Now, we consider the following cases.

Case(I): If $p \geq 2$, is a positive integer constant. Then (3.31) can be rewritten as

$$(3.34) \quad \mathcal{M}'_{7+p} s^{7+p} + \mathcal{M}'_{6+p} s^{6+p} + \dots + \mathcal{M}'_0 = 0.$$

where $\mathcal{M}'_{7+p} := 2nbp(p+1)^3(p-1)^5 c(x)$. Thus $c(x) = 0$ and therefore (3.34) is reduced to

$$(3.35) \quad \mathcal{M}_7 s^7 + \dots + \mathcal{M}_0 = 0.$$

From (3.35) and (3.33) we obtain that $\kappa_1 = 0$.

Case(II): If $p = 1$, then $F = \alpha + \beta$ is a Randers metric. In this case (3.31) is reduced to

$$(3.36) \quad (\kappa_1 - 2bc(x))s^2 + 2(\kappa_1 - 2bc(x))s + b(\kappa_1 b - 2c(x)) = 0.$$

From (3.36) it follows that $c(x) = \kappa_1 = 0$.

Case (III): If p is a positive non-integer constant. Then, from (3.31) and (3.32), we have that $c(x) = \mathcal{M}_i = 0$, $(0 \leq i \leq 7)$. Thus from (3.33), it follows that $\kappa_1 = 0$.

Case (IV): If $p = -1$. In this case $F = \frac{\alpha^2}{\alpha+\beta}$, is a Matsumoto metric. Putting $\phi(s) = \frac{1}{1+s}$ in (3.30) and multiplying by $A_1^4 A_2^2 (1+s)^2$, we get

$$18(2n+1)\kappa_1 s^8 + \mathcal{D}_7 s^6 + \dots + \mathcal{D}_0 = 0.$$

Thus $\kappa_1 = 0$.

Case (V): If $p \leq -2$ is a negative integer constant. In this case, Eq. (3.31) can be rewritten as follows.

$$(3.37) \quad \mathcal{N}_{8-p} s^{8-p} + \mathcal{N}_{7-p} s^{7-p} + \dots + \mathcal{N}_1 s + \mathcal{N}_0 = 0,$$

where $\mathcal{N}_{8-p} = 3(n-1)p(p+1)(p-1)^4\kappa_1$. Thus $\kappa_1 = 0$.

Case (VI): If p is a negative non-integer constant. In this case, (3.31) is reduced to

$$(3.38) \quad A_1 A_2^2 E c(x) b p + (1+s)^{1-p} [\mathcal{M}_7 s^7 + \dots + \mathcal{M}_0] = 0.$$

From (3.33), and (3.38) we have $\kappa_1 = c(x) = 0$.

Therefore, in any case $\kappa_1 = \kappa_A = 0$, which means that κ is a constant. Thus F is a locally Minkowski metric. This completes the proof. $\square$

It would be noted that, in the Case (I), $\phi(s) = (1+s)^p$ is a polynomials of s . In [5] it is proved that every (α, β) -metric $F = \alpha\phi(s)$ with relatively isotropic mean Landsberg curvature, where $\phi(s)$ is a polynomial of s , is either a Riemannian or a locally Minkowski metric.

4. Conclusion

In this paper, we considered conformally flat p -power (α, β) -metrics. We proved that every conformally flat p -power (α, β) -metrics with relatively isotropic mean Landsberg curvature is a Riemannian or a locally Minkowski metric. Therefore, this class of (α, β) -metrics are classified. The readers are encouraged to consider the class of conformally flat p -power (α, β) -metrics with other Riemannian or non-Riemannian quantities, such as weakly isotropic scalar flag curvature, weak Einstein metric, etc.

REFERENCES

1. M. AMINI: *On conformally flat cubic (α, β) -metrics*. J. Finsler Geom. Appl. **2** (2021), 75–85.
2. P. L. ANTONELLI and R. S. INGARDEN and M. MATSUMOTO: *The theory of sprays and Finsler spaces with applications in physics and biology*. FTPH 58, Kluwer Academic Publishers, 1993.
3. G. CHEN, and X. CHENG: *An important class of conformally flat weak Einstein Finsler metrics*. Int. J. Math. **24** (2013), 15 pages.
4. G. CHEN and Q. HE and Z. SHEN: *On conformally flat (α, β) -metrics with constant flag curvature*. Publ. Math. Debrecen **86** (2015), 351–363.
5. X. CHENG and H. LI and Y. ZOU: *On conformally flat (α, β) -metrics with relatively isotropic mean Landsberg curvature*. Publ. Math. Debrecen **85** (2014), 131–144.
6. X. CHENG and Z. SHEN: *Randers metrics with special curvature properties*. Osaka J Math **40** (2003), 87–101.
7. X. CHENG and Z. SHEN: *Finsler Geometry: An Approach via Randers Spaces*. SpringerLink : Bücher, Springer Berlin Heidelberg, 2013.
8. X. CHENG and H. WANG and M. WANG: *(α, β) -metrics with relatively isotropic mean Landsberg curvature*. Publ. Math. Debrecen **72** (2008), 475–485.
9. S. S. CHERN and Z. SHEN: *Riemann-Finsler Geometry*. Singapore, World Scientific, 2005.

10. Y. ICHIJYO and M. HASHUIGUCHI: *On the condition that a Randers space be conformally flat*. Rep. Fac. Sci. Kagoshima Univ. **22** (1989), 7–14.
11. L. Kang: *On conformally flat Randers metrics*. Sci. Sin. Math. **41** (2011), 439–446.
12. S. KIKUCHI: *On the condition that a Finsler space be conformally flat*. Tensor (N.S.) **55** (1994), 97–100.
13. M. S. KNEBELMAN: *Conformal geometry of generalised metric spaces*. Proc. Natl. Acad. Sci. USA **15** (1929), 376–379.
14. B. LI and Z. SHEN: *On a class of weak Landsberg metrics*. Science in China Series A **50** (2007), 75–85.
15. M. MATSUMOTO: *Projective flat Finsler spaces with (α, β) -metric*. Rep. Math. Phys. **30** (1991), 15–20.
16. M. MATSUMOTO: *Conformally Berwald and conformally flat Finsler spaces*. Publ. Math. Debrecen **58** (2001), 275–285.
17. L. I. PISCORAN and M. AMINI: *On conformally flat square-root (α, β) -metrics*. J. Finsler Geom. Appl. **2** (2021), 89–102.
18. G. RANDERS: *On an asymmetric metric in the four-space of general relativity*. Phys. Rev. B **59** (1941), 195–199.
19. H. RUND: *The Differential Geometry of Finsler Spaces*. Berlin, Springer-Verlag, 1959.
20. H. SHIMADA and S. V. SABAU: *Introduction to Matsumoto metric*. Nonlinear Anal.: Theo., Math. and Appl. **63** (2005), 165–168.
21. A. TAYEBI and M. AMINI: *Conformally flat 4-th root (α, β) -metrics with relatively isotropic mean Landsberg curvature*. Math. Analysis. Convex. Optimization **1** (2020), 25–34.
22. A. TAYEBI and B. NAJAFI: *On m -th root metrics with special curvature properties*. C. R. Acad. Sci. Paris, Ser. I. **349** (2011), 691–693.
23. A. TAYEBI and M. RAZGORDANI: *On conformally flat fourth root (α, β) -metrics*. Differential Geom. Appl. **62** (2019), 253–266.
24. G. YANG: *On a class of Einstein-reversible Finsler metrics*. Differ. Geom. Appl. **60** (2018), 80–103.