

$(\psi, \gamma, 2)$ -CHEREDNIK-OPDAM LIPSCHITZ FUNCTIONS IN THE  
SPACE  $L^2_{\alpha, \beta}(\mathbb{R})$

Radouan Daher and Salah El Ouadih

**Abstract.** In this paper, using a generalized translation operator, we obtain an analog of Younis Theorem 5.2 in [3] for the Cherednik-Opdam transform for functions satisfying the  $(\psi, \gamma, 2)$ -Cherednik-Opdam Lipschitz condition in the space  $L^2_{\alpha, \beta}(\mathbb{R})$ .

**Keywords:** Cherednik-Opdam operator; Cherednik-Opdam transform; generalized translation.

1. Introduction and Preliminaries

Various investigators such as V.N. Mishra and L.N. Mishra [7], Mishra and al. [5, 6] have determined the degree of approximation of  $2\pi$ -periodic signals (functions) belonging to various classes  $Lip\alpha$ ,  $Lip(\alpha, r)$ ,  $Lip(\xi(t), r)$  and  $W(L_r, \xi(t))$ , ( $r \geq 1$ ), of functions through trigonometric Fourier approximation using different summability matrices with monotone rows. In this direction, Younis Theorem 5.2 [3] characterized the set of functions in  $L^2(\mathbb{R})$  satisfying the Cauchy Lipschitz condition by means of an asymptotic estimate growth of the norm of their Fourier transforms, namely we have

**Theorem 1.1.** [3] Let  $f \in L^2(\mathbb{R})$ . Then the following are equivalents

$$(i) \quad \|f(x+h) - f(x)\| = O\left(\frac{h^\delta}{(\log \frac{1}{h})^\gamma}\right), \quad \text{as } h \rightarrow 0, 0 < \delta < 1, \gamma \geq 0,$$

$$(ii) \quad \int_{|\lambda| \geq r} |\widehat{f}(\lambda)|^2 d\lambda = O\left(\frac{r^{-2\delta}}{(\log r)^{2\gamma}}\right), \quad \text{as } r \rightarrow \infty,$$

where  $\widehat{f}$  stands for the Fourier transform of  $f$ .

In this paper, we prove the generalization of Theorem 1.1 for the Cherednik-Opdam transform for functions satisfying the  $(\psi, \gamma, 2)$ -Cherednik-Opdam Lipschitz condition in the space  $L^2_{\alpha, \beta}(\mathbb{R})$ . For this purpose, we use the generalized translation operator. We point out that similar results have been established in the Jacobi

transform [8].

In this section, we develop some results from harmonic analysis related to the differential-difference operator  $T^{(\alpha,\beta)}$ . Further details can be found in [1] and [2]. In the following we fix parameters  $\alpha, \beta$  subject to the constraints  $\alpha \geq \beta \geq -\frac{1}{2}$  and  $\alpha > \frac{-1}{2}$ .

Let  $\rho = \alpha + \beta + 1$  and  $\lambda \in \mathbb{C}$ . The Opdam hypergeometric functions  $G_\lambda^{(\alpha,\beta)}$  on  $\mathbb{R}$  are eigenfunctions  $T^{(\alpha,\beta)}G_\lambda^{(\alpha,\beta)}(x) = i\lambda G_\lambda^{(\alpha,\beta)}(x)$  of the differential-difference operator

$$T^{(\alpha,\beta)}f(x) = f'(x) + [(2\alpha + 1) \coth x + (2\beta + 1) \tanh x] \frac{f(x) - f(-x)}{2} - \rho f(-x),$$

that are normalized such that  $G_\lambda^{(\alpha,\beta)}(0) = 1$ . In the notation of Cherednik one would write  $T^{(\alpha,\beta)}$  as

$$T(k_1 + k_2)f(x) = f'(x) + \left\{ \frac{2k_1}{1 + e^{-2x}} + \frac{4k_2}{1 - e^{-4x}} \right\} (f(x) - f(-x)) - (k_1 + 2k_2)f(x),$$

with  $\alpha = k_1 + k_2 - \frac{1}{2}$  and  $\beta = k_2 - \frac{1}{2}$ . Here  $k_1$  is the multiplicity of a simply positive root and  $k_2$  the (possibly vanishing) multiplicity of a multiple of this root. By [1] or [2], the eigenfunction  $G_\lambda^{(\alpha,\beta)}$  is given by

$$G_\lambda^{(\alpha,\beta)}(x) = \varphi_\lambda^{\alpha,\beta}(x) - \frac{1}{\rho - i\lambda} \frac{\partial}{\partial x} \varphi_\lambda^{\alpha,\beta}(x) = \varphi_\lambda^{\alpha,\beta}(x) + \frac{\rho}{4(\alpha + 1)} \sinh(2x) \varphi_\lambda^{\alpha+1,\beta+1}(x),$$

where  $\varphi_\lambda^{\alpha,\beta}(x) = {}_2F_1\left(\frac{\rho+i\lambda}{2}; \frac{\rho-i\lambda}{2}; \alpha + 1; -\sinh^2 x\right)$  is the classical Jacobi function.

**Lemma 1.1.** [4] *The following inequalities are valids for Jacobi functions  $\varphi_\lambda^{\alpha,\beta}(x)$*

- (i)  $|\varphi_\lambda^{\alpha,\beta}(x)| \leq 1$ .
- (ii)  $1 - \varphi_\lambda^{\alpha,\beta}(x) \leq x^2(\lambda^2 + \rho^2)$ .
- (iii) *there is a constant  $c > 0$  such that*

$$1 - \varphi_\lambda^{\alpha,\beta}(x) \geq c,$$

for  $\lambda x \geq 1$ .

Denote  $L_{\alpha,\beta}^2(\mathbb{R})$ , the space of measurable functions  $f$  on  $\mathbb{R}$  such that

$$\|f\|_{2,\alpha,\beta} = \left( \int_{\mathbb{R}} |f(x)|^2 A_{\alpha,\beta}(x) dx \right)^{1/2} < +\infty,$$

where

$$A_{\alpha,\beta}(x) = (\sinh |x|)^{2\alpha+1} (\cosh |x|)^{2\beta+1}.$$

The Cherednik-Opdam transform of  $f \in C_c(\mathbb{R})$  is defined by

$$\mathcal{H}f(\lambda) = \int_{\mathbb{R}} f(x) G_\lambda^{(\alpha,\beta)}(-x) A_{\alpha,\beta}(x) dx \quad \text{for all } \lambda \in \mathbb{C}.$$

The inverse transform is given as

$$\mathcal{H}^{-1}g(x) = \int_{\mathbb{R}} g(\lambda)G_{\lambda}^{(\alpha,\beta)}(x) \left(1 - \frac{\rho}{i\lambda}\right) \frac{d\lambda}{8\pi|c_{\alpha,\beta}(\lambda)|^2},$$

here

$$c_{\alpha,\beta}(\lambda) = \frac{2^{\rho-i\lambda}\Gamma(\alpha+1)\Gamma(i\lambda)}{\Gamma(\frac{1}{2}(\rho+i\lambda))\Gamma(\frac{1}{2}(\alpha-\beta+1+i\lambda))}.$$

The corresponding Plancherel formula was established in [1], to the effect that

$$\int_{\mathbb{R}} |f(x)|^2 A_{\alpha,\beta}(x) dx = \int_0^{+\infty} (|\mathcal{H}f(\lambda)|^2 + |\mathcal{H}\check{f}(\lambda)|^2) d\sigma(\lambda),$$

where  $\check{f}(x) := f(-x)$  and  $d\sigma$  is the measure given by

$$d\sigma(\lambda) = \frac{d\lambda}{16\pi|c_{\alpha,\beta}(\lambda)|^2}.$$

According to [2] there exists a family of signed measures  $\mu_{x,y}^{(\alpha,\beta)}$  such that the product formula

$$G_{\lambda}^{(\alpha,\beta)}(x)G_{\lambda}^{(\alpha,\beta)}(y) = \int_{\mathbb{R}} G_{\lambda}^{(\alpha,\beta)}(z)d\mu_{x,y}^{(\alpha,\beta)}(z)$$

holds for all  $x, y \in \mathbb{R}$  and  $\lambda \in \mathbb{C}$ , where

$$d\mu_{x,y}^{(\alpha,\beta)}(z) = \begin{cases} \mathcal{K}_{\alpha,\beta}(x, y, z)A_{\alpha,\beta}(z)dz, & \text{if } xy \neq 0 \\ d\delta_x(z), & \text{if } y = 0 \\ d\delta_y(z), & \text{if } x = 0 \end{cases}$$

and

$$\begin{aligned} \mathcal{K}_{\alpha,\beta}(x, y, z) &= M_{\alpha,\beta} |\sinh x \cdot \sinh y \cdot \sinh z|^{-2\alpha} \int_0^{\pi} g(x, y, z, \chi)_+^{\alpha-\beta-1} \\ &\times [1 - \sigma_{x,y,z}^{\chi} + \sigma_{x,z,y}^{\chi} + \sigma_{z,y,x}^{\chi} + \frac{\rho}{\beta + \frac{1}{2}} \coth x \cdot \coth y \cdot \coth z (\sin \chi)^2] \times (\sin \chi)^{2\beta} d\chi \end{aligned}$$

if  $x, y, z \in \mathbb{R} \setminus \{0\}$  satisfy the triangular inequality  $||x| - |y|| < |z| < |x| + |y|$ , and  $\mathcal{K}_{\alpha,\beta}(x, y, z) = 0$  otherwise. Here

$$\forall x, y, z \in \mathbb{R}, \chi \in [0, 1], \sigma_{x,y,z}^{\chi} = \begin{cases} \frac{\cosh x + \cosh y - \cosh z \cos \chi}{\sinh x \sinh y}, & \text{if } xy \neq 0 \\ 0, & \text{if } xy = 0 \end{cases}$$

and  $g(x, y, z, \chi) = 1 - \cosh^2 x - \cosh^2 y \cdot \cosh^2 z + 2 \cosh x \cdot \cosh y \cdot \cosh z \cdot \cos \chi$ .

**Lemma 1.2.** [2] For all  $x, y \in \mathbb{R}$ , we have

- (i)  $\mathcal{K}_{\alpha,\beta}(x, y, z) = \mathcal{K}_{\alpha,\beta}(y, x, z)$ .
- (ii)  $\mathcal{K}_{\alpha,\beta}(x, y, z) = \mathcal{K}_{\alpha,\beta}(-x, z, y)$ .
- (iii)  $\mathcal{K}_{\alpha,\beta}(x, y, z) = \mathcal{K}_{\alpha,\beta}(-z, y, -x)$ .

The product formula is used to obtain explicit estimates for the generalized translation operators

$$\tau_x^{(\alpha,\beta)} f(y) = \int_{\mathbb{R}} f(z) d\mu_{x,y}^{(\alpha,\beta)}(z).$$

It is known from [2] that

$$(1.1) \quad \mathcal{H}\tau_x^{(\alpha,\beta)} f(\lambda) = G_\lambda^{(\alpha,\beta)}(x)\mathcal{H}f(\lambda),$$

for  $f \in C_c(\mathbb{R})$ .

### 2. Main Result

In this section we give the main result of this paper. We need first to define  $(\psi, \gamma, 2)$ -Cherednik-Opdam Lipschitz class.

Denote  $N_h$  by

$$N_h = \tau_h^{(\alpha,\beta)} + \tau_{-h}^{(\alpha,\beta)} - 2I,$$

where  $I$  is the unit operator in the space  $L^2_{\alpha,\beta}(\mathbb{R})$ .

**Definition 2.1.** Let  $\gamma \geq 0$ . A function  $f \in L^2_{\alpha,\beta}(\mathbb{R})$  is said to be in the  $(\psi, \gamma, 2)$ -Cherednik-Opdam Lipschitz class, denoted by  $Lip(\psi, \gamma, 2)$ , if

$$\|N_h f(x)\|_{2,\alpha,\beta} = O\left(\frac{\psi(h)}{\left(\log \frac{1}{h}\right)^\gamma}\right) \quad \text{as } h \rightarrow 0,$$

where

- (a)  $\psi$  is a continuous increasing function on  $[0, \infty)$ ,
- (b)  $\psi(0) = 0$ ,  $\psi(ts) = \psi(t)\psi(s)$  for all  $t, s \in [0, \infty)$ ,
- (c) and

$$\int_0^{1/h} s\psi(s^{-2})(\log s)^{-2\gamma} ds = O\left(h^{-2}\psi(h^2)\left(\log \frac{1}{h}\right)^{-2\gamma}\right), \quad h \rightarrow 0.$$

**Lemma 2.1.** If  $f \in C_c(\mathbb{R})$ , then

$$(2.1) \quad \mathcal{H}\tilde{\tau}_x^{(\alpha,\beta)} f(\lambda) = G_\lambda^{(\alpha,\beta)}(-x)\mathcal{H}\check{f}(\lambda).$$

*Proof.* For  $f \in C_c(\mathbb{R})$ , we have

$$\begin{aligned} \mathcal{H}\tilde{\tau}_x^{(\alpha,\beta)} f(\lambda) &= \int_{\mathbb{R}} \tau_x^{(\alpha,\beta)} f(-y) G_\lambda^{(\alpha,\beta)}(-y) A_{\alpha,\beta}(y) dy \\ &= \int_{\mathbb{R}} \tau_x^{(\alpha,\beta)} f(y) G_\lambda^{(\alpha,\beta)}(y) A_{\alpha,\beta}(y) dy \\ &= \int_{\mathbb{R}} \left[ \int_{\mathbb{R}} f(z) \mathcal{K}_{\alpha,\beta}(x, y, z) A_{\alpha,\beta}(z) dz \right] G_\lambda^{(\alpha,\beta)}(y) A_{\alpha,\beta}(y) dy \\ &= \int_{\mathbb{R}} f(z) \left[ \int_{\mathbb{R}} G_\lambda^{(\alpha,\beta)}(y) \mathcal{K}_{\alpha,\beta}(x, y, z) A_{\alpha,\beta}(y) dy \right] A_{\alpha,\beta}(z) dz. \end{aligned}$$

Since  $\mathcal{K}_{\alpha,\beta}(x, y, z) = \mathcal{K}_{\alpha,\beta}(-x, z, y)$ , it follows from the product formula that

$$\begin{aligned} \mathcal{H}\tau_x^{(\alpha,\beta)} f(\lambda) &= G_\lambda^{(\alpha,\beta)}(-x) \int_{\mathbb{R}} f(z) G_\lambda^{(\alpha,\beta)}(z) A_{\alpha,\beta}(z) dz \\ &= G_\lambda^{(\alpha,\beta)}(-x) \int_{\mathbb{R}} f(-z) G_\lambda^{(\alpha,\beta)}(-z) A_{\alpha,\beta}(z) dz \\ &= G_\lambda^{(\alpha,\beta)}(-x) \mathcal{H}\check{f}(\lambda). \end{aligned}$$

□

**Lemma 2.2.** For  $f \in L^2_{\alpha,\beta}(\mathbb{R})$ , then

$$\|N_h f(x)\|_{2,\alpha,\beta}^2 = 4 \int_0^{+\infty} |\varphi_\lambda^{\alpha,\beta}(h) - 1|^2 (|\mathcal{H}f(\lambda)|^2 + |\mathcal{H}\check{f}(\lambda)|^2) d\sigma(\lambda).$$

*Proof.* From formulas (1.1) and (2.1), we have

$$\mathcal{H}(N_h f)(\lambda) = (G_\lambda^{(\alpha,\beta)}(h) + G_\lambda^{(\alpha,\beta)}(-h) - 2)\mathcal{H}(f)(\lambda),$$

and

$$\mathcal{H}(\check{N}_h f)(\lambda) = (G_\lambda^{(\alpha,\beta)}(-h) + G_\lambda^{(\alpha,\beta)}(h) - 2)\mathcal{H}(\check{f})(\lambda).$$

Since

$$G_\lambda^{(\alpha,\beta)}(h) = \varphi_\lambda^{\alpha,\beta}(h) + \frac{\rho}{4(\alpha+1)} \sinh(2h)\varphi_\lambda^{\alpha+1,\beta+1}(h),$$

and  $\varphi_\lambda^{\alpha,\beta}$  is even, then

$$\mathcal{H}(N_h f)(\lambda) = 2(\varphi_\lambda^{\alpha,\beta}(h) - 1)\mathcal{H}(f)(\lambda)$$

and

$$\mathcal{H}(\check{N}_h f)(\lambda) = 2(\varphi_\lambda^{\alpha,\beta}(h) - 1)\mathcal{H}(\check{f})(\lambda).$$

Now by Plancherel Theorem, we have the result. □

**Theorem 2.1.** Let  $f \in L^2_{\alpha,\beta}(\mathbb{R})$ . Then the following are equivalent

(a)  $f \in Lip(\psi, \gamma, 2)$ ,

(b)  $\int_r^{+\infty} (|\mathcal{H}f(\lambda)|^2 + |\mathcal{H}\check{f}(\lambda)|^2) d\sigma(\lambda) = O\left(\frac{\psi(r^{-2})}{(\log r)^{2\gamma}}\right)$ , as  $r \rightarrow \infty$ .

*Proof.* (a)  $\Rightarrow$  (b) Let  $f \in Lip(\psi, \gamma, 2)$ . Then we have

$$\|N_h f(x)\|_{2,\alpha,\beta} = O\left(\frac{\psi(h)}{(\log \frac{1}{h})^\gamma}\right) \text{ as } h \rightarrow 0.$$

From Lemma 2.2, we have

$$\|N_h f(x)\|_{2,\alpha,\beta}^2 = 4 \int_0^{+\infty} |1 - \varphi_\lambda^{\alpha,\beta}(h)|^2 (|\mathcal{H}f(\lambda)|^2 + |\mathcal{H}\check{f}(\lambda)|^2) d\sigma\lambda.$$

If  $\lambda \in [\frac{1}{h}, \frac{2}{h}]$ , then  $\lambda h \geq 1$  and (iii) of Lemma 1.1 implies that

$$1 \leq \frac{1}{c^2} |1 - \varphi_\lambda^{\alpha, \beta}(h)|^2.$$

Then

$$\begin{aligned} \int_{\frac{1}{h}}^{\frac{2}{h}} (|\mathcal{H}f(\lambda)|^2 + |\mathcal{H}\check{f}(\lambda)|^2) d\sigma(\lambda) &\leq \frac{1}{c^2} \int_{\frac{1}{h}}^{\frac{2}{h}} |1 - \varphi_\lambda^{\alpha, \beta}(h)|^2 (|\mathcal{H}f(\lambda)|^2 + |\mathcal{H}\check{f}(\lambda)|^2) d\sigma(\lambda) \\ &\leq \frac{1}{c^2} \int_0^{+\infty} |1 - \varphi_\lambda^{\alpha, \beta}(h)|^2 (|\mathcal{H}f(\lambda)|^2 + |\mathcal{H}\check{f}(\lambda)|^2) d\sigma(\lambda) \\ &\leq \frac{1}{4c^2} \|N_h f(x)\|_{2, \alpha, \beta}^2 \\ &= O\left(\frac{\psi(h^2)}{(\log \frac{1}{h})^{2\gamma}}\right). \end{aligned}$$

We obtain

$$\int_r^{2r} (|\mathcal{H}f(\lambda)|^2 + |\mathcal{H}\check{f}(\lambda)|^2) d\sigma(\lambda) \leq C \frac{\psi(r^{-2})}{(\log r)^{2\gamma}}, \quad r \rightarrow \infty,$$

where  $C$  is a positive constant. Now,

$$\begin{aligned} \int_r^{+\infty} (|\mathcal{H}f(\lambda)|^2 + |\mathcal{H}\check{f}(\lambda)|^2) d\sigma(\lambda) &= \sum_{i=0}^{\infty} \int_{2^i r}^{2^{i+1} r} (|\mathcal{H}f(\lambda)|^2 + |\mathcal{H}\check{f}(\lambda)|^2) d\sigma(\lambda) \\ &\leq C \left( \frac{\psi(r^{-2})}{(\log r)^{2\gamma}} + \frac{\psi((2r)^{-2})}{(\log 2r)^{2\gamma}} + \frac{\psi((4r)^{-2})}{(\log 4r)^{2\gamma}} + \dots \right) \\ &\leq C \frac{\psi(r^{-2})}{(\log r)^{2\gamma}} (1 + \psi(2^{-2}) + (\psi(2^{-2}))^2 + (\psi(2^{-2}))^3 + \dots) \\ &\leq K_\psi \frac{\psi(r^{-2})}{(\log r)^{2\gamma}}, \end{aligned}$$

where  $K_\psi = C(1 - \psi(2^{-2}))^{-1}$  since  $\psi(2^{-2}) < 1$ .

Consequently

$$\int_r^{+\infty} (|\mathcal{H}f(\lambda)|^2 + |\mathcal{H}\check{f}(\lambda)|^2) d\sigma(\lambda) = O\left(\frac{\psi(r^{-2})}{(\log r)^{2\gamma}}\right), \quad \text{as } r \rightarrow \infty.$$

(b)  $\Rightarrow$  (a). Suppose now that

$$\int_r^{+\infty} (|\mathcal{H}f(\lambda)|^2 + |\mathcal{H}\check{f}(\lambda)|^2) d\sigma(\lambda) = O\left(\frac{\psi(r^{-2})}{(\log r)^{2\gamma}}\right), \quad \text{as } r \rightarrow \infty,$$

and write

$$\|N_h f(x)\|_{2, \alpha, \beta}^2 = 4(I_1 + I_2),$$

where

$$I_1 = \int_0^{\frac{1}{h}} |1 - \varphi_\lambda^{\alpha, \beta}(h)|^2 (|\mathcal{H}f(\lambda)|^2 + |\mathcal{H}\check{f}(\lambda)|^2) d\sigma\lambda,$$

and

$$I_2 = \int_{\frac{1}{h}}^{+\infty} |1 - \varphi_\lambda^{\alpha, \beta}(h)|^2 (|\mathcal{H}f(\lambda)|^2 + |\mathcal{H}\check{f}(\lambda)|^2) d\sigma\lambda.$$

Firstly, we use the formula  $|\varphi_\lambda^{\alpha, \beta}(h)| \leq 1$  and

$$I_2 \leq 4 \int_{\frac{1}{h}}^{+\infty} (|\mathcal{H}f(\lambda)|^2 + |\mathcal{H}\check{f}(\lambda)|^2) d\sigma(\lambda) = O\left(\frac{\psi(h^2)}{(\log \frac{1}{h})^{2\gamma}}\right), \quad \text{as } h \rightarrow 0.$$

To estimate  $I_1$ , we use the inequalities (i) and (ii) of Lemma 1.1

$$\begin{aligned} I_1 &= \int_0^{\frac{1}{h}} |1 - \varphi_\lambda^{\alpha, \beta}(h)|^2 (|\mathcal{H}f(\lambda)|^2 + |\mathcal{H}\check{f}(\lambda)|^2) d\sigma\lambda \\ &\leq 2 \int_0^{\frac{1}{h}} |1 - \varphi_\lambda^{\alpha, \beta}(h)| (|\mathcal{H}f(\lambda)|^2 + |\mathcal{H}\check{f}(\lambda)|^2) d\sigma\lambda \\ &\leq 2h^2 \int_0^{\frac{1}{h}} (\lambda^2 + \rho^2) (|\mathcal{H}f(\lambda)|^2 + |\mathcal{H}\check{f}(\lambda)|^2) d\sigma\lambda. \end{aligned}$$

Now, we apply integration by parts for a function

$$\phi(s) = \int_s^{+\infty} (|\mathcal{H}f(\lambda)|^2 + |\mathcal{H}\check{f}(\lambda)|^2) d\sigma(\lambda)$$

to get

$$\begin{aligned} I_1 &\leq -2h^2 \int_0^{1/h} (s^2 + \rho^2) \phi'(s) ds \\ &\leq -2h^2 \int_0^{1/h} s^2 \phi'(s) ds \\ &\leq h^2 \left( -\frac{1}{h^2} \phi\left(\frac{1}{h}\right) + 2 \int_0^{1/h} s \phi(s) ds \right) \\ &\leq -\phi\left(\frac{1}{h}\right) + 2h^2 \int_0^{1/h} s \phi(s) ds \\ &\leq 2h^2 \int_0^{1/h} s \phi(s) ds. \end{aligned}$$

Since  $\phi(s) = O\left(\frac{\psi(s^{-2})}{(\log s)^{2\gamma}}\right)$ , we have  $s\phi(s) = O\left(\frac{s\psi(s^{-2})}{(\log s)^{2\gamma}}\right)$  and

$$\int_0^{1/h} s\phi(s) ds = O\left(\int_0^{1/h} \frac{s\psi(s^{-2})}{(\log s)^{2\gamma}} ds\right) = O\left(\frac{h^{-2}\psi(h^2)}{(\log \frac{1}{h})^{2\gamma}}\right),$$

so that

$$I_1 = O\left(\frac{\psi(h^2)}{(\log \frac{1}{h})^{2\gamma}}\right).$$

Consequently,

$$\|N_h f(x)\|_{2,\alpha,\beta} = O\left(\frac{\psi(h)}{(\log \frac{1}{h})^\gamma}\right) \quad \text{as } h \rightarrow 0,$$

and this ends the proof of the theorem.  $\square$

### 3. Conclusion

In this work we have succeeded to generalise the theorem in [3] for the Cherednik-Opdam transform in the space  $L^2_{\alpha,\beta}(\mathbb{R})$ . We proved that  $f(x)$  belong to  $Lip(\psi, \gamma, 2)$ . Then

$$\int_r^{+\infty} (|\mathcal{H}f(\lambda)|^2 + |\mathcal{H}\check{f}(\lambda)|^2) d\sigma(\lambda) = O\left(\frac{\psi(r^{-2})}{(\log r)^{2\gamma}}\right), \quad \text{as } r \rightarrow \infty.$$

### Acknowledgements

The authors would like to thank the referee for his valuable comments and suggestions.

### REFERENCES

1. E. M. OPDAM, *Harmonic analysis for certain representations of graded Hecke algebras*, Acta Math. Vol. 175, no. 1, (1995), 75-121.
2. J. P. ANKER, F. AYADI and M. SIFI, *Opdams hypergeometric functions: product formula and convolution structure in dimension 1*, Adv. Pure Appl. Math. Vol. 3, no. 1, (2012), 11-44.
3. M. S. YOUNIS, *Fourier transforms of Dini-Lipschitz functions*. Int. J. Math. Math. Sci. Vol. 9, no. 2,(1986), 301-312. doi:10.1155/S0161171286000376.
4. S. S. PLATONOV, *Approximation of functions in  $L_2$ -metric on noncompact rank 1 symmetric space*. Algebra Analiz . Vol. 11, no. 1, (1999), 244-270.
5. L. N. MISHRA, V. N. MISHRA, K. KHATRI and DEEPMALA, *On the trigonometric approximation of signals belonging to generalized weighted Lipschitz  $W(L^r, \xi(t))$ , ( $r \geq 1$ ) class by matrix  $(C^1.N_p)$  Operator of conjugate series of its Fourier series*, Applied Mathematics and Computation, Vol. 237, (2014), 252-263.
6. V. N. MISHRA, K. KHATRI, L. N. MISHRA and DEEPMALA; *Trigonometric approximation of periodic signals belonging to generalized weighted Lipschitz  $W'(L_r, \xi(t))$ , ( $r \geq 1$ )– class by Nörlund-Euler  $(N, p_n)(E, q)$  operator of conjugate series of its Fourier series*, Journal of Classical Analysis, Vol. 5, no. 2 (2014), 91-105. doi:10.7153/jca-05-08.



7. V. N. MISHRA and L. N. MISHRA, *Trigonometric approximation of signals (functions) in  $L_p$  ( $p \geq 1$ )-norm*, International Journal of Contemporary Mathematical Sciences, Vol. 7, no. 19, (2012), 909-918.
8. A. ABOUELAZ, R. DAHER and M. EL HAMMA, *Generalization of Titchmarsh's theorem for the Jacobi transform*, Ser. Math. Inform. Vol. 28, no. 1, (2013), 43-51.

Radouan Daher  
Department of Mathematics  
Faculty of Sciences Ain Chock  
University Hassan II, Casablanca, Morocco  
rjdaher024@gmail.com

Salah El Ouadih  
Department of Mathematics  
Faculty of Sciences Ain Chock  
University Hassan II, Casablanca, Morocco  
salahwadih@gmail.com