

ON GENERALIZED FIBONACCI DIFFERENCE SPACE DERIVED FROM THE ABSOLUTELY p - SUMMABLE SEQUENCE SPACES

Gülşen Kılınc

© 2019 by University of Niš, Serbia | Creative Commons Licence: CC BY-NC-ND

Abstract. In this study, it is specified *the sequence space* $l(F(r, s), p)$, (where $p = (p_k)$ is any bounded sequence of positive real numbers) and researched some algebraic and topological features of this space. Further, α -, β -, γ - duals and its Schauder Basis are given. The classes of *matrix transformations* from the space $l(F(r, s), p)$ to the spaces l_∞, c , and c_0 are qualified. Additionally, acquiring qualifications of some other *matrix transformations* from the space $l(F(r, s), p)$ to the *Euler, Riesz, difference, etc., sequence spaces* is the other result of the paper.

Keywords. Matrix transformations; sequence space; Schauder Basis.

1. Introduction

In the first instance, let's remember some basic concept definitions in summability theory. The symbol w denotes the space of all real or complex valued sequences. A subspace of w is entitled *sequence space*. Some of the most known sequence spaces are l_∞, c, c_0 and l_p ($1 \leq p \leq \infty$) known as classical *sequence spaces*. The spaces which represented by these symbols are all bounded, convergent, null sequences and absolutely p -summable sequences, respectively. The spaces l_∞, c, c_0 are Banach spaces with the norm

$$(1.1) \quad \|z\|_\infty = \sup_{r \in \mathbb{N}} |z_r|,$$

and the space l_p ($1 \leq p \leq \infty$) is Banach space with the norm

$$(1.2) \quad \|z\|_p = \left(\sum_r |z_r|^p \right)^{\frac{1}{p}}.$$

The spaces bs and cs are the space of the sequences that constitute the bounded and convergent series, respectively. The space c_0s is defined

$$(1.3) \quad c_0s = \left\{ y = (y_r) \in w : \sum_r y_r = 0 \right\},$$

the sequence space bv defined by

$$(1.4) \quad \left\{ z = (z_r) \in w : \sum_{r=1}^{\infty} |z_r - z_{r-1}| < \infty \right\}.$$

The space bv_0 denotes $bv_0 = bv \cap c_0$.

To straightforwardness in representation, the symbol \sum_r will be used instead of the symbol $\sum_{r=0}^{\infty}$ throughout this study. Also, the representations e and $e^{(n)}$ denote $(1, 1, \dots)$ and the sequence n -th unit vector, respectively; where $n \in \mathbb{N}$ and $\mathbb{N} = \{0, 1, 2, \dots\}$.

A sequence space ϑ with a linear topology is entitled a K -space ensured that each of the maps $p_i : \vartheta \rightarrow \mathbb{C}$ defined by $p_i(z) = z_i$ is continuous for all $i \in \mathbb{N}$, where \mathbb{C} demonstrates the complex field. A K -space ϑ is entitled an FK -space ensured ϑ is a complete linear metric space. An FK -space whose topology is normable is entitled a BK -space (see [1]) which comprises Φ , the set of all finitely nonzero sequences. [12]

If algebraic operations are continuous in a linear T space fitted with a ρ metric, then (T, ρ) is entitled linear metric space. (see [27]). That is, when (t_n) and (z_n) are any two sequences in T and (k_n) is a sequence of scalars, then provision of conditions $\lim_{n \rightarrow \infty} \rho(t_n, t) = 0$, $\lim_{n \rightarrow \infty} \rho(z_n, z) = 0$ and $\lim_{n \rightarrow \infty} \rho(k_n) = k$ means provision of results $\lim_{n \rightarrow \infty} \rho(t_n + z_n, t + z) = 0$ and $\lim_{n \rightarrow \infty} \rho(k_n t_n, kt) = 0$. If T linear metric space is complete, it is named Frechet sequence space [28].

Let ν and η be a sequence space and $A = (a_{ij})$ an infinite matrix of real or complex numbers and where $i, j \in \mathbb{N}$. If for each sequence $z = (z_j)$ in ν , the A -transform of z is in η , then A - is called a matrix transformation from ν into η and we demonstrate it by writing $A : \nu \rightarrow \eta$, where for all $i \in \mathbb{N}$,

$$(1.5) \quad (Az)_i = \sum_j a_{ij} z_j$$

The class of all matrices specified $A : \nu \rightarrow \eta$ is denoted by the notation $(\nu : \eta)$. The infinite matrix A belongs to the class $(\nu : \eta)$ iff for each $i \in \mathbb{N}$, and every $z \in \nu$, $\sum_j a_{ij} z_j$ series are convergent and $(Az)_i \in \eta$.

The matrix domain ν_A of an infinite matrix A in a sequence space ν is specified by

$$(1.6) \quad \nu_A = \{z = (z_k) \in w : Az \in \nu\}.$$

Let U is an infinite matrix. If the constituents on the principal diagonal of U are non- zero and the constituents on the top of the principal diagonal are zero, U

is entitled the triangular matrix. To study with triangular matrix domains has a special significance because of the features that these matrices have. Here are some of them: It is trivial that $Z(Ty) = (ZT)y$ holds for the triangle matrices Z, T , and a sequence y . Further, a triangular matrix U has an inverse matrix $U^{-1} = V$ that is only one and triangular. Then, $y = U(Vy) = V(Uy)$ holds for all $y \in w$. If Z is triangle and ν is a BK -space, then ν_Z is also a BK -space with the norm given by $\|y\|_{\nu_Z} = \|Zy\|_{\nu}$ for all $y \in \nu_Z$.

If a function $h : T \rightarrow \mathbb{R}$ fulfills the the undermentioned conditions, for all $y, z \in T$

i) $h(y) = 0$ if $y = \theta$,

ii) $h(y) = h(-y)$,

iii) $h(y + z) \leq h(y) + h(z)$

iv) $|\beta_n - \beta| \rightarrow 0$ and $h(y_n - y) \rightarrow 0$ imply $h(\beta_n y_n - \beta y) \rightarrow 0$, for all β 's in \mathbb{R} and all y 's in T , where θ is the zero vector in the linear space T .

Then, a linear topological space T defined on the real field \mathbb{R} is entitled a paranormed space.

Let (p_r) be a bounded sequence of exactly positive real numbers with $\sup_r p_r = H$ and $M = \max\{1, H\}$. Then, absolutely p -summable sequences space $l(p)$ was specified by Maddox [13] (see also [14] and [15]) such as

$$(1.7) \quad l(p) = \left\{ y = (y_r) \in w : \sum_r |y_r|^{p_r} < \infty \right\},$$

where $0 \leq p_r \leq H < \infty$. It is the complete space paranormed by

$$(1.8) \quad h(y) = \left(\sum_r |y_r|^{p_r} \right)^{1/M}.$$

Throughout this study, \mathcal{F} symbolizes the collection of all finite subsets of \mathbb{N} and any constituent with a negative index is considered to be zero.

Let us bring to mind the sequence (f_n) of Fibonacci numbers given by the linear iteration correlates: The first two constituents are taken as 1 and other index constituent are found by summing up the last two constituent that preceded it. Fibonacci numbers have many quirky features and practices in sciences, arts, and architecture. For instance, the ratio sequences of Fibonacci numbers converges to the golden ratio which is significant in sciences and arts. Also, some fundamental features of Fibonacci numbers (f_n) are given as below:

$$(1.9) \quad \lim_{n \rightarrow \infty} \frac{f_{n+1}}{f_n} = \frac{1 + \sqrt{5}}{2} = \phi \quad (\text{Golden Ratio}),$$

$$(1.10) \quad \sum_{k=0}^n f_k = f_{n+2} - 1, \quad \text{for each } n \in \mathbb{N},$$

$$(1.11) \quad \sum_k \frac{1}{f_k} < \infty,$$

$$(1.12) \quad f_{n-1} \cdot f_{n+1} - f_n^2 = (-1)^{n+1} \quad \text{for each } n \geq 1 \quad (\text{Cassini Formula}).$$

The practice forming a new sequence space through the medium of the matrix domain of a special triangle has latterly been utilized by several authors in many exploratory papers. One of them is Fibonacci matrix defined Kara [4]. Now, let us familiarize aforementioned Fibonacci matrix $F = (f_{nk})$

$$(1.13) \quad f_{nk} = \begin{cases} -\frac{f_{n+1}}{f_n} & , \quad (k = n - 1), \\ \frac{f_n}{f_{n+1}} & , \quad (k = n), \\ 0 & , \quad (0 \leq k < n - 1 \text{ or } k > n). \end{cases}$$

It can be demonstrated readily that the matrix F is conservative, but it is neither regular nor coercive.

Let us acquaint some *sequence spaces* defined by the domain of a matrix and will be named in this work:

$$(1.14) \quad bv_\infty = \{y = (y_r) \in w : (y_r - y_{r-1}) \in l_\infty\},$$

was introduced by Başar and Altay [1].

$$(1.15) \quad e_\infty^r = \{y = (y_k) \in w : E^r y \in l_\infty\},$$

was introduced by Altay, Başar and Mursaleen [6].

$$(1.16) \quad e_c^r = \{y = (y_k) \in w : E^r y \in c\},$$

$$(1.17) \quad e_0^r = \{y = (y_k) \in w : E^r y \in c_0\},$$

were introduced by Altay, Başar [8]

$$(1.18) \quad X_\infty = \{y = (y_r) \in w : C_1 y \in l_\infty\},$$

was introduced by Ng and Lee [2]

$$(1.19) \quad \tilde{c} = \{y = (y_r) \in w : C_1 y \in c\},$$

$$(1.20) \quad \tilde{c}_0 = \{y = (y_r) \in w : C_1 y \in c_0\},$$

were introduced by Şengönül and Başar [9]

$$(1.21) \quad r_\infty^t = \{y = (y_k) \in w : R^t y \in l_\infty\},$$

was introduced by Altay and Başar [7].

$$(1.22) \quad r_c^t = \{y = (y_k) \in w : R^t y \in c\},$$

$$(1.23) \quad r_0^t = \{y = (y_k) \in w : R^t y \in c_0\},$$

was introduced by Altay and Başar [10].

$$(1.24) \quad l_\infty(\widehat{F}) = \{y = (y_r) \in w : Fy \in l_\infty\},$$

was introduced by Kara [4].

$$(1.25) \quad c(\widehat{F}) = \{y = (y_r) \in w : Fy \in c\},$$

$$(1.26) \quad c_0(\widehat{F}) = \{y = (y_r) \in w : Fy \in c\},$$

were introduced by Başarır et all [11].

$$(1.27) \quad c(\Delta) = \{y = (y_r) \in w : (y_r - y_{r+1}) \in c\},$$

$$(1.28) \quad c_0(\Delta) = \{y = (y_r) \in w : (y_r - y_{r+1}) \in c_0\},$$

was introduced by Kızılmaz [5] where E^r, C_1, R^t, F denote *Euler mean* of order r , *arithmetic, Riesz means and Fibonacci matrix*, respectively. Now, let us give aforementioned *matrix methods*:

$$(1.29) \quad e_{nk}^r = \begin{cases} \binom{n}{k} (1-r)^{n-k} r^k, & 0 \leq k \leq n, \\ 0, & k > n, \end{cases}$$

$$(1.30) \quad \delta_{nk} = \begin{cases} (-1)^{n-k}, & n-1 \leq k \leq n, \\ 0, & 0 \leq k < n-1 \text{ or } k > n, \end{cases}$$

the matrix $R^t = (r_{nk}^t)$ is specified by

$$(1.31) \quad r_{nk}^t = \begin{cases} \frac{t_k}{T_n}, & 0 \leq k \leq n, \\ 0, & k > n, \end{cases}$$

Ng and Lee specified the *Cesaro sequence spaces* X_p and X_∞ of non-absolute type by the set of the sequences whose C -transforms are in l_p and l_∞ , respectively.

The matrix $C = (c_{nk})$ is specified as follow:

$$(1.32) \quad c_{nk} = \begin{cases} \frac{1}{n+1}, & 0 \leq k \leq n, \\ 0, & k > n, \end{cases}$$

Now, let us present the matrix method which will be used to specify the sequence space which is subject of this work. It is entitled the *generalized Fibonacci band matrix* $F(r, s)$ specified by Candan at first [3]. It is formed by using *Fibonacci sequence* and non-zero real numbers r and s .

$$(1.33) \quad f_{nk}(r, s) = \begin{cases} s \frac{f_{n+1}}{f_n}, & k = n - 1, \\ r \frac{f_n}{f_{n+1}}, & k = n, \\ 0, & \text{in other cases.} \end{cases}$$

It can be seen that, the matrix $F(r, s)$ is degraded to the matrix F , for $r = 1$ and $s = -1$. Therefore, the data acquired from the matrix $F(r, s)$ is more general than the data acquired from the \hat{F} matrix. The inverse $F^{-1}(r, s)$ of the matrix $F(r, s)$ is calculated as

$$(1.34) \quad f_{nk}^{-1}(r, s) = \begin{cases} \frac{1}{r} \left(-\frac{s}{r}\right)^{n-k} \frac{f_{n+1}^2}{f_k \cdot f_{k+1}}, & 0 \leq k \leq n, \\ 0, & k > n. \end{cases}$$

The other significant principal paper are here: [3], [16], [6], [17], [18], [26], [19], [20], [21], [22], [23], [24], [25], [31], [32], [34].

This paper is organized as follow: In first chapter, it is familiarized generalized Fibonacci difference space $l(F(r, s), p)$ and is given its some algebraic and topological features. In second chapter, its α -, β -, γ - duals and Schauder Basis are determined. In third chapter, the qualifications of some other *matrix transformations* from the space $l(F(r, s), p)$ to the *Euler, Riesz, difference, etc., sequence spaces* are acquired.

2. Some properties of the sequence space $l(F(r, s), p)$

In this chapter, firstly, we familiarize *the Fibonacci difference sequence spaces* $l(F(r, s), p)$, $l_p(F(r, s))$ as the set of all sequences whose $F(r, s)$ -transforms are in the spaces $\ell(p)$ and l_p , respectively. Later, some algebraic and topological particularity of aforementioned these new spaces will be proved and a *Schauder basis* constructed. Let $p = (p_k)$ be any bounded sequence of positive real numbers. Then,

$$(2.1) \quad l(F(r, s), p) = \left\{ y = (y_n) \in w : \sum_n \left| r \frac{f_n}{f_{n+1}} y_n + s \frac{f_{n+1}}{f_n} y_{n-1} \right|^{p_k} < \infty \right\},$$

where $0 < p_k \leq H < \infty$. In the case, $p_k = p$ for all $k \in \mathbb{N}$, the space $l(F(r, s), p)$ is degraded to space $l_p(F(r, s))$, i.e., for $(p \geq 1)$

$$(2.2) \quad l_p(F(r, s)) = \left\{ y = (y_n) \in w : \sum_n \left| r \frac{f_n}{f_{n+1}} y_n + s \frac{f_{n+1}}{f_n} y_{n-1} \right|^p < \infty \right\},$$

By means of the notation of (1.6), the spaces $l(F(r, s), p)$ and $l_p(F(r, s))$ can be redescribed as follows:

$$(2.3) \quad l(F(r, s), p) = (l(p))_{F(r,s)} \text{ and } l_p(F(r, s)) = (l_p)_{F(r,s)}.$$

Define the sequence $y = (y_k)$ by the $F(r, s)$ -transform of a sequence $x = (x_k)$, i.e.,

for $n = 0, y_0 = rx_0$ and for $n \geq 1$

$$(2.4) \quad y_n = (F(r, s)x)_n = r \frac{f_n}{f_{n+1}} x_n + s \frac{f_{n+1}}{f_n} x_{n-1},$$

for all $k \in \mathbb{N}$. So, It can be acquired by a simplistic calculation that

$$(2.5) \quad x_n = \sum_{j=0}^n \frac{1}{r} \left(-\frac{s}{r}\right)^{n-j} \frac{f_{n+1}^2}{f_j \cdot f_{j+1}} y_j$$

Throughout the text, we assume that the sequences $x = (x_k)$ and $y = (y_k)$ are attached to the connection (2.4)

The following inequality will be employed throughout the paper. Let $p = (p_k)$ be a sequence of positive real numbers with $0 < p_k \leq \sup_k p_k = H$, and let $D = \max\{1, 2^{H-1}\}$. Then, for the factorable sequences (c_k) and (d_k) in the complex plane, we have

$$(2.6) \quad |c_k + d_k|^{p_k} \leq D \left(|c_k|^{p_k} + |d_k|^{p_k} \right)$$

The other inequality is *Minkowsky inequality*, which will be employed in this paper. Its expression is here:

$$(2.7) \quad \left(\sum_{k=1}^{\infty} |c_k + d_k|^p \right)^{1/p} \leq \left(\sum_{k=1}^{\infty} |c_k|^p \right)^{1/p} + \left(\sum_{k=1}^{\infty} |d_k|^p \right)^{1/p}$$

Theorem 2.1. $l(F(r, s), p)$ is a linear, complete and paranormed space with the h function specified as

$$(2.8) \quad h(x) = \left(\sum_n \left| r \frac{f_n}{f_{n+1}} x_n + s \frac{f_{n+1}}{f_n} x_{n-1} \right|^{p_n} \right)^{\frac{1}{M}},$$

where $0 < p_n \leq H < \infty$, for all $n \in \mathbb{N}$.

Proof. Let $x, y \in l(F(r, s), p)$. Then

$$(2.9) \quad \sum_n \left| r \frac{f_n}{f_{n+1}} x_n + s \frac{f_{n+1}}{f_n} x_{n-1} \right|^{p_n} < \infty,$$

$$(2.10) \quad \sum_n \left| r \frac{f_n}{f_{n+1}} y_n + s \frac{f_{n+1}}{f_n} y_{n-1} \right|^{p_n} < \infty.$$

For $\lambda, \mu \in \mathbb{C}$, there comes into being integers M_λ , and N_μ such that $|\lambda| \leq M_\lambda$ and $|\mu| \leq N_\mu$. Employing Inequality (2.1), we have

$$\begin{aligned} & \sum_n \left| \lambda \left(r \frac{f_n}{f_{n+1}} x_n + s \frac{f_{n+1}}{f_n} x_{n-1} \right) + \mu \left(r \frac{f_n}{f_{n+1}} y_n + s \frac{f_{n+1}}{f_n} y_{n-1} \right) \right|^{p_n} \\ & \leq \sum_n \left(|\lambda| \left| r \frac{f_n}{f_{n+1}} x_n + s \frac{f_{n+1}}{f_n} x_{n-1} \right| \right)^{p_n} \\ & \quad + \sum_n \left(|\mu| \left| r \frac{f_n}{f_{n+1}} y_n + s \frac{f_{n+1}}{f_n} y_{n-1} \right| \right)^{p_n} \\ & \leq D.M_\lambda^H \sum_n \left(\left| r \frac{f_n}{f_{n+1}} x_n + s \frac{f_{n+1}}{f_n} x_{n-1} \right| \right)^{p_n} \\ & \quad + D.N_\mu^H \sum_n \left(\left| r \frac{f_n}{f_{n+1}} y_n + s \frac{f_{n+1}}{f_n} y_{n-1} \right| \right)^{p_n} \\ & < \infty. \end{aligned}$$

So that $\lambda x + \mu y \in l(F(r, s), p)$. This substantiates that $l(F(r, s), p)$ is a linear space.

Clearly, $h(x) = h(-x)$, for all $x \in l(F(r, s), p)$. It is unconcealed that $r \frac{f_n}{f_{n+1}} x_n + s \frac{f_{n+1}}{f_n} x_{n-1} = 0$, for $x = 0$. Since $\frac{p_n}{M} \leq 1$, employing *Minkowski Inequality*, we have

$$\begin{aligned} h(x+y) &= \left[\sum_n \left| r \frac{f_n}{f_{n+1}} (x_n + y_n) + s \frac{f_{n+1}}{f_n} (x_{n-1} + y_{n-1}) \right|^{p_n} \right]^{\frac{1}{M}} \\ &= \left[\sum_n \left(\left| \left(r \frac{f_n}{f_{n+1}} x_n + s \frac{f_{n+1}}{f_n} x_{n-1} \right) + \left(r \frac{f_n}{f_{n+1}} y_n + s \frac{f_{n+1}}{f_n} y_{n-1} \right) \right|^{\frac{p_n}{M}} \right)^M \right]^{\frac{1}{M}} \\ &\leq \left[\sum_n \left(\left| r \frac{f_n}{f_{n+1}} x_n + s \frac{f_{n+1}}{f_n} x_{n-1} \right|^{\frac{p_n}{M}} + \left| r \frac{f_n}{f_{n+1}} y_n + s \frac{f_{n+1}}{f_n} y_{n-1} \right|^{\frac{p_n}{M}} \right)^M \right]^{\frac{1}{M}} \\ &\leq \left(\sum_n \left| r \frac{f_n}{f_{n+1}} x_n + s \frac{f_{n+1}}{f_n} x_{n-1} \right|^{p_n} \right)^{\frac{1}{M}} + \left(\sum_n \left| r \frac{f_n}{f_{n+1}} y_n + s \frac{f_{n+1}}{f_n} y_{n-1} \right|^{p_n} \right)^{\frac{1}{M}} \\ &= h(x) + h(y). \end{aligned}$$

Also, since the inequality $|\alpha|^{p_n} \leq \max \{1, |\alpha|^M\}$ is ensured for $\alpha \in \mathbb{R}$, we get

$$\begin{aligned} h(\alpha x) &= \left[\sum_n \left| r \frac{f_n}{f_{n+1}} (\alpha x_n) + s \frac{f_{n+1}}{f_n} (\alpha x_{n-1}) \right|^{p_n} \right]^{\frac{1}{M}}, \\ &= \left(\sum_n |\alpha|^{p_k} \cdot \left| r \frac{f_n}{f_{n+1}} x_n + s \frac{f_{n+1}}{f_n} x_{n-1} \right|^{p_n} \right)^{\frac{1}{M}}, \\ &\leq \max \{1, |\alpha|\} \cdot h(x). \end{aligned}$$

Let (α_n) be a sequence of scalars with $\alpha_n \rightarrow \alpha$, as $n \rightarrow \infty$ and $\{x^{(n)}\}_{n=0}^\infty$ be a sequence of components $x^{(n)} \in l(F(r, s), p)$ with $h(x^{(n)} - x) \rightarrow 0$, as $n \rightarrow \infty$. Then, we follow up that

$$\begin{aligned} (2.11) \quad 0 \leq h(\alpha_n x^{(n)} - \alpha x) &= h(\alpha_n x^{(n)} - \alpha x^{(n)} + \alpha x^{(n)} - \alpha x) \\ &= h((\alpha_n - \alpha) x^{(n)} + \alpha (x^{(n)} - x)), \\ &\leq h((\alpha_n - \alpha) x^{(n)}) + h(\alpha (x^{(n)} - x)), \\ &= |\alpha_n - \alpha| \cdot h(x^{(n)}) + \max \{1, |\alpha|\} \cdot h(x^{(n)} - x). \end{aligned}$$

If we combine the facts $\alpha_n - \alpha \rightarrow 0$, as $n \rightarrow \infty$ and $h(x^{(n)} - x) \rightarrow 0$, as $n \rightarrow \infty$ with (2.11) we acquire that $h(\alpha_n x^{(n)} - \alpha x) \rightarrow 0$, as $n \rightarrow \infty$. That is, scalar multiplication is continuous. This shows that h is a paranorm on $l(F(r, s), p)$. Moreover, if we assume $h(x) = 0$, then we get

$$(2.12) \quad \left| r \frac{f_n}{f_{n+1}} x_n + s \frac{f_{n+1}}{f_n} x_{n-1} \right| = 0,$$

for each $n \in \mathbb{N}$, we acquire that $x = \theta = (0, 0, 0, \dots)$. It demonstrates that h is total paranorm. Now, we indicate that $l(F(r, s), p)$ is complete. Let $(x^{(n)})$ be any Cauchy sequence in $l(F(r, s), p)$, where $x^{(n)} = \{x_1^{(n)}, x_2^{(n)}, x_3^{(n)}, \dots\}$. Then, for a given $\varepsilon > 0$, there comes into being a positive integer $n_0(\varepsilon)$ such that $[h(x^n - x^m)]^M < \varepsilon^M$ for all $n, m > n_0(\varepsilon)$. Since for each immutable $k \in \mathbb{N}$

$$\begin{aligned} &|(F(r, s) x^n)_k - (F(r, s) x^m)_k|^{p_k} \leq \sum_k |(F(r, s) x^n)_k - (F(r, s) x^m)_k|^{p_k}, \\ &= \sum_k \left| \left(r \frac{f_k}{f_{k+1}} x_k^{(n)} + s \frac{f_{k+1}}{f_k} x_{k-1}^{(n)} \right) - \left(r \frac{f_k}{f_{k+1}} x_k^{(m)} + s \frac{f_{k+1}}{f_k} x_{k-1}^{(m)} \right) \right|^{p_k}, \\ &= \sum_k \left| \left(r \frac{f_k}{f_{k+1}} (x_k^{(n)} - x_k^{(m)}) + s \frac{f_{k+1}}{f_k} (x_{k-1}^{(n)} - x_{k-1}^{(m)}) \right) \right|^{p_k}, \\ &= [h(x^n - x^m)]^M < \varepsilon^M. \end{aligned}$$

For every $n, m > n_0(\varepsilon)$, $\{(F(r, s)x^0)_k, (F(r, s)x^1)_k, (F(r, s)x^2)_k, \dots\}$ is a *Cauchy sequence* of real numbers for every immutable $k \in \mathbb{N}$. Since \mathbb{R} is complete, it is convergent, therefore we say

$$(F(r, s)x^n)_k \rightarrow (F(r, s)x)_k$$

as $n \rightarrow \infty$. Employing these infinitely many limits $(F(r, s)x)_0, (F(r, s)x)_1, (F(r, s)x)_2, \dots$ we specify the sequence $\{(F(r, s)x)_0, (F(r, s)x)_1, (F(r, s)x)_2, \dots\}$. For each $k \in \mathbb{N}$ and $n > n_0(\varepsilon)$

$$\begin{aligned} [h(x^n - x)]^M &= \sum_k \left| r \frac{f_k}{f_{k+1}} (x_k^{(n)} - x_k) + s \frac{f_{k+1}}{f_k} (x_{k-1}^{(n)} - x_{k-1}) \right|^{p_k}, \\ &= \sum_k \left| \left(r \frac{f_k}{f_{k+1}} x_k^{(n)} + s \frac{f_{k+1}}{f_k} x_{k-1}^{(n)} \right) - \left(r \frac{f_k}{f_{k+1}} x_k + s \frac{f_{k+1}}{f_k} x_{k-1} \right) \right|^{p_k}, \\ &= \sum_k |(F(r, s)x^n)_k - (F(r, s)x)_k|^{p_k} < \varepsilon^M. \end{aligned}$$

This shows that $(x^n - x) \in l(F(r, s), p)$. Since $l(F(r, s), p)$ is a linear space, we conclude that $x \in l(F(r, s), p)$. It follows that $x^n \rightarrow x$, as $n \rightarrow \infty$, in $l(F(r, s), p)$, it means that $l(F(r, s), p)$ is complete. Now, someone can readily check that the absolute feature does not verify on the space $l(F(r, s), p)$, that is

(2.13)

$$\begin{aligned} h(x) &= \left(\sum_k \left| r \frac{f_k}{f_{k+1}} x_k + s \frac{f_{k+1}}{f_k} x_{k-1} \right|^{p_k} \right)^{\frac{1}{M}} \neq \left(\sum_k \left| r \frac{f_k}{f_{k+1}} |x_k| + s \frac{f_{k+1}}{f_k} |x_{k-1}| \right|^{p_k} \right)^{\frac{1}{M}} \\ &= h(|x|), \end{aligned}$$

where $|x| = (|x_k|)$. This says that $l(F(r, s), p)$ is the sequence space of non-absolute type. \square

Theorem 2.2. *Convergence in $l(F(r, s), p)$ is strictly stronger than coordinate-wise convergence, but the contrary isn't actual, ordinarily.*

Proof. First, we indicate that $h(x^n - x) \rightarrow 0$, as $n \rightarrow \infty$ purportes $x_k^{(n)} \rightarrow x_k$, as $n \rightarrow \infty$, for all $k \in \mathbb{N}$. If we fix k , then we have

$$\begin{aligned} 0 &\leq \left| \left(r \frac{f_k}{f_{k+1}} x_k^{(n)} + s \frac{f_{k+1}}{f_k} x_{k-1}^{(n)} \right) - \left(r \frac{f_k}{f_{k+1}} x_k + s \frac{f_{k+1}}{f_k} x_{k-1} \right) \right|^{p_k}, \\ &\leq \sum_k \left| \left(r \frac{f_k}{f_{k+1}} x_k^{(n)} + s \frac{f_{k+1}}{f_k} x_{k-1}^{(n)} \right) - \left(r \frac{f_k}{f_{k+1}} x_k + s \frac{f_{k+1}}{f_k} x_{k-1} \right) \right|^{p_k}, \\ &= \sum_k \left| r \frac{f_k}{f_{k+1}} (x_k^{(n)} - x_k) + s \frac{f_{k+1}}{f_k} (x_{k-1}^{(n)} - x_{k-1}) \right|^{p_k}, \\ &= [h(x^n - x)]^M. \end{aligned}$$

Hence, we have for $k = 0$,

$$(2.14) \quad \lim_{n \rightarrow \infty} \left| \left(r \frac{f_0}{f_1} x_0^{(n)} + s \frac{f_1}{f_0} x_{-1}^{(n)} \right) - \left(r \frac{f_0}{f_1} x_0 + s \frac{f_1}{f_0} x_{-1} \right) \right| = 0,$$

that is, $\left| r \frac{f_0}{f_1} [x_0^{(n)} - x_0] \right| \rightarrow 0$, as $n \rightarrow \infty$ and $\frac{f_0}{f_1} = 1$ and $r \frac{f_0}{f_1} = r \neq 0$, then $|x_0^{(n)} - x_0| \rightarrow 0$ as $n \rightarrow \infty$. Likewise, for each $k \in \mathbb{N}$, we have $|x_k^{(n)} - x_k| \rightarrow 0$, as $n \rightarrow \infty$. Now, we demonstrate that the contrary isn't actual, ordinarily. We theorize $x_k^{(n)} \rightarrow x_k$, as $n \rightarrow \infty$. Then, there comes into being an $N \in \mathbb{N}$, such that $|x_k^{(n)} - x_k| < 1$, for each immutable k and for all $n \geq N$. Therefore, we see that,

$$(2.15) \quad \begin{aligned} 0 \leq h(x^n - x) &= \left(\sum_k \left| r \frac{f_k}{f_{k+1}} (x_k^{(n)} - x_k) + s \frac{f_{k+1}}{f_k} (x_{k-1}^{(n)} - x_{k-1}) \right|^{p_k} \right)^{\frac{1}{M}}, \\ &= \left\{ \sum_k \left[\left| r \frac{f_k}{f_{k+1}} (x_k^{(n)} - x_k) + s \frac{f_{k+1}}{f_k} (x_{k-1}^{(n)} - x_{k-1}) \right|^{\frac{p_k}{M}} \right]^M \right\}^{\frac{1}{M}}, \\ &\leq \left\{ \sum_k \left[\left| r \frac{f_k}{f_{k+1}} (x_k^{(n)} - x_k) \right|^{\frac{p_k}{M}} + \left| s \frac{f_{k+1}}{f_k} (x_{k-1}^{(n)} - x_{k-1}) \right|^{\frac{p_k}{M}} \right]^M \right\}^{\frac{1}{M}}, \\ &\leq \left[\sum_k \left| r \frac{f_k}{f_{k+1}} (x_k^{(n)} - x_k) \right|^{p_k} \right]^{\frac{1}{M}} + \left[\sum_k \left| s \frac{f_{k+1}}{f_k} (x_{k-1}^{(n)} - x_{k-1}) \right|^{p_k} \right]^{\frac{1}{M}}, \\ &\leq \left(\sum_k \left| r \frac{f_k}{f_{k+1}} \right|^{p_k} \cdot |x_k^{(n)} - x_k|^{p_k} \right)^{\frac{1}{M}} + \left(\sum_k \left| s \frac{f_{k+1}}{f_k} \right|^{p_k} \cdot |x_{k-1}^{(n)} - x_{k-1}|^{p_k} \right)^{\frac{1}{M}}, \\ &\leq \left(\sum_k \left| r \frac{f_k}{f_{k+1}} \right|^{p_k} \right)^{\frac{1}{M}} + \left(\sum_k \left| s \frac{f_{k+1}}{f_k} \right|^{p_k} \right)^{\frac{1}{M}} \end{aligned}$$

for all k and N . Since $\left| \frac{f_{k+1}}{f_k} \right| \rightarrow 1, 6 \Rightarrow \left| s \frac{f_{k+1}}{f_k} \right| \rightarrow |s|, 6$ and $\left| \frac{f_k}{f_{k+1}} \right| \rightarrow 0, 6 \Rightarrow \left| r \frac{f_k}{f_{k+1}} \right| \rightarrow |r|, 6$ as $k \rightarrow \infty$. In (2.15), $h(x^n - x)$ doesn't convergence for each immutable $k \in \mathbb{N}$ and for all $n \geq N$. This purports that the contrary isn't actual. Let us consider the elements of the sequence x^n be equal, then we follow up that $h(x^n - x) = 0$, that is to say that coordinatewise convergence requires convergence. Hence, we can say that the contrary is not actual, ordinarily. \square

Theorem 2.3. $l(F(r, s), p)$ is a K -space.

Proof. Firstly, we show that $q_i(x) = x_i$ is linear for all $i \in \mathbb{N}$. Let $x = (x_i), y = (y_i) \in l(F(r, s), p)$ and $\alpha \in \mathbb{C}$. Then, we get

$$(2.16) \quad q_i(x + y) = (x + y)_i = x_i + y_i = q_i(x) + q_i(y)$$

and

$$(2.17) \quad q_i(\alpha x) = (\alpha x)_i = \alpha x_i = \alpha q_i(x)$$

for all $i \in \mathbb{N}$. Hence q_i is linear. Now, we substantiate that q_i is continuous. For this, it is sufficient to show that q_i is bounded. Let $x = (x_i) \in l(F(r, s), p)$ be any vector. Then, since $|q_i(x)| = |x_i|$ for all $i \in \mathbb{N}$, one can see that

$$\begin{aligned} \|q_i\| &= \sup_{x \neq 0} \frac{|q_i(x)|}{\|x\|_{l(F(r,s),p)}} = \sup_{x \neq 0} \frac{|x_i|}{\|x\|_{l(F(r,s),p)}}, \\ &\leq \sup_{x \neq 0} \frac{\|x\|_{l(F(r,s),p)}}{\|x\|_{l(F(r,s),p)}} = 1 < \infty, \end{aligned}$$

i.e., q_i is bounded. Hence, q_i is a linear and continuous operator. That is to say that $l(F(r, s), p)$ is a K -space. \square

Theorem 2.4. $l(F(r, s), p)$ is an FK -space.

Proof. It is easy to see by Theorem (2.1) and Theorem (2.2) that $l(F(r, s), p)$ is complete sequence space and convergence requires coordinatewise convergence. Hence, $l(F(r, s), p)$ is an FK -space. \square

Theorem 2.5. $l_p(F(r, s))$ is linear space and a BK -space with the following norm

$$(2.18) \quad \|x\| = \left(\sum_n \left| r \frac{f_n}{f_{n+1}} x_n + s \frac{f_{n+1}}{f_n} x_{n-1} \right|^p \right)^{1/p},$$

where $x = (x_n) \in l_p(F(r, s))$ and $1 \leq p < \infty$.

Proof. it will not be done because the proof of linearity of the space is routine operations. Because l_p is a BK -space with known norm and $F(r, s)$ is a triangle matrix, according to Theorem 4.3.2 of Wilansky [28], we acquire that $l_p(F(r, s))$ is a BK -space. \square

Theorem 2.6. When p fulfilled the condition $1 \leq p < \infty$. The newly specified sequence space C is a BK -space with the norm $\|x\|_{l_p(F(r,s))} = \|f(r, s)x\|_p$, in other words,

$$(2.19) \quad \|x\|_{l_p(F(r,s))} = \left(\sum_n |F_n(r, s)x|^p \right)^{\frac{1}{p}}, \quad 1 \leq p < \infty.$$

Theorem 2.7. $l_p(F(r, s))$ is a Frechet space.

Proof. To avoid the repetition of the similar expressions, we demonstrate that the algebraic operations are continuous in the space $l_p(F(r, s))$. Let (x_n) and (y_n) be two sequences in $l_p(F(r, s))$ and (α_n) be a sequence of scalars such that $d(x_n, x) \rightarrow 0$, $d(y_n, y) \rightarrow 0$ and $\alpha_n \rightarrow \alpha$, as $n \rightarrow \infty$. Then, we get that

$$\begin{aligned}
 (2.20) \quad & 0 \leq \lim_{n \rightarrow \infty} d(x_n + y_n, x + y), \\
 & = \lim_{n \rightarrow \infty} (\|x_n + y_n - (x + y)\|) \\
 & \leq \lim_{n \rightarrow \infty} (\|x_n - x\| + \|y_n - y\|) \\
 & = \lim_{n \rightarrow \infty} d(x_n, x) + \lim_{n \rightarrow \infty} d(y_n, y) \\
 & = 0
 \end{aligned}$$

$$\begin{aligned}
 (2.21) \quad & 0 \leq \lim_{n \rightarrow \infty} d(\alpha_n x_n, \alpha x) \\
 & = \lim_{n \rightarrow \infty} \|\alpha_n x_n - \alpha x\| \\
 & = \lim_{n \rightarrow \infty} \|(\alpha_n - \alpha)x_n + \alpha(x_n - x)\| \\
 & \leq \lim_{n \rightarrow \infty} (|\alpha_n - \alpha| \|x_n\| + |\alpha| \|x_n - x\|) \\
 & = \lim_{n \rightarrow \infty} |\alpha_n - \alpha| \|x_n\| + |\alpha| \lim_{n \rightarrow \infty} d(x_n, x) \\
 & = 0
 \end{aligned}$$

It is ready to see from (2.20) and (2.21) that the algebraic operations are continuous on the linear metric space $l_p(F(r, s))$. Hence $l_p(F(r, s))$ is a Frechet space. \square

With the notation of (2.4), the transformation T specified from $l(F(r, s), p)$ to $l(p)$ by $x \rightarrow y = Tx$ is linear bijection, so we have the following:

Corollary 2.1. *The sequence space $l(F(r, s), p)$ of the non-absolute type is linearly paranorm isomorphic to the space $l(p)$, where $0 < p_k \leq H < \infty$, for all $k \in N$.*

Due to the well known fact that the matrix domain ν_A of the normed sequence space denoted by ν , has got a base iff ν has got a base, whenever a matrix A is a triangle [31](remark 2.4). Accordingly, we can give the following result:

Corollary 2.2. *Let $0 < p_k \leq H < \infty$ and $\lambda_k = (Fx)_k$, for all $k \in N$. Specify the sequence $b^{(k)} = \{b_n^{(k)}\}_{n \in N}$ of the elements of the spaces $l(F(r, s), p)$ by*

$$(2.22) \quad b_n^{(k)} = \begin{cases} \frac{1}{r} \left(-\frac{s}{r}\right)^{k-n} \frac{f_{k+1}^2}{f_n \cdot f_{n+1}}, & 0 \leq n \leq k \\ 0, & n > k \end{cases}$$

for every immutable $k \in N$. Then, the sequence $\{b^{(k)}\}_{k \in N}$ is a basis for the space $l(F(r, s), p)$ and any $x \in l(F(r, s), p)$ has a sole representation of the shape

$$(2.23) \quad x = \sum_k \lambda_k b^{(k)}$$

3. The α -, β -, γ - Duals of the space $l(F(r, s), p)$

The α -, β - and γ duals of the sequence space X are specified as follows:

If x and y are sequences and X and Y are subsets of w , then, we write $x.y = (x_k y_k)_{k=0}^{\infty}$, and

$$(3.1) \quad x^{-1} * Y = \{a \in w : ax \in Y\}$$

$$(3.2) \quad M(X, Y) = \bigcap_{x \in X} x^{-1} * Y = \{a \in w : ax \in Y, \text{ for all } x \in X\},$$

for the multiplier space of X and Y . One can readily follow up for a sequence space Z with $Y \subset Z$ and $M(X, Y) \subset M(X, Z)$ that the containments $M(X, Y) \subset M(Z, Y)$ hold. The α -, β -, and γ - duals of a sequence space X , which are, respectively, denoted by

$$(3.3) \quad X^\alpha = M(X, l_1), X^\beta = M(X, cs) \text{ and } X^\gamma = M(X, bs)$$

It is obvious that $X^\alpha \subset X^\beta \subset X^\gamma$. Also, it can be readily seen that the containments $X^\alpha \subset Y^\alpha$, $X^\beta \subset Y^\beta$ and $X^\gamma \subset Y^\gamma$ hold, whenever $Y \subset X$.

Lemma 3.1. [30] Let $A = (a_{nk})$ be an infinite matrix over the complex field. The following expressions are ensured:

i) Let $0 < p_k \leq 1$ for all $k \in N$. Then, an infinite matrix A transforms all sequences belong to $l(p)$ into l_1 iff

$$(3.4) \quad \sup_{N \in \mathcal{F}} \sup_{k \in N} \left| \sum_{n \in N} a_{nk} \right|^{p_k} < \infty.$$

ii) Let $1 < p_k \leq H < \infty$, for all $k \in N$. Then, an infinite matrix A transforms all sequences belong to $l(p)$ into l_1 iff there comes into being an integer $B > 1$ such that

$$(3.5) \quad \sup_{N \in \mathcal{F}} \sum_k \left| \sum_{n \in N} a_{nk} B^{-1} \right|^{q_k} < \infty.$$

Lemma 3.2. [29] Let $A = (a_{nk})$ be an infinite matrix over the complex field. The following expressions ensure:

i) Let $0 < p_k \leq 1$ for all $k \in N$. Then, an infinite matrix A transforms all sequences belong to $l(p)$ into l_∞ iff

$$(3.6) \quad \sup_{n, k \in N} |a_{nk}|^{p_k} < \infty$$

ii) Let $1 < p_k \leq H < \infty$, for all $k \in N$. Then, an infinite matrix A transforms all sequences belong to $l(p)$ into l_∞ iff there comes into being an integer $B > 1$ such that

$$(3.7) \quad \sup_{n \in N} \sum_k |a_{nk} B^{-1}|^{q_k} < \infty$$

Lemma 3.3. [29] Let $A = (a_{nk})$ be an infinite matrix over the complex field and $0 < p_k \leq H < \infty$ for all $k \in N$. Then, an infinite matrix A transforms all sequences belong to $l(p)$ into c iff (3.6), (3.7) and for all $k \in N$

$$(3.8) \quad \lim_{n \rightarrow \infty} a_{nk} = \beta_k$$

ensures.

Let us specify the following sets:

$$(3.9) \quad E_1 = \left\{ a = (a_k) \in w : \sup_{N \in \mathcal{F}} \sup_{k \in N} \left| \sum_{n \in N} \frac{1}{r} \left(-\frac{s}{r} \right)^{n-k} \frac{f_{n+1}^2}{f_k \cdot f_{k+1}} a_n \right|^{p_k} < \infty \right\},$$

$$(3.10) \quad E_2 = \cup_{B > 1} \left\{ a = (a_k) \in w : \sup_{n \in \mathcal{F}} \sum_k \left| \sum_{n \in N} \frac{1}{r} \left(-\frac{s}{r} \right)^{n-k} \frac{f_{n+1}^2}{f_k \cdot f_{k+1}} a_n \cdot B^{-1} \right|^{q_k} < \infty \right\}$$

$$(3.11) \quad E_3 = \left\{ a = (a_k) \in w : \sup_{k, n \in N} \left| \sum_{j=k}^n \frac{1}{r} \left(-\frac{s}{r} \right)^{j-k} \frac{f_{j+1}^2}{f_k \cdot f_{k+1}} a_j \right|^{p_k} < \infty \right\}$$

$$(3.12) \quad E_4 = \left\{ a = (a_k) \in w : \sum_{j=k}^{\infty} \frac{1}{r} \left(-\frac{s}{r} \right)^{j-k} \frac{f_{j+1}^2}{f_k \cdot f_{k+1}} a_j < \infty \right\}$$

$$(3.13) \quad E_5 = \cup_{B > 1} \left\{ a = (a_k) \in w : \sup_{n \in N} \sum_k \left| \sum_{j=k}^n \frac{1}{r} \left(-\frac{s}{r} \right)^{j-k} \frac{f_{j+1}^2}{f_k \cdot f_{k+1}} a_j \cdot B^{-1} \right|^{q_k} < \infty \right\}$$

Since proof of the first chapter of the following theorems can be made similar to the proof of the second chapter, we will only give proof of the second part.

Theorem 3.1. The following expressions are ensured:

i) Let $0 < p_k \leq 1$ for all $k \in N$. Then,

$$(3.14) \quad \{l(F(r, s), p)\}^\alpha = E_1$$

ii) Let $1 < p_k \leq H < \infty$, for all $k \in N$. Then,

$$(3.15) \quad \{l(F(r, s), p)\}^\alpha = E_2$$

Proof. Let us take any $a = (a_n) \in w$. By employing (2.5) we acquire that, for all $n \in \mathbb{N}$

$$(3.16) \quad a_n x_n = \sum_{k=0}^n \left(\frac{1}{r}\right) \left(-\frac{s}{r}\right)^{n-k} \frac{f_{n+1}^2}{f_k \cdot f_{k+1}} a_n \cdot y_k = (Ey)_n,$$

where $E = (e_{nk})$ is specified by

$$(3.17) \quad e_{nk} = \begin{cases} \left(\frac{1}{r}\right) \left(-\frac{s}{r}\right)^{n-k} \frac{f_{n+1}^2}{f_k \cdot f_{k+1}} a_n, & 0 \leq k \leq n, \\ 0, & k > n, \end{cases}$$

for all $k, n \in \mathbb{N}$. Thus, we follow up by unification (3.16) with the proviso (3.5). of part (ii) of Lemma 3.1 that $ax = (a_n x_n) \in l_1$ whenever $x = (x_k) \in l(F(r, s), p)$ iff $Ey \in l_1$, whenever $y = (y_k) \in l(F(r, s), p)$. This leads to fact that $\{l(F(r, s), p)\}^\alpha =$

E_2 , as claimed. \square

Theorem 3.2. *The following expressions are ensured:*

i) *Let $0 < p_k \leq 1$ for all $k \in \mathbb{N}$. Then,*

$$(3.18) \quad \{l(F(r, s), p)\}^\beta = E_3 \cap E_4.$$

ii) *Let $1 < p_k \leq H < \infty$, for all $k \in \mathbb{N}$. Then,*

$$(3.19) \quad \{l(F(r, s), p)\}^\beta = E_4 \cap E_5.$$

Proof. Take any $a = (a_j) \in w$, then, one can acquire by (2.5) that

$$(3.20) \quad \begin{aligned} \sum_{j=0}^n a_j x_j &= \sum_{j=0}^n \left(\sum_{k=0}^j \frac{1}{r} \left(-\frac{s}{r}\right)^{j-k} \frac{f_{j+1}^2}{f_k \cdot f_{k+1}} y_k \right) a_j \\ &= \sum_{k=0}^n \left(\sum_{j=k}^n \frac{1}{r} \left(-\frac{s}{r}\right)^{j-k} \frac{f_{j+1}^2}{f_k \cdot f_{k+1}} a_j \right) y_k \\ &= (Dy)_n, \end{aligned}$$

for all $n \in \mathbb{N}$, where $D = (d_{nk})$ is specified by

$$(3.21) \quad d_{nk} = \begin{cases} \sum_{j=k}^n \left(\frac{1}{r}\right) \left(-\frac{s}{r}\right)^{j-k} \frac{f_{j+1}^2}{f_k \cdot f_{k+1}} a_j, & 0 \leq k \leq n, \\ 0, & k > n, \end{cases}$$

for all $n, k \in \mathbb{N}$. Thus, we make inferences from Lemma (3.3) with (3.20) that $ax = (a_j x_j) \in cs$ whenever $x = (x_j) \in l(F(r, s), p)$ iff $Dy \in c$, whenever $y = (y_k) \in l(p)$. Therefore, we derive from (3.7) and (3.8) that

$$(3.22) \quad \sup_{n \in \mathbb{N}} \sum_k \left| \sum_{j=k}^n \frac{1}{r} \left(-\frac{s}{r}\right)^{j-k} \frac{f_{j+1}^2}{f_k \cdot f_{k+1}} a_j \cdot B^{-1} \right|^{q_k} < \infty,$$

$$(3.23) \quad \sum_{j=k}^{\infty} \frac{1}{r} \left(-\frac{s}{r}\right)^{j-k} \frac{f_{j+1}^2}{f_k \cdot f_{k+1}} a_j < \infty.$$

This shows that $\{l(F(r, s), p)\}^\beta = E_4 \cap E_5 \quad \square$

Theorem 3.3. *The following expressions are ensured:*

i) Let $0 < p_k \leq 1$ for all $k \in N$. Then,

$$(3.24) \quad \{l(F(r, s), p)\}^\gamma = E_3.$$

ii) Let $1 \leq p_k \leq H < \infty$, for all $k \in N$. Then,

$$(3.25) \quad \{l(F(r, s), p)\}^\gamma = E_5.$$

Proof. From Lemma (3.2) and (3.20), we acquire that $ax = (a_j x_j) \in bs$ whenever $x = (x_j) \in l(F(r, s), p)$ iff $Dy \in l_\infty$, whenever $y = (y_k) \in l(p)$, where $D = (d_{nk})$ is acquired by (3.21). Therefore, we acquire (3.6) and (3.7) that

$$(3.26) \quad \{l(F(r, s), p)\}^\gamma = \begin{cases} E_3, & p_k \leq 1, \\ E_5, & p_k > 1 \end{cases},$$

as desired. \square

4. Some Matrix Transformations on the space $l(F(r, s), p)$

In this section, we characterize some *matrix transformations* on the space $l(F(r, s), p)$, since the cases $0 < p_k \leq 1$ and $1 < p_k \leq H < \infty$ are integrated. The following theorem gives the exact provisos of the general case $0 < p_k \leq H < \infty$. We consider only the case $1 < p_k \leq H < \infty$ and omit the proof of the case $0 < p_k \leq 1$, since it can be ensured in an alike way.

Theorem 4.1. *The following expressions are ensured:*

i) Let $0 < p_k \leq 1$ for all $k \in N$. Then, an infinite matrix A transforms all sequences belong to $l(F(r, s), p)$ into l_∞ iff

$$(4.1) \quad \sup_{k, n \in N} \left| \sum_{j=k}^{\infty} \frac{1}{r} \left(-\frac{s}{r}\right)^{j-k} \frac{f_{j+1}^2}{f_k \cdot f_{k+1}} a_{nj} \right|^{p_k} < \infty,$$

$$(4.2) \quad \sum_{j=k}^{\infty} \frac{1}{r} \left(-\frac{s}{r}\right)^{j-k} \frac{f_{j+1}^2}{f_k \cdot f_{k+1}} a_{nj} < \infty.$$

ii) Let $1 < p_k \leq H < \infty$, for all $k \in N$. Then, an infinite matrix A transforms all sequences belong to $l(F(r, s), p)$ into l_∞ iff (4.1) ensures and there comes into being an integer $B > 1$ such that

$$(4.3) \quad \sup_{n \in N} \sum_k \left| \sum_{j=k}^{\infty} \frac{1}{r} \left(-\frac{s}{r}\right)^{j-k} \frac{f_{j+1}^2}{f_k \cdot f_{k+1}} a_{nj} B^{-1} \right|^{q_k} < \infty.$$

Proof. Let $A \in (l(F(r, s), p) : l_\infty)$ and $1 < p_k \leq H < \infty$, for all $k \in \mathbb{N}$. Then, Ax

comes into being for every $x \in l(F(r, s), p)$ and this implies that $A_n \in \{l(F(r, s), p)\}^\beta$ for each immutable $n \in \mathbb{N}$. Therefore, the necessities of (4.2) and (4.3) are immediate.

Conversely, suppose that the provisos (4.2) and (4.3) ensure, and take any $x \in l(F(r, s), p)$. Since, $A_n \in \{l(F(r, s), p)\}^\beta$, for each $n \in \mathbb{N}$, Ax comes into being. By employing (2.5), we acquire that

$$(4.4) \quad \sum_{j=0}^m a_{nj}x_j = \sum_{j=0}^m \sum_{k=0}^j \frac{1}{r} \left(-\frac{s}{r}\right)^{j-k} \frac{f_{j+1}^2}{f_k \cdot f_{k+1}} y_k a_{nj} = \sum_{k=0}^m \sum_{j=k}^m \frac{1}{r} \left(-\frac{s}{r}\right)^{j-k} \frac{f_{j+1}^2}{f_k \cdot f_{k+1}} a_{nj} y_k,$$

for all $m, n \in \mathbb{N}$. Taking into account the hypothesis, we reproduce from (4.4), as $m \rightarrow \infty$ that for all $n \in \mathbb{N}$,

$$(4.5) \quad \sum_j a_{nj}x_j = \sum_k \sum_{j=k}^{\infty} \frac{1}{r} \left(-\frac{s}{r}\right)^{j-k} \frac{f_{j+1}^2}{f_k \cdot f_{k+1}} a_{nj} y_k,$$

By integrating (4.5) and the inequality which ensures for any complex numbers a, b and any $B > 0$,

$$(4.6) \quad |ab| \leq B \left(|aB^{-1}|^q + |b|^p \right),$$

where $p > 1$ and $p^{-1} + q^{-1} = 1$, we acquire that,

$$\begin{aligned} \sup_{n \in \mathbb{N}} \left| \sum_j a_{nj}x_j \right| &= \sup_{n \in \mathbb{N}} \left| \sum_k \sum_{j=k}^{\infty} \frac{1}{r} \left(-\frac{s}{r}\right)^{j-k} \frac{f_{j+1}^2}{f_k \cdot f_{k+1}} a_{nj} y_k \right| \\ &\leq \sup_{n \in \mathbb{N}} \sum_k \left| \sum_{j=k}^{\infty} \frac{1}{r} \left(-\frac{s}{r}\right)^{j-k} \frac{f_{j+1}^2}{f_k \cdot f_{k+1}} a_{nj} y_k \right|, \\ &\leq \sup_{n \in \mathbb{N}} \sum_k B \left(\left| \sum_{j=k}^{\infty} \frac{1}{r} \left(-\frac{s}{r}\right)^{j-k} \frac{f_{j+1}^2}{f_k \cdot f_{k+1}} a_{nj} B^{-1} \right|^{q_k} + |y_k|^{p_k} \right), \\ &= B \left(\sup_{n \in \mathbb{N}} \sum_k \left| \sum_{j=k}^{\infty} \frac{1}{r} \left(-\frac{s}{r}\right)^{j-k} \frac{f_{j+1}^2}{f_k \cdot f_{k+1}} a_{nj} B^{-1} \right|^{q_k} + \sup_{n \in \mathbb{N}} \sum_k |y_k|^{p_k} \right) \\ &< \infty. \end{aligned}$$

This shows that $Ax \in l_\infty$. \square

Theorem 4.2. *The following expressions are ensured:*

i) Let $0 < p_k \leq 1$ for all $k \in \mathbb{N}$. Then, an infinite matrix A transforms all sequences belong to $l(F(r, s), p)$ into c iff (4.1) and (4.2) ensure and there is a sequence $\alpha = (\alpha_k)$ of scalars such that

$$(4.7) \quad \lim_{n \rightarrow \infty} \sum_{j=k}^{\infty} \frac{1}{r} \left(-\frac{s}{r}\right)^{j-k} \frac{f_{j+1}^2}{f_k \cdot f_{k+1}} a_{nj} = \alpha_k,$$

for all $k \in \mathbb{N}$.

ii) Let $1 < p_k \leq H < \infty$, for all $k \in \mathbb{N}$. Then, an infinite matrix A transforms all sequences belong to $l(F(r, s), p)$ into c iff (4.2), (4.3) and (4.7), ensure.

Proof. Let $A \in (l(F(r, s), p) : c)$ and $1 < p_k \leq H < \infty$ for all $k \in \mathbb{N}$. Then, since the containment $c \subset l_\infty$ ensures, the necessities of (4.2) and (4.3) are immediately acquired from Theorem (4). To ensure the the necessities of (4.7), consider the sequence $b^{(k)}$ defined by (2.22) which belongs to the space $l(F(r, s), p)$ for every immutable $k \in \mathbb{N}$. Since, the A - transform of every $x \in l(F(r, s), p)$ comes into being and is in c by the hypothesis, we have

$$(4.8) \quad Ab^{(k)} = \left(\sum_{j=0}^{\infty} a_{ij} b_j^{(k)} \right)_{i=0}^{\infty} = \left(\sum_{j=k}^{\infty} \frac{1}{r} \left(-\frac{s}{r}\right)^{j-k} \frac{f_{j+1}^2}{f_k \cdot f_{k+1}} a_{ij} \right)_{i=0}^{\infty} \in c,$$

for every fixed $k \in \mathbb{N}$, which shows the necessity (4.7).

Conversely, suppose that the provisos (4.2), (4.3) and (4.7) ensure and take $x = (x_k)$ in the space $l(F(r, s), p)$. Then, Ax exists. We follow up for all $m, n \in \mathbb{N}$ that

$$\begin{aligned} & \sum_{k=0}^m \left| \sum_{j=k}^n \frac{1}{r} \left(-\frac{s}{r}\right)^{j-k} \frac{f_{j+1}^2}{f_k \cdot f_{k+1}} a_{nj} B^{-1} \right|^{q_k} \\ & \leq \sup_{n \in \mathbb{N}} \sum_k \left| \sum_{j=k}^{\infty} \frac{1}{r} \left(-\frac{s}{r}\right)^{j-k} \frac{f_{j+1}^2}{f_k \cdot f_{k+1}} a_{nj} B^{-1} \right|^{q_k} < \infty \end{aligned}$$

which gives the fact that by letting $m, n \rightarrow \infty$ with (4.3) and (4.7)

$$\begin{aligned} & \lim_{m, n \rightarrow \infty} \sum_{k=0}^m \left| \sum_{j=k}^n \frac{1}{r} \left(-\frac{s}{r}\right)^{j-k} \frac{f_{j+1}^2}{f_k \cdot f_{k+1}} a_{nj} B^{-1} \right|^{q_k} \\ & \leq \sup_{n \in \mathbb{N}} \sum_k \left| \sum_{j=k}^{\infty} \frac{1}{r} \left(-\frac{s}{r}\right)^{j-k} \frac{f_{j+1}^2}{f_k \cdot f_{k+1}} a_{nj} B^{-1} \right|^{q_k} < \infty. \end{aligned}$$

This shows that $\sum_k |\alpha_k B^{-1}|^{q_k} < \infty$ and $(\alpha_k) \in \{l(F(r, s), p)\}^\beta$ which implies that the series $\sum_k \alpha_k x_k$ converges for all $x \in \{l(F(r, s), p)\}$.

Now, let us consider the equality acquired from 4.5 with $(a_{nj} - \alpha_j)$ instead of a_{nj}

$$(4.9) \quad \sum_j (a_{nj} - \alpha_j) x_j = \sum_k \sum_{j=k}^{\infty} \frac{1}{r} \left(-\frac{s}{r}\right)^{j-k} \frac{f_{j+1}^2}{f_k \cdot f_{k+1}} (a_{nj} - \alpha_j) y_k = \sum_k c_{nk} y_k,$$

where $c = (c_{nk})$ described by $c_{nk} = \sum_{j=k}^{\infty} \frac{1}{r} \left(-\frac{s}{r}\right)^{j-k} \frac{f_{j+1}^2}{f_k \cdot f_{k+1}} (a_{nj} - \alpha_j)$ for all $k, n \in \mathbb{N}$. From Lemma 3.3, $c_{nk} \rightarrow 0$, as $n \rightarrow \infty$. This means that $Ax \in c$, whenever $x \in l(F(r, s), p)$ and this step completes the proof. \square

Corollary 4.1. *The following expressions are ensured:*

i) Let $0 < p_k \leq 1$ for all $k \in \mathbb{N}$. Then, an infinite matrix A transforms all sequences belong to $l(F(r, s), p)$ into c_0 iff (4.1), and (4.2) ensure and (4.7) ensures with $\alpha_k = 0$, for all $k \in \mathbb{N}$.

ii) Let $1 < p_k \leq H < \infty$, for all $k \in \mathbb{N}$. Then, an infinite matrix A transforms all sequences belong to $l(F(r, s), p)$ into c_0 iff (4.1), and (4.2) ensure and (4.7) also ensures with $\alpha_k = 0$, for all $k \in \mathbb{N}$.

Lemma 4.1. [1] Let λ, μ be any two sequence spaces, A be an infinite matrix and B be a triangle matrix. Then, $A \in (\lambda : \mu_B)$ iff $BA \in (\lambda : \mu)$.

By considering Theorem , Theorem and Lemma together, it can be acquired following outcome:

Corollary 4.2. *Let $A = (a_{nk})$ be an infinite matrix of complex constituent. Then, the following expressions are ensured:*

i) $E = (e_{nk}) \in (l(F(r, s), p) : bv_{\infty})$ iff (4.1), (4.3) ensure with d_{nk} instead of a_{nk} ; where $d_{nk} = e_{nk} - e_{n-1, k}$ for all $k, n \in \mathbb{N}$.

ii) $E = (e_{nk}) \in (l(F(r, s), p) : e_{\infty}^r)$ iff (4.1), (4.3) ensure with d_{nk} instead of a_{nk} ; where $d_{nk} = \sum_{j=0}^n \binom{n}{j} (1-r)^{n-j} r^j e_{jk}$ for all $k, n \in \mathbb{N}$.

iii) $E = (e_{nk}) \in (l(F(r, s), p) : X_{\infty})$ iff (4.1), (4.3) ensure with d_{nk} instead of a_{nk} ; where $d_{nk} = \sum_{j=0}^n \frac{e_{jk}}{(n+1)}$ for all $k, n \in \mathbb{N}$.

iv) $E = (e_{nk}) \in (l(F(r, s), p) : r_{\infty}^t)$ iff (4.1), (4.3) ensure with d_{nk} instead of a_{nk} ; where $d_{nk} = \sum_{j=0}^n t_j e_{jk} / T_n$ for all $k, n \in \mathbb{N}$.

v) $E = (e_{nk}) \in (l(F(r, s), p) : bs)$ iff (4.1), (4.3) ensure with d_{nk} instead of a_{nk} ; where $d_{nk} = \sum_{j=0}^n e_{jk}$ for all $k, n \in \mathbb{N}$.

vi) $E = (e_{nk}) \in \left(l(F(r, s), p) : l_{\infty} \left(\widehat{F} \right) \right)$ iff (4.1), (4.3) ensure with d_{nk} instead of a_{nk} ; where $d_{nk} = s \frac{f_{n+1}}{f_n} e_{n-1, k} + r \frac{f_n}{f_{n+1}} e_{nk}$ for all $k, n \in \mathbb{N}$.

Corollary 4.3. *Let $A = (a_{nk})$ be an infinite matrix of complex constituent. Then, the following expressions are ensured:*

i) $E = (e_{nk}) \in (l(F(r, s), p) : c(\Delta))$ iff (4.1), (4.3) and (4.5) with d_{nk} instead of a_{nk} ; where $d_{nk} = e_{nk} - e_{n+1,k}$ for all $k, n \in N$.

ii) $E = (e_{nk}) \in (l(F(r, s), p) : e_c^r)$ iff (4.1), (4.3) and (4.5) also ensures with d_{nk} instead of a_{nk} ;

where $d_{nk} = \sum_{j=0}^n \binom{n}{j} (1-r)^{n-j} r^j e_{jk}$ for all $k, n \in N$.

iii) $E = (e_{nk}) \in (l(F(r, s), p) : \tilde{c})$ iff (4.1), (4.3) and (4.5) also ensures with d_{nk} instead of a_{nk} ; where $d_{nk} = \sum_{j=0}^n \frac{e_{jk}}{(n+1)}$ for all $k, n \in N$.

iv) $E = (e_{nk}) \in (l(F(r, s), p) : r_c^t)$ iff (4.1), (4.3) and (4.5) also ensures with d_{nk} instead of a_{nk} ; where $d_{nk} = \sum_{j=0}^n t_j e_{jk} / T_n$ for all $k, n \in N$.

v) $E = (e_{nk}) \in (l(F(r, s), p) : c(\hat{F}))$ iff (4.1), (4.3) and (4.5) also ensures with d_{nk} instead of a_{nk} ; where $d_{nk} = s \frac{f_{n+1}}{f_n} e_{n-1,k} + r \frac{f_n}{f_{n+1}} e_{nk}$ for all $k, n \in N$.

vi) $E = (e_{nk}) \in (l(F(r, s), p) : cs)$ iff (4.1), (4.3) and (4.5) also ensures with d_{nk} instead of a_{nk} ; where $d_{nk} = \sum_{j=0}^n e_{jk}$ for all $k, n \in N$.

Corollary 4.4. Let $A = (a_{nk})$ be an infinite matrix of complex constituent. Then, the following expressions are ensured:

i) $E = (e_{nk}) \in (l(F(r, s), p) : c_0(\Delta))$ iff (4.1), (4.3) and (4.5) also ensures with $\alpha_k = 0$, for all $k \in N$ and d_{nk} instead of a_{nk} ; where $d_{nk} = e_{nk} - e_{n+1,k}$ for all $k, n \in N$.

ii) $E = (e_{nk}) \in (l(F(r, s), p) : e_0^r)$ iff (4.1), (4.3) and (4.7) also ensures with $\alpha_k = 0$ for all $k \in N$ and d_{nk} instead of a_{nk} ; where $d_{nk} = \sum_{j=0}^n \binom{n}{j} (1-r)^{n-j} r^j e_{jk}$ for all $k, n \in N$.

iii) $E = (e_{nk}) \in (l(F(r, s), p) : \tilde{c}_0)$ iff (4.1), (4.3) and (4.7) also ensures with $\alpha_k = 0$ for all $k \in N$ and d_{nk} instead of a_{nk} ; where $d_{nk} = \sum_{j=0}^n \frac{e_{jk}}{(n+1)}$ for all $k, n \in N$.

iv) $E = (e_{nk}) \in (l(F(r, s), p) : r_0^t)$ iff (4.1), (4.3) and (4.7) also ensures with $\alpha_k = 0$ for all $k \in N$ and d_{nk} instead of a_{nk} ; where $d_{nk} = \sum_{j=0}^n t_j e_{jk} / T_n$ for all $k, n \in N$.

v) $E = (e_{nk}) \in (l(F(r, s), p) : c_0(\hat{F}))$ iff (4.1), (4.3) and (4.7) also ensures with $\alpha_k = 0$ for all $k \in N$ and d_{nk} instead of a_{nk} ; where $d_{nk} = s \frac{f_{n+1}}{f_n} e_{n-1,k} + r \frac{f_n}{f_{n+1}} e_{nk}$ for all $k, n \in N$.

vi) $E = (e_{nk}) \in (l(F(r, s), p) : c_0s)$ iff (4.1), (4.3) and (4.7) also ensures with $\alpha_k = 0$ for all $k \in N$ and d_{nk} instead of a_{nk} ; where $d_{nk} = \sum_{j=0}^n e_{jk}$ for all $k, n \in N$.

REFERENCES

1. F. BAŞAR and B. ALTAY: On the spaces of sequences of p -bounded variation and related matrix mappings. Ukrainian Math. J. **55:1** (2003),136–147.

2. P. N. NG. and P. Y. LEE: *TCesaro sequence spaces of non-absolute type*. Comment. Math. Prace Mat. **20:2** (1978), 429–433.
3. M. CANDAN and E. E. KARA: *A Study On topological and geometrical characteristics of new Banach Sequence Spaces*. Gulf Journal Of Mathematics **3:4** (2015), 67–84.
4. E. E. KARA: *Some topological and geometrical properties of new Banach sequence spaces*. J. Inequal. Appl. **38** (2013), 1–15.
5. H. KIZMAZ: *On certain sequence spaces*. Canad. Math. Bull. **24:2** (1981), 169–176.
6. B. ALTAY, F. BAŞAR and M. MURSALEEN : *On the Euler sequence spaces which include the spaces l_p and l_∞* . Inform. Sci. **176:10** (2006), 1450–1462.
7. B. ALTAY and F. BAŞAR : *On the Paranormed Riesz sequence spaces of non-absolute type*. Southeast Asian Bull. Math. **26:5** (2002), 701–715.
8. B. ALTAY and F. BAŞAR : *Some Euler sequence spaces of non-absolute type*. Ukrainian Math. J. **57:1** (2005), 1–17
9. M. ŞENGÖNÜL and F. BAŞAR : *Some new Cesaro sequence spaces of non-absolute type which include the spaces c_0 and c* . Soochow J. Math. **31:1** (2005), 107–119.
10. B. ALTAY and F. BAŞAR : *Some Paranormed Riesz sequence spaces of non-absolute type*. Southeast Asian Bull. Math. **30:5** (2006), 591–608.
11. M. BAŞARIR, F. BAŞAR and E. E. KARA: *On the spaces of Fibonacci Difference null and convergent sequences*. arXiv:1309.0150v1 [math. FA], 2013.
12. B. CHOUDHARY and S. NANDA: *Functional Analysis with Applications*. JohnWiley , Sons, NewDelhi, India, 1989.
13. I. J. MADDOX: *Spaces of strongly summable sequences*. Quart. J. Math. Oxford. **18:2** (1967), 345–355.
14. S. SIMONS: *The sequence spaces $\ell(p_v)$ and $m(p_v)$* . Proc. London Math. Soc. **15:3** (1965), 422–436.
15. H. NAKANO: *Modulared sequence spaces*. Proc. Japan Acad. **27:2** (1951), 508–512.
16. M. MURSALEEN: *On some geometric properties of a sequence space related to l_p* . Bull. Aust. Math. Soc. **67** (2003), 343–347.
17. C. AYDIN and F. BAŞAR: *Some new sequence spaces which include the spaces l_p and l_∞* . Demonstr. Math. **38** (2005), 641–656.
18. B. ALTAY and F. BAŞAR : *The matrix domain and the fine spectrum of the difference operator Δ on the sequence space l_p ; ($0 < p < 1$)*. Commun. Math. Anal. **2** (2007), 1–11.
19. F. BAŞAR: *Summability Theory and Its Applications*,. Bentham Science Publishers, e-books, Monographs, xi+405 pp., İstanbul, (2012) ISB:978-1-60805-252-3,
20. M. MURSALEEN and A. K. NOMAN : *On some new sequence spaces of non-absolute type related to the spaces l_p and l_∞ ,I*. Filomat. **25:2** (2011), 33–51.
21. M. MURSALEEN and A. K. NOMAN: *On some new sequence spaces of non-absolute type related to the spaces l_p and l_∞ ,II*. Mathematical Communications. **16:2** (2011), 383–398.
22. M. CANDAN and K. KAYADUMAN: *Almost convergent sequence space derived by generalized Fibonacci matrix and Fibonacci core*. Brithish J. Math. Comput. Sci **7:2** (2015), 150–167.

23. M. CANDAN: *A new sequence space isomorphic to the space $l(p)$ and compact operators*. J. Math. Comput. Sci **4:2** (2014), 306–334.
24. M. CANDAN: *A new approach on the spaces of generalized Fibonacci difference null and convergent sequences*. Math. Aeterna. **1:5** (2015), 191–210.
25. A. ALOTAIBI, M. MURSALEEN, B. A. ALAMRI and S. A. MOHIUDDINE : *Compact operators on some Fibonacci difference sequence spaces*. Journal of Inequalities and Applications. (2015), 2015:203 DOI10.1186s13660-015-0713-5.
26. M. CANDAN: *Domain of Double Sequential Band Matrix in the Spaces of Convergent and Null Sequences*. Advanced in Difference Equations. **1** (2014), 1–18.
27. E. MALKOWSKY and V. RAKOČEVIĆ: *An Introduction into the theory of sequence spaces and measures of noncompactness*. Zbornik Radova, Matematički Institut SANU, Belgrade, **9:17** (2000), 143–234.
28. A. WILANSKY: *Functional Analysis*. Blaisdell Publishing Company, New York-Toronto-London, 1964.
29. C. G. LASCARIDES and I. J. MADDOX: *Matrix transformations between some classes of sequences*. Proc. Camb. Phil. Soc. **68** (1970), 99–104.
30. K. G. GROSSE- ERDMAN: *Matrix transformations between the sequence spaces of Maddox*. J. Math. Anal. Apply **180** (1993), 223–238.
31. A. JARRAH and E. MALKOWSKY: *BK-spaces, bases and linear operators*. Rendiconti Circ. Mat. Palermo II **52** (1990), 177–191.
32. H. ÇAPAN and F. BAŞAR: *Domain of The Double Band Matrix Defined By Fibonacci Numbers in The Maddox's Space $l(p)$* Electronic Journal of Mathematical Analysis and Applications **3:2** (2015), 31–45.
33. M. CANDAN: *Almost convergence and double sequential band matrix*. Acta Math. Scientia. **34:2** (2014), 354–366.
34. M. CANDAN and A. GÜNEŞ: *Paranormed sequence spaces of Non Absolute Type Founded Using Generalized Difference Matrix*. Proceedings of the National Academy of Sciences; India Section A:Physical Sciences. **85:2** (2014), 269–276.

Gülşen Kılınç
Faculty of Education
Department of Elementary Education
Adiyaman University,
The University Campus,
02040-Adiyaman/TURKEY
gkilinc@adiyaman.edu.tr