

## APPENDICITIS IN CHILDREN: SYMPTOMS AND SIGNS, LABORATORY AND HISTOPATHOLOGY FINDINGS IN 67 PATIENTS

Dragoljub Živanović<sup>1,2</sup>, Ivona Đorđević<sup>1,2</sup>, Milan Petrović<sup>1\*</sup>

<sup>1</sup>University of Niš Faculty of Medicine, Niš, Serbia

<sup>2</sup>Clinical Center Niš, Clinic for Pediatric Surgery and Orthopedics, Niš, Serbia

**Abstract.** Acute abdominal pain is a reason for hospital admission of about 20% of children. Typical clinical presentation of appendicitis may be significantly different in children. Diagnosis is based on the combination of symptoms, clinical signs, and results of laboratory and radiology examinations. The objective of the present study was to analyze symptoms, signs, laboratory and histopathology findings in children who underwent surgery for acute appendicitis. Sixty-seven patients (37 males and 30 females) with mean age of 9.77 years, operated on for clinical diagnosis of acute appendicitis were enrolled in the study. Abdominal pain was present in all patients, followed by vomitus and fever. Laboratory markers of inflammation varied significantly with severity of inflammation, but were normal in chronic appendicitis. Clinical and histopathology assessments of inflammation were concordant in 22 – 43% depending of the degree of appendicitis. Perforation occurred in 26.86% and negative appendectomy rate was 6%.

**Key words:** appendicitis, children, appendectomy, histopathology.

### Introduction

Acute abdominal pain constitutes 20% of all pediatric surgery admissions [1]. Typical clinical presentation of appendicitis that includes pain, nausea, vomiting and low-grade fever sometimes accompanied by constipation or diarrhea may significantly differ in children [2]. A child may present with very few symptoms and in good general health or in very severe general condition with signs of septic shock and/or bowel obstruction. Decision making is based on the clinical course, physical signs, laboratory and radiology findings. Overall rupture rates varied from 20% to 76%, with a median of 36% in a recent analysis of data from 30 pediatric hospitals in the United States [3]. Perforation rate is reported to be higher in children younger than 5, and is nearly 100% in those younger than 3 years [4,5]. Delay in diagnosis is associated with increased risk of perforation, and it correlates more with pre-hospital than with in-hospital delay [6,7].

Despite the improvement in diagnosis, some of the appendices removed under high clinical suspicion of acute appendicitis turn out to be normal on histopathology (negative appendectomies - NA). Recent researches revealed that some of them had manifested the signs of

inflammation on more detailed examination or on immunohistochemistry [8–10].

White blood cell (WBC) count, percentage of neutrophils (Ne%) and C-reactive protein level (CRP) are the most commonly used laboratory markers [11,12], with combined specificity of 95% for acute and 100% for perforated appendicitis [13]. Recently, neutrophil/lymphocyte ratio (NLR) was reported as more reliable marker of acute appendicitis than WBC count [14], and values of NLR higher than 3.5 were strongly associated with acute appendicitis.

Objective of the present study was to analyze symptoms and signs, laboratory and histopathology findings in children who underwent surgery for acute appendicitis.

### Materials and Methods

Sixty-seven patients, operated on for clinical diagnosis of acute appendicitis, were enrolled in the study.

Data on patients' demographics, history, symptoms and signs at presentation, physical and laboratory findings were obtained from the medical records. Clinical classification of the degree of inflammation made by treating surgeon has been obtained from operative reports. All removed appendices were routinely sent to histopathology examination. The histopathology classification was based on pathologists' reports. Pathologists were blinded for clinical classification of the degree of inflammation.

Collected data were entered in prepared MS Excel 2007© spread sheet. Statistical analysis was performed using Graph Pad Prism 5© statistical software.

Correspondence to: Dragoljub Živanović, M.D., PhD  
Faculty of Medicine, University of Niš  
81 Dr. Zorana Đinđića, 18000 Niš, Serbia  
Phone: +381 64 6389 999  
E-mail: dzivanovic.nis@gmail.com  
Received May 9<sup>th</sup>, 2017, accepted for publication September 26<sup>th</sup>, 2017  
<sup>\*</sup>PhD Student

## Results

Mean age of patients was  $9.77 \pm 2.88$  years (range 4.25 – 15.75). There were 37 males and 30 females. Patients' demographics are summarized in Table 1. Forty-one (61%) patients came from urban, and 26 (39%) from the rural areas. Forty-nine (73%) patients were from local community while 18 patients (27%) had been referred from the surrounding secondary level regional general hospitals (RGH). Ten out of 18 referred patients (55%) presented with perforation.

All patients complained of abdominal pain. Symptoms and their duration are summarized in Table 2. Forty-three patients vomited before admission. Thirty-four patients had normal body temperature. After admission, only 10 patients remained afebrile. Seventeen patients presented with constipation and 11 with diarrhea. Table 3 summarizes the presence of the signs of appendicitis.

Mean value of WBC count was  $16.24 \times 10^3/\mu\text{L}$ . Mean percentage of neutrophils was 73.79% and average NLR ratio was 6.08. Mean value of CRP was 54.38 mg/ml (Table 4).

Time from admission to operation varied with severity of inflammation (Figure 1). Clinical and histopathology diagnoses, as well as concordance of clinical and histopathology assessment of the degree of inflammation are given in Table 5 and Figure 2. Pediatric surgeon's and pathologist's reports were concordant in 35% of acute, 43% of phlegmonous, 33% of gangrenous and 22% of perforated appendices. Overall concordance in clinical and histopathology reports was 33.34% (21 of 67 patients). According to Cohen's kappa coefficient ( $\text{Kappa} = 0.141$ ), the strength of agreement was considered to be poor.

**Table 1** Demographic data of patients operated on with diagnosis of acute appendicitis

Histopathology grade of inflammation	Age (mean $\pm$ SD)	Gender		Living area		Referral	
		Male	Female	Urban	Rural	Local	RGH
Acute	$8.88 \pm 2.81$	4	6	2	4	6	4
Phlegmonous	$9.38 \pm 3.09$	16	15	19	12	23	8
Gangrenous	$10.78 \pm 2.52$	9	4	8	5	9	4
Perforated	$10.80 \pm 2.23$	4	1	1	4	3	2
Chronic	$11.54 \pm 1.65$	1	3	3	1	4	0
Normal (NA)	$8.73 \pm 3.56$	3	1	4	0	4	0
Total	$9.77 \pm 2.88$	37	30	41	26	49	18

**Table 2** Symptoms of appendicitis and its duration

Duration	Symptoms							
	Vomitus		Body temperature					
	Yes	No	Before admission			After admission		
			< 36.9	37.0-37.9	>38.0	< 36.9	37.0-37.9	>38.0
<6h	3	5	6	1	1	1	6	1
6 - 12h	7	1	4	1	3	1	4	3
12 - 24h	23	7	17	9	4	4	17	9
24 - 48 h	6	5	2	2	7	1	3	7
>48h	4	6	5	3	2	3	3	4
Total	43	24	34	16	17	10	33	24

**Table 3** Presence of signs of appendicitis


Sign	no	yes	n/a	(%)
Abdominal tenderness	0	67	0	100.0
Rebound tenderness (Blumberg's sign)	16	28	23	41.8
Localized guarding	59	8	0	11.9
Diffuse guarding (defense)	58	9	0	13.4
Diarrhea	53	11	3	16.4
Constipation	47	17	3	25.4

n/a – not applicable (presence or absence of sign was not recorded)


**Table 4** Mean values of laboratory markers of inflammation (WBC, % of neutrophils, Ne/Ly ratio and CRP) in study groups and in different grades of appendicitis

Histopathology grade of inflammation	Laboratory marker of inflammation (mean ± SD)			
	WBC count (n x 10 <sup>3</sup> /μL)	Ne% <sup>†</sup>	NLR	CRP <sup>§</sup> (mg/ml)
Acute	13.55 ± 6.23	67.53 ± 11.92 <sup>‡</sup>	7.1 ± 10.92	35.40 ± 44.07
Phlegmonous	17.36 ± 6.39	78.08 ± 9.74 <sup>‡, §</sup>	6.0 ± 3.20 <sup>€</sup>	49.85 ± 71.68
Gangrenous	16.45 ± 4.03	76.41 ± 18.65	6.8 ± 3.17 <sup>€</sup>	66.41 ± 67.28
Perforated	22.34 ± 7.88	81.60 ± 11.58 <sup>¥</sup>	16.2 ± 10.87	66.41 ± 67.28
Chronic	8.175 ± 1.87 <sup>*</sup>	61.45 ± 9.85 <sup>§, ¥</sup>	2.2 ± 1.31 <sup>€, £</sup>	13.41 ± 11.98
Normal (NA)	14.00 ± 9.34	68.15 ± 14.91	3.9 ± 2.74	55.86 ± 67.26
Total	16.24 ± 6.59	73.79 ± 10.41	6.1 ± 4.98	54.38 ± 67.45

WBC – White blood cells; Ne% – percent of neutrophils; NLR – Neutrophil to lymphocyte ratio; CRP – C-reactive protein; NA – negative appendectomy. \*Mean WBC count was significantly lower in chronic appendicitis (p<0.05). †Differences in Ne% among histology groups were significant (p<0.05; One-way ANOVA) ‡p<0.05, §p<0.01; ¥p<0.05. Significant differences in NLR were observed between phlegmonous and chronic € p<0.05 and gangrenous and chronic appendicitis £ p<0.05. § Differences in mean values of CRP were not significant (p>0.05).

**Fig. 1** Time from admission to surgery in relation to intensity of inflammation**Table 5** Distribution of clinical and histopathology degree of inflammation and concordance between clinical and histopathology findings.

Degree of inflammation	Clinical diagnosis		Histopathology diagnosis					Concordant diagnoses		
	No	%	Acute	Phlegmonous	Gangrenous	Perforated	Chronic	Normal (N/A)	No	%
Acute	20	29.85	6	9	0	0	1	4	6	30.00
Phlegmonous	14	20.90	4	6	1	0	3	0	6	42.86
Gangrenous	15	22.39	0	9	5	1	0	0	5	33.33
Perforated	18	26.87	0	7	7	4	0	0	4	22.22
Chronic	0	0.00	0	0	0	0	0	0	0	0.00
Normal (N/A)	0	0.00	0	0	0	0	0	0	0	0.00
Total	67	100	10	31	13	5	4	4	21	
%			14.93	46.27	19.40	7.46	5.97	5.97	31.34	



**Fig. 2** Histopathology diagnoses in different groups of clinically assessed degree of inflammation and concordance of clinical and histology reports. A – Acute appendicitis, concordance 30% ; B – Phlegmonous appendicitis, concordance 43%; C – Gangrenous appendicitis, concordance 33%; D – Perforated appendicitis, concordance 22%.

## Discussion

Appendicitis is the most common reason for emergency abdominal surgery in children [15] with peak incidence in the second and third decade of life. Average age of patients in our study was 9.77 years. Appendicitis is rare in children younger than 5 and most of them present with perforation. The reasons for low incidence of appendicitis in young children are not clear, but distinct anatomical, pathophysiological and social characteristics of children contribute to high perforation rate in children younger than 5 years. Four of our patients (5.97%) were in that age group and half of them had perforations. Slight male preponderance was observed (male: female ratio 1.23), with the highest predominance of boys in group of perforated appendicitis (72%).

### Symptoms and signs

Abdominal pain was a universally present symptom. Most frequently, it was located in the right lower quadrant. Description of its migratory nature had not been easily obtained, particularly in younger children. Common duration of pain was 12-24h before admission. Very short duration of pain, less than 6 hours, was associated with more severe inflammation, while duration of pain more than 48h corresponded to either acute or per-

forated appendicitis. These findings slightly differ from those of Rothrock and Pagane [16]. Recent suggestions that acute appendicitis in fact may be the result of repeated episodes of inflammation and that acute and perforated appendicitis might be different entities may support our findings [6,17]. Vomiting had low sensitivity and specificity in limited number of reports on pediatric patients [18,19]. Two-thirds of our patients vomited. Most of the patients with 3-5 episodes of vomitus had ruptured or gangrenous appendicitis. Less than 2 or more than 5 episodes of vomitus corresponded with acute inflammation.

O'Shea et al. [20] reported sensitivity of 0.75 and specificity 0.78 for fever as a symptom of appendicitis, while Andersson et al. [21] found that body temperature provided important diagnostic information, particularly in advanced appendicitis. Normal body temperature from onset of symptoms to admission was present in 52% of our patients. Nevertheless, 85% of them had fever after admission. Ten patients (15%) remained afebrile during hospital course but histopathology showed only one normal appendix (NA).

All children in our series had abdominal tenderness in right lower quadrant (RLQ). Rebound tenderness (Blumberg's sign) was positive in 28 patients (41.79%). In adults, rebound tenderness is the single most accurate physical sign, with accuracy of 86% [18], but its accu-

racy decreases with age to 43% in pediatric patients [18,21]. Localized guarding in RLQ was present in 8 children (11.94%) and 6 of them had clinically perforated appendicitis.

### Laboratory markers of inflammation

Mean values of WBC were elevated in patients with acute inflammation and in negative appendectomies but were normal in children with chronic appendicitis ( $p < 0.05$ ; one-way ANOVA; Table 4). Values of Ne% were significantly different among histology groups ( $p = 0.05$ ; one-way ANOVA; Table 4). Mean values of Ne% were statistically different between acute and phlegmonous ( $p < 0.05$ ), phlegmonous and chronic ( $p < 0.01$ ) and perforated and chronic ( $p < 0.05$ ) appendicitis. Differences in NLR were statistically different between groups with phlegmonous and chronic ( $p < 0.05$ ) and gangrenous and chronic ( $p < 0.05$ ) inflammation. Differences in values of CRP were not significant ( $p > 0.05$ ; one-way ANOVA; Table 4) Three out of 4 patients with chronic appendicitis had normal WBC, Ne% and NLR, but only one had normal CRP. Among children with negative appendectomy, all inflammatory markers were normal in 2 children and 2 had high levels of WBC and CRP. Results of Siddique et al. [13] as well as our findings, demonstrated that mean values of WBC and CRP increase with severity of inflammation. However, both markers are general markers of inflammation, not specific for appendicitis. Nevertheless, appendicitis cannot be ruled out solely on the basis of normal laboratory findings, in the presence of other suggestive clinical signs.

Accurate assessment of the degree of inflammation has implication on postoperative therapy protocol and complication rate. Overall reported accuracy of macroscopic assessment of the appendix at surgery was 87.3% [22] with 100% consensus for perforation, abscess, or gangrene of the appendix. Conversely, Bliss et al. [23] reported greater agreement of surgeon's and pathologist's assessment for normal and acutely inflamed appendices than for more advanced forms. In our study, pediatric surgeons tend to underestimate acute and overestimate more advanced forms of appendicitis (Figure 2). Clinical diagnosis of perforation was established in 26.86% of patients, with histopathology confirmation in 7.46% (22% of concordance). The remaining 14 clinically perforated appendices were histologically either gangrenous or phlegmonous (39% each). Surgeons tend to classify appendix as perforated in the presence of turbid fluid in abdomen, even if the actual hole in appendix wall has not been visualized. On the other hand, pathologists may omit small holes particularly if the whole appendix has not been examined. Strict definition of perforation as a hole in appendix or free appendicolith in abdomen may improve concordance of surgeon's and pathologist's

reports in advanced forms of appendicitis [24]. Diagnosis of chronic appendicitis and negative appendectomy could not be established without histopathology.

Time from admission to operation varied with the severity of inflammation from average of 10h 49 min for acute to 7h 45 min for perforated appendicitis. Patients with negative appendectomies were operated on 15h 36 min and those with chronic inflammation 21h 36 min after admission. Narsule et al. [6] reported similar results. In patients with milder or equivocal symptoms and signs repeated examinations, laboratory tests and imaging studies may be necessary. On the other hand, children who were septic, dehydrated and with signs of generalized peritonitis on admission, may benefit from short period of IV fluids and antibiotic administration before operation, but this may result in intra-hospital delay of surgery. Yardeni et al. [25] reported that overnight delay of surgery in those patients who presented with non-perforated appendicitis did not significantly affect operating time, rate of perforations or frequency of complications.

Negative appendectomy rate of 6% in our study is low and comparable with other reports [6,26]. Likewise, clinical perforation rate of 26.86% is similar to other reports [6]. It is biased to some extent with the fact that more than half of children with perforation (10 of 18) was referred from other RGH. The main reason for referral was severe clinical presentation. Six referred patients with perforation of the appendix had clinical signs of acute abdomen and x-ray findings of intestinal obstruction. None of the referred patients were treated in RGH before referral. Perforation was found in 3 patients admitted less than 6 hours after onset of the symptoms. Two of them were operated on within 3 hours after admission. These results may support theories [17] that two distinct populations of patients with appendicitis may exist, and that in some patients, inflammation progresses more rapidly to perforation [6]. Furthermore, this implies that perforation correlates more with pre-hospital than with in-hospital delay, and that postponing operation for several hours in order to clarify diagnosis in equivocal cases would not increase the risk of perforation [25,27].

In conclusion, clinical presentation of appendicitis in children may vary, from those with very mild symptoms to those with life threatening septic shock and bowel obstruction. Unfortunately, there is neither a single diagnostic test nor a combination of clinical, laboratory, and imaging studies, that have 100% of sensitivity and/or specificity. Furthermore, there is still no way to distinguish simple acute and perforated appendicitis with certainty before surgery. Delaying surgery for several hours in doubtful cases to clarify diagnosis may further reduce negative appendectomy rates without increasing the risk of perforation.

## References

1. Sudhakaran N, Ade-Ajayi N. Appendicitis in children. *Surgery*. Elsevier Ltd; 2010; 28:16–21.
2. Benabbas R, Hanna M, Shah J, Sinert R. Diagnostic Accuracy of History, Physical Examination, Laboratory Tests, and Point-of-care Ultrasound for Pediatric Acute Appendicitis in the Emergency Department: A Systematic Review and Meta-analysis. *Alpern E, editor. Acad Emerg Med* 2017; 24:523–551.
3. Newman K, Ponsky T, Kittle K, Dyk L, Throop C, Giesecker K, et al. Appendicitis 2000: variability in practice, outcomes, and resource utilization at thirty pediatric hospitals. *J Pediatr Surg* 2003; 38:372-9-9.
4. Alloo J, Gerstle T, Shilyansky J, Ein SH. Appendicitis in children less than 3 years of age: a 28-year review. *Pediatr Surg Int* 2004; 19:777–779.
5. Marjanović Z, Spasić Z, Zivanović D, Kostić A, Djordjević I ZD. Akutni appendicitis kod dece uzrasta do tri godine. *Srp Arh Celok Lek* 2006; 134:203–207.
6. Narsule CK, Kahle EJ, Kim DS, Anderson AC, Luks FI. Effect of delay in presentation on rate of perforation in children with appendicitis. *Am J Emerg Med. Elsevier B.V.*; 2010;2–5.
7. Stevenson MD, Dayan PS, Dudley NC, Bajaj L, Macias CG, Bachur RG, et al. Time From Emergency Department Evaluation to Operation and Appendiceal Perforation. *Pediatrics*. 2017;139.
8. Tsuji M, Puri P, Reen DJ. Characterisation of the local inflammatory response in appendicitis. *J Pediatr Gastroenterol Nutr* 1993; 16:43–48.
9. Wang Y, Reen D, Puri P. Is a histologically normal appendix following emergency appendectomy always normal? *Lancet. Elsevier*. 1996; 347:1076–1079.
10. Nemeth L, Reen DJ, O'Briain DS, McDermott M, Puri P. Evidence of an inflammatory pathologic condition in “normal” appendices following emergency appendectomy. *Arch Pathol Lab Med* 2001; 125:759–764.
11. Wu Hp, Chen C, Kuo I, Wu Y, Fu Y. Diagnostic Values of a Single Serum Biomarker at Different Time Points Compared with Alvarado Score and Imaging Examinations in Pediatric Appendicitis. *J Surg Res. Elsevier Inc*; 2012;174:272–177.
12. Andersson REB. Meta-analysis of the clinical and laboratory diagnosis of appendicitis. *Br J Surg*. 2004;91:28–37.
13. Siddique K, Baruah P, Bhandari S, Mirza S, Harinath G. Diagnostic accuracy of white cell count and C-reactive protein for assessing the severity of paediatric appendicitis. *JRSM Short Rep* 2011; 2:59.
14. Yazici M, Ozkisacik S, Oztan MO, Gürsoy H. Neutrophil/lymphocyte ratio in the diagnosis of childhood appendicitis. *Turk J Pediatr* 2010; 52:400–403.
15. Morrow SE, Newman KD. Current management of appendicitis. *Semin Pediatr Surg* 2007; 16:34–40.
16. Rothrock SG, Pagane J. Acute appendicitis in children: emergency department diagnosis and management. *Ann Emerg Med* 2000; 36:39–51.
17. Livingston EH, Woodward W a, Sarosi G a, Haley RW. Disconnect between incidence of nonperforated and perforated appendicitis: implications for pathophysiology and management. *Ann Surg* 2007; 245:886–892.
18. Kwok MY, Kim MK, Gorelick MH. Evidence-based approach to the diagnosis of appendicitis in children. *Pediatr Emerg Care* 2004; 20:690-8-701.
19. Golledge J, Scriven M. Peritonism in appendicitis. *Ann R Coll Surg Engl* 1996; 78:11–14.
20. O'Shea JS, Bishop ME, Alario AJ, Cooper JM, others. Diagnosing appendicitis in children with acute abdominal pain. *Pediatr Emerg Care* 1988; 4:172.
21. Andersson RE, Hugander a P, Ghazi SH, Ravn H, Offenbartl SK, Nyström PO, et al. Diagnostic value of disease history, clinical presentation, and inflammatory parameters of appendicitis. *World J Surg* 1999; 23:133–140.
22. Roberts JK, Behraves M, Dmitrevski J. Macroscopic findings at appendectomy are unreliable: implications for laparoscopy and malignant conditions of the appendix. *Int J Surg Pathol* 2008; 16:386–390.
23. Bliss D, Mckee J, Cho D, Krishnaswami S, Zallen G, Harrison M, et al. Discordance of the pediatric surgeon's intraoperative assessment of pediatric appendicitis with the pathologists report. *J Pediatr Surg. Elsevier Inc.*; 2010; 45:1398–1403.
24. St Peter SD, Sharp SW, Holcomb GW, Ostlie DJ. An evidence-based definition for perforated appendicitis derived from a prospective randomized trial. *J Pediatr Surg. Elsevier Inc.*; 2008; 43:2242–2245.
25. Yardeni D, Hirschl RB, Drongowski RA, Teitelbaum DH, Geiger JD CA. Delayed versus immediate surgery in acute appendicitis: do we need to operate during the night? *J Pediatr Surg* 2004; 39:464–469.
26. Oyetunji TA, Ong'uti SK, Bolorunduro OB, Cornwell EE 3rd NB. Pediatric negative appendectomy rate: trend, predictors, and differentials. *J Surg Res. Elsevier Inc*; 2012; 173:16–20.
27. Almström M, Svensson JF, Patkova B, Svenningsson A, Wester T. In-hospital Surgical Delay Does Not Increase the Risk for Perforated Appendicitis in Children. *Ann Surg* 2017; 265: 616–621.