

**SUPERMASSIVE BLACK HOLE AT THE GALACTIC CENTER \*****Alexander F. Zakharov**

**Abstract.** We derive an analytical expression of a shadow size as a function of a charge in the Reissner – Nordström (RN) metric. Using the derived expression we consider shadows for negative tidal charges and charges corresponding to naked singularities  $q = Q^2/M^2 > 1$ , where  $Q$  and  $M$  are black hole charge and mass, respectively. An introduction of a negative tidal charge  $q$  can describes black hole solutions in theories with extra dimensions, so following the approach we consider an opportunity to extend RN metric to negative  $Q^2$ , while for the standard RN metric  $Q^2$  is always non-negative. We found that for  $q > 9/8$  black hole shadows disappear. Significant tidal charges  $q = -6.4$  are not consistent with observations of a minimal spot size at the Galactic Center observed in mm-band, moreover, these observations demonstrate that in comparison with the Schwarzschild black hole a Reissner – Nordström black hole with a significant charge  $q \approx 1$  provides a better fit of recent observational data for the black hole at the Galactic Center.

**1. Introduction**

Theories with extra dimensions admit astrophysical objects (supermassive black holes in particular) which are rather different from standard ones. Tests have been proposed when it would be possible to discover signatures of extra dimensions in supermassive black holes since the gravitational field may be different from the standard one in the GR approach. So, gravitational lensing features are different for alternative gravity theories with extra dimensions and general relativity.

Recently, Bin-Nun [1] discussed an opportunity that the black hole at the Galactic Center is described by the tidal Reissner– Nordström metric which may be admitted by the Randall–Sundrum II braneworld scenario [2]. Bin-Nun suggested an opportunity of evaluating the black hole metric analyzing (retro-)lensing of bright stars around the black hole in the Galactic Center. Doeleman et al. evaluated a size of the smallest spot for the black hole at the Galactic Center with VLBI technique

---

Received June 20, 2014.

UDC 524.882

\*The author was supported in part by the NSF (HRD-0833184) and NASA (NNX09AV07A).

in mm-band [3]. Theoretical studies showed that the size of the smallest spot near a black hole practically coincides with shadow size because the spot is the envelope of the shadow [4, 5, 6]. Measurements of the shadow size around the black hole may help to evaluate parameters of black hole metric [5, 6]. We derive an analytic expression for the black hole shadow size as a function of the tidal charge for the Reissner–Nordström metric. We conclude that observational data concerning shadow size measurements are not consistent with significant negative charges, in particular, the significant tidal charge  $q = Q/M^2 = -6.4$  discussed [1], where the author used a little bit different notations, namely  $q' = q/4$ , is practically ruled out with a very high probability (the tidal charge is roughly speaking is far beyond  $9\sigma$  confidence level). We also show a smaller shadow sizes in respect to estimates obtained with the Schwarzschild black hole model can be explained with the Reissner – Nordström metric with a significant charge. It was found a critical  $q$  value for shadow existence, namely for  $q \leq 9/8$ , Reissner – Nordström black holes have shadows while for  $q > 9/8$  the shadows do not exist.

Now there are two basic observational techniques to investigate a gravitational potential at the Galactic Center, namely, a) monitoring the orbits of bright stars near the Galactic Center to reconstruct a gravitational potential [7] (see also a discussion about an opportunity to evaluate black hole dark matter parameters in [8] and an opportunity to constrain some class of alternative theory of gravity [9]); b) in mm-band with VLBI-technique measuring a size and a shape of shadows around black hole giving an alternative possibility to evaluate black hole parameters. The formation of retro-lensing images (also known as mirage, shadows or "faces" in the literature) due to the strong gravitational field effects nearby black holes has been investigated by several authors [5, 6].

As J. A. Wheeler coined "Black holes have no hair": it means that a black hole is characterized by only three parameters ("hairs"), its mass  $M$ , angular momentum  $J$  and charge  $Q$  (see, e.g. [10]). Therefore, in principle, charged black holes can be formed, although astrophysical conditions that lead to their formation may look rather problematic. Nevertheless, one could not claim that their existence is forbidden by theoretical or observational arguments. Moreover, we will show below that observations give a hint about an existence of a significant charge, but its origin is not clear at the moment.

## 2. Basic Equations

The expression for the Reissner - Nordström metric in natural units ( $G = c = 1$ ) has the form

$$(2.1) \quad ds^2 = - \left( 1 - \frac{2M}{r} + \frac{Q^2}{r^2} \right) dt^2 + \left( 1 - \frac{2M}{r} + \frac{Q^2}{r^2} \right)^{-1} dr^2 + r^2 (d\theta^2 + \sin^2\theta d\phi^2).$$

Applying the Hamilton-Jacobi method to the problem of geodesics in the Reissner - Nordström metric, the motion of a test particle in the  $r$ -coordinate can be described

by following equation (see, for example, [10])

$$(2.2) \quad r^4(dr/d\lambda)^2 = R(r),$$

where

$$(2.3) \quad \begin{aligned} R(r) &= P^2(r) - \Delta(\mu^2 r^2 + L^2), \\ P(r) &= Er^2 - eQr, \\ \Delta &= r^2 - 2Mr + Q^2. \end{aligned}$$

Here, the constants  $\mu, E, L$  and  $e$  are associated with the particle, i.e.  $\mu$  is its mass,  $E$  is energy at infinity,  $L$  is its angular momentum at infinity and  $e$  is the particle's charge. We shall consider the motion of uncharged particles ( $e = 0$ ) below. In this case, the expression for the polynomial  $R(r)$  takes the form

$$(2.4) \quad R(r) = (E^2 - \mu^2)r^4 + 2M\mu r^3 - (Q^2\mu^2 + L^2)r^2 + 2ML^2r - Q^2L^2.$$

Depending on the multiplicities of the roots of the polynomial  $R(r)$ , we can have three types of motion in the  $r$  - coordinate [11]. In particular, by defining  $r_+ = 1 + \sqrt{1 - Q^2}$ , we have: (1) if the polynomial  $R(r)$  has no roots for  $r \geq r_+$ , a test particle is captured by the black hole; (2) if  $R(r)$  has roots and  $(\partial R/\partial r)(r_{max}) \neq 0$  with  $r_{max} > r_+$  ( $r_{max}$  is the maximal root), a particle is scattered after approaching the black hole; (3) if  $R(r)$  has a root and  $R(r_{max}) = (\partial R/\partial r)(r_{max}) = 0$ , the particle now takes an infinite proper time to approach the surface  $r = const$ . If we are considering a photon ( $\mu = 0$ ), its motion in the  $r$ -coordinate depends on the root multiplicity of the polynomial  $\hat{R}(\hat{r})$

$$(2.5) \quad \hat{R}(\hat{r}) = R(r)/(M^4E^2) = \hat{r}^4 - \xi^2\hat{r}^2 + 2\xi^2\hat{r} - \hat{Q}^2\xi^2.$$

where  $\hat{r} = r/M, \xi = L/(Me)$  and  $\hat{Q} = Q/M$ . One could see from Eq. (2.5) and Eqs. (2.3) as well that the black hole charge may influence substantially the photon motion at small radii ( $r \approx 1$ ), while the charge effect is almost negligible at large radial coordinates of photon trajectories ( $r \gg 1$ ). In the last case we should keep in mind that the charge may cause only small corrections on photon motion.

### 3. Derivation of shadow size as a function of charge

Let us consider the problem of the capture cross section of a photon by a charged black hole. It is clear that the critical value of the impact parameter for a photon to be captured by a Reissner - Nordström black hole depends on the multiplicity root condition of the polynomial  $R(r)$ . This requirement is equivalent to the vanishing discriminant condition [12]. To find the critical value of impact parameter for Schwarzschild and RN metrics the condition has been used for corresponding cubic and quartic equations [13, 14, 15]. In particular, it was shown that for these cases

the vanishing discriminant condition approach is more powerful in comparison with the procedure excluding  $r_{max}$  from the following system

$$(3.1) \quad R(r_{max}) = 0, \quad \frac{\partial R}{\partial r}(r_{max}) = 0,$$

as it was done, for example, by Chandrasekhar [16] to solve similar problems, because  $r_{max}$  is automatically excluded in the condition for vanishing discriminant. Introducing the notation  $\xi^2 = l$ ,  $Q^2 = q$ , we obtain

$$(3.2) \quad R(r) = r^4 - lr^2 + 2lr - ql.$$

We remind basic algebraic definitions and relations. If we consider an arbitrary polynomial  $f(X)$  with degree  $n$

$$(3.3) \quad f(X) = X^n + a_1X^{n-1} + \dots + a_{n-1}X + a_n,$$

the elementary symmetric polynomials  $s_k$  have the following form, where  $X_1, \dots, X_n$  are roots of the polynomial  $f(X)$  [12]

$$(3.4) \quad s_k(X_1, \dots, X_n) = \sum_{1 \leq i_1 < i_2 < \dots < i_k \leq n} X_{i_1} X_{i_2} \dots X_{i_k},$$

where  $k = 1, 2, \dots, n$ . The symmetrical  $k$ -power sum polynomial  $p_k$  have the following expression [12]

$$(3.5) \quad p_k(X_1, \dots, X_n) = X_1^k + X_2^k + \dots + X_n^k, \quad \text{for } k \geq 0.$$

To express  $p_k$  through  $s_k$  one can use Newton's equations [12]

$$p_k - p_{k-1}s_1 + p_{k-2}s_2 + \dots + (-1)^{k-1}p_1s_{k-1} + (-1)^kks_k = 0, \quad \text{for } 1 \leq k \leq n;$$

$$p_k - p_{k-1}s_1 + p_{k-2}s_2 + \dots + (-1)^{n-1}p_{k-n+1}s_{n-1} + (-1)^np_{k-n}s_n = 0, \quad \text{for } k > n.$$

We introduce the following polynomial

$$(3.6) \quad \Delta_n(X_1, \dots, X_n) = \prod_{1 \leq i < j \leq n} (X_i - X_j),$$

which can be represented as the Vandermonde determinant

$$(3.7) \quad \Delta_n(X_1, \dots, X_n) = \begin{vmatrix} 1 & 1 & \dots & 1 \\ X_1 & X_2 & \dots & X_n \\ \dots & \dots & \dots & \dots \\ X_1^{n-1} & X_2^{n-1} & \dots & X_n^{n-1} \end{vmatrix}.$$

According to the discriminant  $Dis$  definition we have the  $Dis(s_1, \dots, s_n)$  polynomial

$$(3.8) \quad Dis(s_1, \dots, s_n) = \Delta_n^2(X_1, \dots, X_n) = \prod_{1 \leq i < j \leq n} (X_i - X_j)^2,$$

one can find [12]

$$(3.9) \quad Dis(s_1, \dots, s_n) = \begin{vmatrix} n & p_1 & p_2 & \dots & p_{n-1} \\ p_1 & p_2 & p_3 & \dots & p_n \\ p_2 & p_3 & p_4 & \dots & p_{n+1} \\ \dots & \dots & \dots & \dots & \dots \\ p_{n-1} & p_n & p_{n+1} & \dots & p_{2n-2} \end{vmatrix}.$$

Clearly, that the vanishing discriminant condition is equivalent to an existence of multiple roots among roots  $X_1, \dots, X_n$ . We apply this technique for the quartic polynomial  $R(r)$  in Eq. (3.2). So, the symmetric k-power polynomials for  $n = 4$  have the form

$$(3.10) \quad p_k = X_1^k + X_2^k + X_3^k + X_4^k, k \geq 0.$$

The symmetric elementary polynomials for  $n = 4$  have the form

$$(3.11) \quad \begin{aligned} s_1 &= X_1 + X_2 + X_3 + X_4, \\ s_2 &= X_1X_2 + X_1X_3 + X_1X_4 + X_2X_3 + X_2X_4 + X_3X_4, \\ s_3 &= X_1X_2X_3 + X_2X_3X_4 + X_2X_3X_4, \\ s_4 &= X_1X_2X_3X_4. \end{aligned}$$

We calculate the discriminant of the family  $X_1, X_2, X_3, X_4$

$$Dis(s_1, s_2, s_3, s_4) = \begin{vmatrix} 1 & 1 & 1 & 1 \\ X_1 & X_2 & X_3 & X_4 \\ X_1^2 & X_2^2 & X_3^2 & X_4^2 \\ X_1^3 & X_2^3 & X_3^3 & X_4^3 \end{vmatrix}^2 = \begin{vmatrix} 4 & p_1 & p_2 & p_3 \\ p_1 & p_2 & p_3 & p_4 \\ p_2 & p_3 & p_4 & p_5 \\ p_3 & p_4 & p_5 & p_6 \end{vmatrix}.$$

Expressing the polynomials  $p_k (1 \leq k \leq 6)$  in terms of the polynomials  $s_k (1 \leq k \leq 4)$  and using Newton's equations we calculate the polynomials and discriminant of the family  $X_1, X_2, X_3, X_4$  in roots of the polynomial  $R(r)$ ; we obtain

$$(3.12) \quad \begin{aligned} p_1 &= s_1 = 0, & p_2 &= -2s_2, & p_3 &= 3s_3, & p_4 &= 2s_2^2 - 4s_4, \\ p_5 &= -5s_3s_2, & p_6 &= -2s_2^3 + 3s_2^2 + 6s_4s_2, \end{aligned}$$

where  $s_1 = 0, s_2 = -l, s_3 = -2l, s_4 = -ql$ , corresponding to the polynomial  $R(r)$  in Eq. (3.2). The discriminant  $Dis$  of the polynomial  $R(r)$  has the form:

$$(3.13) \quad \begin{aligned} Dis(s_1, s_2, s_3, s_4) &= \begin{vmatrix} 4 & 0 & 2l & -6l \\ 0 & 2l & -6l & 2l(l+2q) \\ 2l & -6l & 2l(l+2q) & -10l^2 \\ -6l & 2l(l+2q) & -10l^2 & 2l^2(l+6+3q) \end{vmatrix} = \\ &= 16l^3[l^2(1-q) + l(-8q^2 + 36q - 27) - 16q^3]. \end{aligned}$$

The polynomial  $R(r)$  thus has a multiple root if and only if

$$(3.14) \quad l^3[l^2(1-q) + l(-8q^2 + 36q - 27) - 16q^3] = 0.$$

Excluding the case  $l = 0$ , which corresponds to a multiple root at  $r = 0$ , we find that the polynomial  $R(r)$  has a multiple root for  $r \geq r_+$  if and only if

$$(3.15) \quad l^2(1-q) + l(-8q^2 + 36q - 27) - 16q^3 = 0.$$

If  $q = 0$ , we obtain the well-known result for a Schwarzschild black hole [10],  $l = 27$ , or  $L_{cr} = 3\sqrt{3}$ . If  $q = 1$ , then  $l = 16$ , or  $L_{cr} = 4$ . The photon capture cross section for an extreme charged black hole turns out to be considerably smaller than the capture cross section of a Schwarzschild black hole. The critical value of the impact parameter, characterizing the capture cross section for a RN black hole, is determined by the equation

$$(3.16) \quad l_{cr} = \frac{(8q^2 - 36q + 27) + \sqrt{D_1}}{2(1-q)},$$

where  $D_1 = (8q^2 - 36q + 27)^2 + 64q^3(1-q) = -512 \left(q - \frac{9}{8}\right)^3$ . It is clear from the last relation that there are circular unstable photon orbits only for  $q \leq \frac{9}{8}$ . Substituting Eq.(3.16) into the expression for the coefficients of the polynomial  $R(r)$  it is easy to calculate the radius of the unstable circular photon orbit (which is the same as the minimum periastron distance). The orbit of a photon moving from infinity with the critical impact parameter, determined in accordance with Eq.(3.16) spirals into circular orbit. To find a radius of photon unstable orbit we will solve Eq. (3.1) substituting  $l_{cr}$  in the relation. From trigonometric formula for roots of cubic equation we have

$$(3.17) \quad r_{crit} = 2\sqrt{\frac{l_{cr}}{6}} \cos \frac{\alpha}{3},$$

where  $\cos \alpha = -\sqrt{\frac{27}{2l_{cr}}}$ . As it was explained [6] this leads to the formation of shadows described by the critical value of  $L_{cr}$  or, in other words, in the spherically symmetric case, shadows are circles with radii  $L_{cr}$ . Therefore, measuring the shadow size, one could evaluate the black hole charge in black hole mass units  $M$ . In Fig. 3.1 shadow size and radius of last unstable orbit for photons are given as a function of charge (including possible tidal charge with a negative  $q$  and super-extreme charge  $q > 1$ ).




FIG. 3.1: Shadow (mirage) radius (solid line) and radius of the last circular unstable photon orbit (dot-dashed line) in  $M$  units as a function of  $q$ . The critical value  $q = 9/8$  is shown with dashed vertical line.

#### 4. Consequences

In spite of the cosmic censorship hypothesis that a singularity has to be shielded by a horizon, properties of naked singularities are a subject of intensive theoretical studies. As usual spherical symmetrical cases are more easier for analysis and RN metrics with super extreme charge  $q > 1$  are investigated in a number of papers.

So, if we assume that  $q > 1$ , we can see from Eq. (3.16) that for  $q \leq 9/8$  we have shadows, while for  $q > 9/8$  the shadows do not exist. For these charges ( $q > 9/8$ ) incoming photons always scattering by black holes for  $l \neq 0$  because the polynomial  $R(r)$  has no multiple roots but it has a single positive root (it means scattering) since for great positive  $r$  we have  $R(r) > 0$  while  $R(0) < 0$ . The degenerate case of radial trajectories of photons ( $l = 0$ ) can be ignored as the case with "zero measure" or the structural unstable case using the Poincaré – Pontryagin – Andronov – Anosov – Arnold terminology. It means that in any small vicinity a behavior of other geodesics from the radial ones is qualitatively different, therefore, such objects can not be observed in nature. Therefore, shadows exist only for  $q \leq 9/8$ . So,  $q = 9/8$  is critical value which is characterized "catastrophe" or the qualitatively different behavior of the system (appearance and disappearance of shadows). For the critical  $q = 9/8$  we have the smallest shadow with  $l = 27/2$  and a shadow size  $\xi = \sqrt{13.5} \approx 3.674$  (in  $M$  units) or  $38.4 \mu\text{as}$  in diameter for the

black hole at the Galactic Center. For this impact parameter we have corresponding circular unstable orbit for photons with  $r = 3/2$  (in  $M$  units).

If we adopt the distance toward the Galactic Center  $d_* = 8.3$  kpc and mass of the black hole  $M_{BH} = 4.4 * 10^6 M_\odot$  [17], then we have the angle  $10.45 \mu as$  for the corresponding Schwarzschild radius  $R_g = 2.95 * \frac{M_{BH}}{M_\odot} * 10^5$  cm, so a shadow size for the Schwarzschild black hole is around  $54.2 \mu as$ , for a black hole with a tidal charge ( $q = -6.4$ ) suggested by Bin-Nun [1] a shadow size is about  $88.1 \mu as$ , while for the extreme charge ( $q = 1$ ) and critical charge ( $q = 9/8$ ) the shadow sizes are  $41.8 \mu as$  and  $38.4 \mu as$ , respectively. A couple of years ago Doeleman et al. [3] claimed that intrinsic diameter of Sgr  $A^*$  is  $37_{-10}^{+16} \mu as$  at the  $3 \sigma$  confidence level. If we believe in GR and the central object is a black hole, then we have to conclude that a shadow is practically coincides with the intrinsic diameter, so in spite of the fact that a Schwarzschild black hole is marginally consistent with observations, a Reissner – Nordström black hole provides much better fit of a shadow size, while a black hole with a significant tidal charge ( $q = -6.4$ ) is out of  $9.6 \sigma$  level interval. Later on, an accuracy of intrinsic size measurements was significantly improved, so Fish et al. [18] gave  $41.3_{-4.3}^{+5.4} \mu as$  (at  $3 \sigma$  level) on day 95,  $44.4_{-3.0}^{+3.0} \mu as$  on day 96 and  $42.6_{-2.9}^{+3.1} \mu as$  on day 97, so a tidal charge ( $q = -6.4$ ) is out of  $26 \sigma$  level for day 95 and even less probable for other observations.

The black hole in the elliptical galaxy M87 looks also perspective to evaluate shadow size [19] (probably even its shape in the future to estimate a black hole spin) because the distance toward the galaxy is  $16 \pm 0.6$  Mpc, black hole mass is  $M_{M87} = (6.2 \pm 0.4) * 10^9 M_\odot$ , so that an angle  $(7.3 \pm 0.5) \mu as$  corresponds to the Schwarzschild radius [19], so the angle is comparable with the corresponding value considered earlier for our Galactic Center case. In paper [19] it was reported that smallest shadow size is  $5.5 \pm 0.4 R_{SCH}$  with  $1 \sigma$  errors (where  $R_{SCH} = 2GM_{M87}/c^2$ ), so that at the moment the shadow size is consistent with the Schwarzschild metric for the object.

## 5. Conclusions

Based on observations [3, 18] one can say that between for the Schwarzschild black hole model we have tensions between evaluations of black hole mass done with observations of bright star orbits near the Galactic Center and the evaluated shadow size. To reduce tensions between estimates of the black hole mass and the intrinsic size measurements, one can use the Reissner – Nordström metric with a significant charge which is comparable with the critical one. We do not claim that the corresponding charge has an electric origin because an interstellar environment is electrically neutral, so the corresponding charge may be induced (like a tidal charge induced by extra dimension) and has a non-electric origin. Recent estimates of the smallest structure in the M87 published in paper [19] do not need an introduction of charge (tidal or normal) to fit observational data because sizes of the smallest

spot near the black hole at the object are consistent with the shadow size evaluated for the Schwarzschild metric.

The author thanks D. Borka, V. Borka Jovanović, F. De Paolis, G. Djordjević, G. Ingrosso, P. Jovanović, A. A. Nucita, B. Vlahovic for useful discussions and organizers of the 6th Balkan Workshop (BW2013) for the kind attention to the contribution.

## REFERENCES

1. A. Y. BIN-NUN: *Gravitational lensing of stars orbiting Sgr A\* as a probe of the black hole metric in the Galactic center*. Phys. Rev. D **82** (2010), 064009-8.
2. N. DADHICH, R. MAARTENS, PH. PAPAPOPOULOS, V. REZANIA: *Black holes on the brane*. Phys. Lett. B **487** (2000), 1–6.
3. S.S. DOELEMAN et al.: *Event-horizon-scale structure in the supermassive black hole candidate at the Galactic Centre*. Nature, **455** (2008), 78–80.
4. H. FALCKE, F. MELIA and E. AGOL: *Viewing the Shadow of the Black Hole at the Galactic Center*. ApJ. **528** (2000), L13-L16; F. MELIA and H. FALCKE: *The Supermassive Black Hole at the Galactic Center*. Annual Rev. Astron. & Astrophys. **39** (2001), 309–352.
5. A.F. ZAKHAROV, A.A. NUCITA, F. DE PAOLIS and G. INGROSSO: *Measuring the black hole parameters in the galactic center with RADIOASTRON*. New Astronomy. **10** (2005), 479 – 489.
6. A.F. ZAKHAROV, A.A. NUCITA, F. DE PAOLIS and G. INGROSSO: *Direct measurements of black hole charge with future astrometrical missions*. Astron. & Astrophys. **10** (2005), 795–799; A.F. ZAKHAROV, A.A. NUCITA, F. DE PAOLIS and G. INGROSSO: *Shadows as a tool to evaluate black hole parameters and a dimension of spacetime*. New Astron. Rev. **56** (2012), 64–73; A.F. ZAKHAROV: *Constraints on a charge in the Reissner-Nordström metric for the black hole at the Galactic Center*. Phys. Rev. D **90** (2007), 062007-8.
7. A. M. GHEZ, M. MORRIS, E. E. BECKLIN, A. TANNER and T. KREMENEK: *The accelerations of stars orbiting the Milky Way's central black hole*. Nature. **407** (2000), 349–351.
8. A. F. ZAKHAROV, A. A. NUCITA, F. DE PAOLIS AND G. INGROSSO: *Apoastron shift constraints on dark matter distribution at the Galactic Center*. Phys. Rev. D **76** (2007), 062001-6; A.F. ZAKHAROV, S. CAPOZZIELLO, F. DE PAOLIS, G. INGROSSO and A.A. NUCITA: *The Role of Dark Matter and Dark Energy in Cosmological Models: Theoretical Overview*. Space Sci. Rev. **48** (2009), 301–313.
9. D. BORKA, P. JOVANOVIĆ, V. BORKA JOVANOVIĆ and A. F. ZAKHAROV: *Constraints on  $R^n$  gravity from precession of orbits of S2-like stars*. Phys. Rev. D **85** (2012), 124004-11.
10. C. MISNER, K. THORNE and J.A. WHEELER: *Gravitation*, W.H. Freeman and Company, San Francisco, 1973.
11. A.F. ZAKHAROV: *Types of unbound geodesics in the Kerr metric*. Sov. Phys. JETP. **64** (1986), 1–3.

12. A.I. KOSTRIKIN: *Introduction to Algebra*. Springer, 1982.
13. A.F. ZAKHAROV: *Effective Particle Cross Section of a Schwarzschild Black Hole*. *Sov. Astron.* **32** (1988), 456–457.
14. A.F. ZAKHAROV: *Capture Cross Sections for a Charged Reissner – Nordström Black Hole: Photons, Ultrarelativistic Unchanged Particles and Slow Uncharged Particles*. *Sov. Astron.* **35** (1991), 147–149.
15. A.F. ZAKHAROV: *Particle Capture Cross Section for a Reissner – Nordström Black Hole*. *Class. Quant. Grav.* **11**, 1027 (1994), 1027 – 1033.
16. S. CHANDRASEKHAR: *Mathematical Theory of Black Holes*. Clarendon Press, Oxford, 1983.
17. S. GILLESSEN et al.: *The Orbit of the Star S2 around Sgr A\* from Very Large Telescope and Keck Data*: *ApJ* **707** (2009), L114–L117.
18. V. L. FISH et al.: *1.3 mm Wavelength VLBI of Sagittarius A\*: Detection of Time-Variable Emission on Event Horizon Scales*. *ApJ* **727** (2011), L36 (6pp.).
19. S. S. DOELEMAN et al.: *Jet-Launching Structure Resolved Near the Supermassive Black Hole in M87*. *Science* **338** (2012), 355–358.

Alexander F. Zakharov  
Institute of Theoretical and Experimental Physics  
B. Cheremushkinskaya, 25  
117218 Moscow, Russia  
[zakharov@itep.ru](mailto:zakharov@itep.ru)

Joint Institute for Nuclear Research  
Bogoliubov Laboratory of Theoretical Physics  
Joliot-Curie 6, 141980 Dubna  
Moscow region, Russia

North Carolina Central University  
1801 Fayetteville Rd,  
Durham, NC 27707, USA

National Research Nuclear University MEPhI  
(Moscow Engineering Physics Institute),  
115409, Moscow, Russia