

**Original research article**

**HAEMATOLOGICAL AND IRON STATUS  
FOLLOWING A SOCCER MATCH**

*UDC 616.15:796.332*

**Goran Sporiš<sup>1</sup>, Tomislav Vlahović<sup>1</sup>, Nebojša Trajković<sup>2</sup>,  
Zoran Milanović<sup>2</sup>, Dejan Madić<sup>3</sup>**

<sup>1</sup>Faculty of Kinesiology, University of Zagreb, Zagreb, Croatia

<sup>2</sup>Faculty of Sport and Physical Education, University of Niš, Serbia

<sup>3</sup>Faculty of Sport and Physical Education, University of Novi Sad, Serbia

**Abstract.** *The aim of the present investigation was to verify the responses of haematological and iron concentrations in Croatian U-21 soccer players following an official soccer match. A group of 19 trained, healthy male soccer players from the U-21 category of the Croatian soccer association took part in this study. An analysis of red blood cell concentration (RBC), haemoglobin (Hb), haematocrit (Hct), mean corpuscular volume (MCV), mean cell haemoglobin (MCH), mean cell haemoglobin corpuscular (MCHC), and Thrombocytes was performed to indicate the haematological status. In addition, iron, total iron binding capacity (TIBC), the unsaturated iron-binding capacity (UIBC) and ferritin analyses were obtained immediately after the soccer match. A significant decrease in Hct (%) was observed immediately after the game. However, there were no significant changes in the haemoglobin levels. Iron concentration increased significantly ( $p < 0.01$ ) after the soccer match. Ferritin levels were also higher at the end of the match compared to baseline measures. It can be reported that a soccer match may induce an iron increase immediately after the game which was contradictory to recent studies. In summary, this study provided a valuable opportunity to study biochemical parameters that could lead to a possible illness, injuries and severity in match-play.*

**Key words:** *biochemical analysis, soccer, professional players, fatigue.*

---

Received April 13, 2016 / Accepted July 13, 2016

Corresponding author: Nebojša Trajković

University of Niš, Faculty of Sport and Physical Education, Čarojevića 10a, Niš, Serbia

Phone: +381 18 510 900 • E-mail: nele\_trajce@yahoo.com

## INTRODUCTION

Soccer is played by over 250 million people in more than 200 countries (Dunning, 1999). Therefore, its popularity leads to a constant effort for improving performance. Male soccer players draw all the attention and the biochemical impact of a soccer match on male players has extensively been researched (Ascensão et al., 2008). However, it is essential to separate age groups based on the potential for differences.

When focusing on physiological demands, several different blood markers within the athletes can be very helpful to not only get an overall better understanding of the sports' demands, but also to identify certain norms and tendencies present in these particular athletes, who unfortunately are often subject to iron and haemoglobin deficiencies (Landahl, Adolfsson, Borjesson, Mannheimer, & Rodjer, 2005). Studies have shown that results of iron status in sport are rather contradictory. Ostojic and Ahmetovic (2009) stated that iron status indicators were stable and poorly related to the training phase during the sports season. Tan, Dawson, and Peeling (2012) found that iron stores and serum levels were significantly decreased after training sessions. In contrast, some studies have indicated that exercise does not affect the haematology variables (Akgün, 1994; Spiropoulos & Trakada, 2003). Changes in haematological characteristics have been linked to aerobic exercise, such as a decrease in haematocrit and increase in haemoglobin values in young and old individuals (El-Sayed, Sale, Jones, & Chester, 2000). Increase in the leukocytes count was reported in young people following acute resistance exercise (Ramel, Wagner, & Elmadfa, 2003).

Recent studies found reduction in physical performance (Ascensão et al., 2008; Ispirlidis et al., 2008; Magalhães et al., 2010; Rampinini, Coutts, Castagna, Sassi, & Impellizzeri, 2007), increase of muscular markers damage (Andersson, Ekblom, & Krstrup, 2008; Ascensão et al., 2008; Ispirlidis et al., 2008; Magalhães et al., 2010), robust pro- and anti-inflammatory cytokine response (Andersson, Karlsen, Blomhoff, Raastad, & Kadi, 2010; Ispirlidis et al., 2008), increased catabolic (Ispirlidis et al., 2008; Kraemer et al., 2004) and pro oxidant state (Ascensão et al., 2008; Magalhães et al., 2010) are described just after a "friendly" soccer match and during recovery after the match. However, there is evidence that physical intensity during a friendly soccer match (absolute values and percentage of maximum heart rate) can be lower than during official matches (Rodrigues et al., 2007) which is probably due to motivational matters. Following a soccer match there is considerable muscle damage and oxidative stress as well as a marked rise in antioxidant status markers for up to 48 h to 72 h post exercise (Andersson et al., 2010; Ascensão et al., 2008; Ispirlidis et al., 2008). Haematological parameters are influenced by several factors, of which training, age, sex, ethnicity, nutrition, and altitude (Ostojic & Ahmetovic, 2009; Schumacher, Jankovits, Bültmann, Schmid, & Berg, 2002) are the most common. Most of these factors can have a positive or negative influence on haematocrit (Hct), haemoglobin (Hb), and red blood cell (RBC) count.

Soccer is an endurance-based sport during which athletes require large amounts of oxygen within the different systems of the body in order for them to function. For this, high iron and haemoglobin values would be very beneficial for these athletes and their performance. Consequently, regular monitoring of selected psychological, biochemical and physiological markers can prevent or avoid overreaching and overtraining throughout a training program (Silva, Santhiago, Papoti, & Gobatto, 2008). Therefore, the main aim of the present investigation was to verify the responses of haematological and iron concentrations in Croatian U-21 soccer players following an official soccer match.

## METHODS

### Participants

A group of 19 trained, healthy male soccer players from the U-21 category of the Croatian soccer association took part in this study. The players were informed about the experimental procedures and possible discomforts associated with the study, and a written informed consent was obtained. The study was approved by the Ethics Committee of the Faculty of Kinesiology, University of Zagreb in accordance with the Helsinki Declaration. The participants were aware that they could withdraw from the study at any time. The participants' characteristics were on average: age  $20.26 \pm 0.65$  years, body mass:  $71.3 \pm 5.9$ ; body height:  $1.77 \pm 0.07$  and maximal oxygen uptake (VO<sub>2</sub>max)  $64.95 \pm 3.99$  ml·kg<sup>-1</sup>·min<sup>-1</sup>. The selection criteria included: (1) participation at professional (top three division leagues) level of soccer competition for at least 5 years, (2) all players participated in at least 75% training sessions per week and played in at least 16 matches during the season, (3) no consumption of exogenous anabolic-androgenic steroids or other drugs that might have affected their physical performance or hormonal balance during the study (for at least 6 months) (5) no recent history of febrile illness, muscle lesions, lower limb trauma, and metabolic diseases.

The soccer players were instructed not to change their normal eating habits during the entire period of data collection. Nutritional supplements were not included in their diets. In addition, players were instructed to refrain from drinking beverages containing caffeine or alcohol and from consuming food 3 h before testing.

### Blood collection and analysis

Blood samples were assessed pre-match and immediately after a match in response to a competitive (2×45 min) soccer match. On the day of the game, the players arrived at the laboratory after an overnight fast of between 10 and 12 h. A resting blood sample was taken after the participants had been standing for at least 15 min, after which the participants consumed a light standardized meal and drink and rested for 2 h. The meal consisted of 1.7 g white bread and 0.3 g of a low-fat spread; both values are per kilogram of body mass (Thompson et al., 2003). The participants abstained from alcohol and caffeine consumption for at least 24 h, and did not perform any exercise for the last 72 h before testing.

Whole blood was collected in EDTA tubes then divided for biochemical, antioxidant, and oxidative analyses. EDTA blood samples were sent for immediate analysis of red blood cell concentration (RBC), haemoglobin (Hb), haematocrit (Htc), mean corpuscular volume (MCV), mean corpuscular haemoglobin (MCH), mean corpuscular haemoglobin concentration (MCHC), and Thrombocytes on automatic analyser (Dimension Xpand Plus Analyzer, Siemens, Munich, Germany). The remaining freshly withdrawn blood was immediately centrifuged at 3000 rpm during 10 min for careful removal of the plasma. The plasma was separated into several aliquots and rapidly frozen at  $-20$  °C for later biochemical analysis. Iron, total iron binding capacity (TIBC), the unsaturated iron-binding capacity (UIBC) and ferritin analyses were automatically performed in Dimension analyser (Dimension Xpand Plus Analyzer, Siemens, Munich, Germany).

### Statistical analysis

All statistical analyses were performed using the Statistica v.8.0 software (StatSoft Inc., Tulsa, OK, USA). Descriptive statistics and Kolmogorov–Smirnov tests (normality of the distribution) were calculated for all experimental data before inferential testing. Data were expressed as mean values  $\pm$  standard deviation. Sample t-test was used where significant variations occurred. The level of significance was set at  $p < 0.05$ .

## RESULTS

**Table 1** Hematological parameters before and after a soccer match. The values are shown as the mean and Standard Deviation.

	Prematch	Postmatch	p
RBC (mill/mm <sup>3</sup> )	5.07 $\pm$ 0.27	5.04 $\pm$ 0.28	0.46
Hemoglobin (g/L)	150.42 $\pm$ 9.06	149.95 $\pm$ 9.31	0.54
Thrombocytes ( $\times 10^9$ /L)	207.79 $\pm$ 33.14	242.89 $\pm$ 41.32	0.89
HCT (%)	0.45 $\pm$ 0.03	0.43 $\pm$ 0.02	<b>0.00</b>
MCV (fL)	89.68 $\pm$ 2.55	84.03 $\pm$ 2.29	0.25
MCH (pg)	29.70 $\pm$ 0.87	29.72 $\pm$ 0.88	0.81
MCHC (g/l)	331.16 $\pm$ 6.75	334.89 $\pm$ 6.51	<b>0.00</b>

(RBC) red blood cell, (HCT) haematocrit, (MCV) mean corpuscular volume, (MCH) mean corpuscular haemoglobin, (MCHC) mean corpuscular haemoglobin concentration; significant difference at  $p < 0.05$ .

A significant decrease in Hct (%) was observed immediately after the game ( $p < 0.05$ ). As shown in Table 1, only MCHC increased significantly ( $p < 0.05$ ) at the end of the soccer match. There were no significant changes in the haemoglobin, red blood cells, thrombocytes, mean corpuscular volume and mean corpuscular haemoglobin following a soccer match.

**Table 2** Iron status and ferritin before and after soccer match. The values are shown as the mean and standard deviation.

	Prematch	Postmatch	p
Iron ( $\mu\text{mol/L}$ )	19.42 $\pm$ 7.14	25.30 $\pm$ 6.66	<b>0.00</b>
UIBC ( $\mu\text{mol/L}$ )	39.96 $\pm$ 7.65	38.50 $\pm$ 7.70	0.13
TIBC ( $\mu\text{mol/l}$ )	57.8 $\pm$ 8.3	55.94 $\pm$ 6.8	0.24
Ferritin ( $\mu\text{g/L}$ )	40.63 $\pm$ 18.74	49.01 $\pm$ 24.34	<b>0.00</b>

(TIBC) total iron binding capacity, (UIBC) the unsaturated iron-binding capacity; significant difference at  $p < 0.05$ .

Iron concentration ( $\mu\text{mol/L}$ , Table 2) increased significantly ( $p < 0.01$ ) after a soccer match. Ferritin levels were also higher at the end of the match compared to baseline measures (Table 2). However, TIBC ( $p = 0.24$ ) and UIBC ( $p = 0.13$ ) remained unchanged immediately after a soccer match.

## DISCUSSION

Given the fact that soccer match effort implies several acute physiological changes, we tested the hypothesis that an official soccer match induces changes in haematological and iron concentrations in Croatian U-21 soccer players. We found that serum iron concentration demonstrates a significant increase in response to a soccer game. In addition, a significant decrease in Haematocrit values was observed immediately after the soccer match. UIBC and TIBC showed similar values pre to post match testing ( $p > 0.05$ ).

Several studies have reported that the decrease in haemoglobin and haematocrit is a sign of physical exertion and heavy participation (Fallon, 2008; Malcovati, Pascutto, & Cazzola, 2003; Rietjens, Kuipers, Hartgens, & Keizer, 2002). Also, Al'Hazzaa, Almuzaini, Al-Refae, and Sulaiman (2001) stated that decrease in haemoglobin and haematocrit is evident in both endurance oriented sports and in sports that require both aerobic and anaerobic contributions to the energy supply for muscle contraction. We observed that haemoglobin, haematocrit and MCHC diminished in the soccer players after a soccer match. However, only haematocrit and MCHC values showed significant decrease. Despite the decrease in Hb and Hct, we did not find any decrease in the red blood cell count. Accordingly, other studies showed a decrease in Hb and Hct and no changes in RBC (Ostojic & Ahmetovic, 2009; Robach et al., 2014).

Iron, TIBC, and ferritin levels are traditional biomarkers for screening athletes during the training season (Fallon, 2008). We have found a significant increase in Iron and Ferritin values following a 90 min soccer match. Similarly, it was previously reported that eccentric type of exercise induces higher levels of Iron in post compared to pre training measurement (Childs, Jacobs, Kaminski, Halliwell, & Leeuwenburgh, 2001). Iron deficiency may have a negative impact on oxygen transport and immune defense, thus influencing athletic performance (Peeling, Dawson, Goodman, Landers, & Trinder, 2008). Although several studies (Mercer & Densmore, 2005; Ostojic & Ahmetovic, 2009; Resina et al., 1991) have reported a range of 11 to 15% of the players suffering from anemia over a season, none of the players in the present study was iron deficient ( $>13.0 \text{ g}\cdot\text{dl}^{-1}$ ). This finding suggests that the players were subjected to an appropriate level of training intensity and duration. However, a recent study showed that prolonged intense running may reduce iron levels and increase Ferritin values, as a response to a rise in inflammatory and muscle damage markers (Kłapcińska et al., 2013). Discrepancies in the results may be attributed to the fact that football match-play may represent a lower inflammatory stimulus than long-distance running. Possible increased levels of serum ferritin might be an adaptive response to increased oxidative stress mediated by iron, because ferritin sequesters iron in blood or in cells and attenuates its pro-oxidant activity (Meneghini, 1997; Orino et al., 2001).

## CONCLUSION

We can report that a soccer match may induce an iron increase immediately after the game. Ferritin and Haematocrit are also affected by a soccer game. During the competitive season, soccer players are exposed to a variety of physical and psychological stresses from practice, conditioning, and competition. It is essential for players to let their bodies recover from previous practice sessions and games. In summary, this study provided a valuable opportunity to study biochemical parameters that could lead to possible illness, injury and

severity in match-play. Accordingly, it is also essential to educate the athletes about the importance of iron for their performance and to make sure they incorporate iron rich foods in their diet to help prevent too low iron stores and haemoglobin presence.

#### REFERENCES

- Akgün, N. (1994). *Exercise and sports physiology*. Printing. University of the Aegean. Printing Office, Bornova.
- Al'Hazzaa, H., Almuzaini, K., Al-Refae, S., & Sulaiman, M. (2001). Aerobic and anaerobic power characteristics of Saudi elite soccer players. *Journal of Sports Medicine and Physical Fitness*, 41(1), 54.
- Andersson, H., Ekblom, B., & Krstrup, P. (2008). Elite football on artificial turf versus natural grass: Movement patterns, technical standards, and player impressions. *Journal of Sports Sciences*, 26(2), 113-122.
- Andersson, H., Karlsen, A., Blomhoff, R., Raastad, T., & Kadi, F. (2010). Plasma antioxidant responses and oxidative stress following a soccer game in elite female players. *Scandinavian Journal of Medicine and Science in Sports*, 20(4), 600-608.
- Ascensão, A., Rebelo, A., Oliveira, E., Marques, F., Pereira, L., & Magalhães, J. (2008). Biochemical impact of a soccer match-analysis of oxidative stress and muscle damage markers throughout recovery. *Clinical Biochemistry*, 41(10), 841-851.
- Childs, A., Jacobs, C., Kaminski, T., Halliwell, B., & Leeuwenburgh, C. (2001). Supplementation with vitamin C and N-acetyl-cysteine increases oxidative stress in humans after an acute muscle injury induced by eccentric exercise. *Free Radical Biology and Medicine*, 31(6), 745-753.
- Dunning, E. (1999). *Sport matters: Sociological studies of sport, violence, and civilization*. Psychology Press.
- El-Sayed, M. S., Sale, C., Jones, P., & Chester, M. (2000). Blood hemostasis in exercise and training. *Medicine and Science in Sports and Exercise*, 32(5), 918-925.
- Fallon, K. E. (2008). The clinical utility of screening of biochemical parameters in elite athletes: Analysis of 100 cases. *British Journal of Sports Medicine*, 42(5), 334-337.
- Ispirliidis, I., Fatouros, I. G., Jamurtas, A. Z., Nikolaidis, M. G., Michailidis, I., Douroudos, I., et al. (2008). Time-course of changes in inflammatory and performance responses following a soccer game. *Clinical Journal of Sport Medicine*, 18(5), 423-431.
- Klapcińska, B., Waśkiewicz, Z., Chrapusta, S. J., Sadowska-Krepa, E., Czuba, M., & Langfort, J. (2013). Metabolic responses to a 48-h ultra-marathon run in middle-aged male amateur runners. *European Journal of Applied Physiology*, 113(11), 2781-2793.
- Kraemer, W. J., French, D. N., Paxton, N. J., Häkkinen, K., Volek, J. S., Sebastianelli, W. J., et al. (2004). Changes in exercise performance and hormonal concentrations over a big ten soccer season in starters and nonstarters. *The Journal of Strength & Conditioning Research*, 18(1), 121-128.
- Landahl, G., Adolfsson, P., Borjesson, M., Mannheimer, C., & Rodger, S. (2005). Iron deficiency and anemia: a common problem in female elite soccer players. *International Journal of Sport Nutrition and Exercise Metabolism*, 15(6), 689.
- Magalhães, J., Rebelo, A., Oliveira, E., Silva, J. R., Marques, F., & Ascensão, A. (2010). Impact of Loughborough Intermittent Shuttle Test versus soccer match on physiological, biochemical and neuromuscular parameters. *European Journal of Applied Physiology*, 108(1), 39-48.
- Malcovati, L., Pascutto, C., & Cazzola, M. (2003). Hematologic passport for athletes competing in endurance sports: a feasibility study. *Haematologica*, 88(5), 570-581.
- Meneghini, R. (1997). Iron homeostasis, oxidative stress, and DNA damage. *Free Radical Biology and Medicine*, 23(5), 783-792.
- Mercer, K. W., & Densmore, J. J. (2005). Hematologic disorders in the athlete. *Clinics in Sports Medicine*, 24(3), 599-621.
- Orino, K., Lehman, L., Tsuji, Y., Ayaki, H., Torti, S. V., & Torti, F. M. (2001). Ferritin and the response to oxidative stress. *Biochemical Journal*, 357(1), 241-247.
- Ostojic, S., & Ahmetovic, Z. (2009). Indicators of iron status in elite soccer players during the sports season. *International Journal of Laboratory Hematology*, 31(4), 447-452.
- Peeling, P., Dawson, B., Goodman, C., Landers, G., & Trinder, D. (2008). Athletic induced iron deficiency: new insights into the role of inflammation, cytokines and hormones. *European Journal of Applied Physiology*, 103(4), 381-391.
- Ramel, A., Wagner, K.-H., & Elmadfa, I. (2003). Acute impact of submaximal resistance exercise on immunological and hormonal parameters in young men. *Journal of Sports Science*, 21(12), 1001-1008.

- Rampinini, E., Coutts, A., Castagna, C., Sassi, R., & Impellizzeri, F. (2007). Variation in top level soccer match performance. *International Journal of Sports Medicine*, 28(12), 1018-1024.
- Resina, A., Gatteschi, L., Giamberardino, M., Imreh, F., Rubenni, M., & Vecchiet, L. (1991). Hematological comparison of iron status in trained top-level soccer players and control subjects. *International Journal of Sports Medicine*, 12(5), 453-456.
- Rietjens, G., Kuipers, H., Hartgens, F., & Keizer, H. (2002). Red blood cell profile of elite olympic distance triathletes. A three-year follow-up. *International Journal of Sports Medicine*, 23(6), 391-396.
- Robach, P., Boisson, R. C., Vincent, L., Lundby, C., Moutereau, S., Gergel , L., et al. (2014). Hemolysis induced by an extreme mountain ultra-marathon is not associated with a decrease in total red blood cell volume. *Scandinavian Journal of Medicine and Science in Sports*, 24(1), 18-27.
- Rodrigues, V., Mortimer, L., Condessa, L., Coelho, D., Soares, D., & Garcia, E. (2007). Exercise intensity in training sessions and official games in soccer. *Journal of Sports Science & Medicine*, 1, 57-61.
- Schumacher, Y. O., Jankovits, R., B ltermann, D., Schmid, A., & Berg, A. (2002). Hematological indices in elite cyclists. *Scandinavian Journal of Medicine and Science in Sports*, 12(5), 301-308.
- Silva, A., Santhiago, V., Papoti, M., & Gobatto, C. (2008). Psychological, biochemical and physiological responses of Brazilian soccer players during a training program. *Science & Sports*, 23(2), 66-72.
- Spiropoulos, K., & Trakada, G. (2003). Hematologic and biochemical laboratory parameters before and after a marathon race. *Lung*, 181(2), 89-95.
- Tan, D., Dawson, B., & Peeling, P. (2012). Hemolytic effects of a football-specific training session in elite female players. *International Journal of Sports Physiology and Performance*, 7, 271-276.
- Thompson, D., Williams, C., Garcia-Roves, P., McGregor, S., McArdle, F., & Jackson, M. (2003). Post-exercise vitamin C supplementation and recovery from demanding exercise. *European Journal of Applied Physiology*, 89(3-4), 393-400.

## HEMATOLOŠKI PROFIL I STATUS GVOŽDA NAKON FUDBALSKOG MEĀA

Cilj ovog istraŹivanja je bio da se provjeri hematološki odgovor i koncentracija gvoŹda kod hrvatskih U-21 fudbalera nakon zvaniĉne fudbalske utakmice. Grupa od 19 obuĉenih, zdravih muškaraca fudbalera iz selekcije U-21 Hrvatskog fudbalskog saveza je uĉestvovalo u ovoj studiji. Analiza koncentracije crvenih krvnih zrnaca (RBC), hemoglobin (Hb), hematokrit (Hct), srednja vrednost zapremine eritrocita (MCV), proseĉna koliĉina hemoglobina u eritrocitu (MCH), srednja koncentracija hemoglobina u eritrocitima (MCHC), i trombocita je izvedena u cilju otkrivanja hematološkog statusa. Pored toga, gvoŹde, ukupni kapacitet vezivanja gvoŹda u serumu (TIBC), nezasiĉeni kapacitet vezivanja gvoŹda (UIBC) i analiza feritina dobijeni su odmah posle fudbalske utakmice. Znaĉajan pad Hct (%) uoĉen je odmah posle utakmice. Međutim, nije bilo znaĉajnih promena u nivou hemoglobina. Koncentracija gvoŹda znaĉajno je poveĉana ( $p < 0,01$ ) posle fudbalske utakmice. Nivoi Feritina su takođe veći na kraju utakmice u odnosu na polazne mere. MoŹe se zakljuĉiti da je fudbalski meĀ izazvao poveĉanje gvoŹda odmah nakon utakmice što je u suprotnosti sa nedavnim studijama. Ukratko, ova studija pruŹa dragocenu priliku za izuĉavanje biohemijskih parametara koji bi mogli dovesti do nekih bolesti, ili povreda tokom i nakon igre.

Ključne reĉi: biohemijske analize, fudbal, profesionalni igraĉi, zamor